


**Potential Wildlife-Highway Interactions at Select Mile-Markers
Along Montana Department of Transportation Routes:
An ArcView GIS Analysis**



Submitted by:

**Lynn R. Irby
Kevin M. Podruzny
Fish and Wildlife Management Program
Department of Ecology
Montana State University
Bozeman, MT 59717**

Submitted to:

**MONTANA DEPARTMENT OF TRANSPORTATION
RESEARCH MANAGEMENT UNIT
2701 PROSPECT AVENUE
HELENA, MT 59620**

(February 2001)

CONTENTS

**Combining MDT Roads Information with
GAP Predicted Species Distributions**

Acknowledgements

GIS Process

Data Projection Conversion for MDT Data Layers

Figures

Montana Natural Heritage Program Species of Special Concern

List of Montana Vertebrate Species with Predicted GAP Distributions

**Predicted Distributions for Species of Special Concern and Watch List
Species and Potential Conflicts with Montana Highways**

Combining MDT Roads Information with GAP Predicted Species Distributions

A potentially useful tool for MDT biologists and planners is a database table indicating where highways throughout the state show the potential for conflict with various vertebrate species. Given a list of potential species at a specified location, MDT biologists can be alert to the possibility of conflicts and/or mitigation. To develop a potential conflict database, areas around mile markers first defined various Montana road sections. Around each mile marker, we created a 10-mile radius buffer. 10-mile buffers have 2 major benefits: 1) they extend completely to the furthest neighboring mile marker, allowing for more complete coverage of roads between mile markers; 2) they are conservative in predicting potential species occurrences by indicating when mile markers fall within species' general distributions (i.e., they are less likely to miss rarer species). These buffered mile markers were then merged with GAP predicted species distributions via GIS.

GAP data provided the series of predicted species distributions based on habitat availability throughout the state. The National GAP Program supplied Montana GAP data layers for terrestrial vertebrates. GAP data will likely be the most widespread and available database of predicted species distributions available to the greatest number of groups and organizations and managers, especially for little studied species.

Predicted species distributions intersected with MDT mile marker layers were constructed for Montana Natural Heritage Program species of special concern or watch list species. Similarly, predicted species distributions intersected with MDT mile marker layers were constructed for all mammalian species. Output in the form of a MS Excel spreadsheet, an ASCII text file table, and an interactive ArcView data layer were constructed for species of special concern and watch list species. An interactive ArcView data layer was also constructed for all mammalian species.

It should be noted that the resulting conflict databases only provide a list of **potential** species that could occur at or near various locations. Like the GAP data from which they are produced, mile markers that indicate a species do not absolutely confirm the presence or absence of that species, they simply indicated that the species might occur near the point based on availability of appropriate habitat. MDT biologists should closely examine identified sites for all indicated species to determine if specific local habitats could indeed support those species.

Acknowledgements

We thank the following individuals and organizations for providing GIS layers for GAP analysis: Ree Bannon and Melissa Hart with the National GAP Program, and John O'Mara with Montana Department of Transportation. We also thank Shannon Podruzny, Edward Olexa, and Peter Gogan with the USGS Northern Rocky Mountain Science Center for GIS assistance and facilities.

GIS Process

To perform the following GIS processes in ArcView, the following extensions must first be installed: Animal Movement SA v 2.04 beta; Buffer Theme Builder Extension; Database Access; Geoprocessing; Getgrid Value Extension v2; Graticules and Measured Grids; Model Builder; and Spatial Analyst

- 1) Create/Open ArcView Project (wildlife.apr in folder Irby_MDT).
- 2) Re-project MDT routes, mile-markers, counties, cities, and streams themes to conform with GAP data layer specifications (See Data Projection Conversion Section below). This step has been performed and does not need to be repeated.
- 3) Add above re-projected themes to Project wildlife.apr.
- 4) Construct 10-mile radius buffers around mile-markers (Figure 1). This layer has been created (Buffered Mile Markers theme in wildlife.apr) and will not need to be duplicated unless a buffer of a different radius is desired.
- 5) Add Theme (data source=GRID data source) for desired species from CD or hard drive.
- 6) Convert GAP predicted species distributions from GRID format to Shapefile format (Figure 1). Activate species GRID layer created in step 5. Click Theme menu – Convert to Shapefile. Enter filename in an appropriate directory. Click <OK>. (This step can be time consuming ... wait).. At prompt “Add Shapefile as Theme to View?”, click <Yes>.
- 7) Intersect Shapefiles of buffered mile markers and predicted species distributions (Figure 1). Activate GRID theme for desired species. Click <Theme Properties>. Click <Theme Query Builder>. Enter “Gridcode=1”. Click <OK> and then click <OK> again to run Query and exit Theme Properties. Activate “Buffered Mile Markers” theme. Click Theme menu – Select by Theme. Then choose “Select Features that Intersect the selected features of (Shapefile of species distribution)” and click <New Set>.
- 8) Develop database table of potential species occurrence for each mile marker. Click <Open Theme Table> of activated “Buffered Mile Markers” theme. Click Table menu – Start Editing. Click Edit Menu – Add Field. Enter Field Name (Species ID etc. up to 8 characters) and Field Width. Click <OK> to exit Field Definitions. Highlight Field Name. Click <Calculate>. Enter “fieldname” = 1. Click <Switch Selection>. Click <Calculate>. Enter “fieldname” = 0. Click <Clear Selected Features>. Click Table menu – Stop Editing. Click <Yes> to Save Edits. Close table. Delete Grid and Shapefiles for species distribution from view. Click <Save>. Repeat steps for additional species. To make “Buffered Mile Markers” theme table available to use in various database managing software environments (e.g., Access, Oracle, dBase) - open the table and choose File menu – Export and save as a dBase file (with a new name).
- 9) Import database table as Shapefile to allow for interactive onscreen query of mile markers (Figure 2). Go to “Tables” in wildlife.apr, click <Add> and choose the new dBase file exported in step 8. Go to View window, Choose

View menu – Add Event Theme. Select the new dBase file from the list of tables and use the UTM fields for the X and Y coordinates. Long species names can be added to field names in the new theme's attribute table. With the table open, choose Table menu – Properties to enter aliases. In the View window with the new theme activated, use the <Identity> tool to view attributes of a particular mile marker.

Data Projection Conversion for MDT Data Layers

Input Coordinate System:

Name: NAD_1983_Montana
POSC: 32100
Unit: Meter
Geographic CSYS: GCS_North_American_1983
Datum: D_North_American_1983
Prime Meridian: Greenwich
False Easting: 600000
False Northing: 0
Base Projection: Lambert_Conformal_Conic
Central_Meridian: -109.5
Central_Parallel: 44.25
Standard_Parallel_1: 45.0
Standard_Parallel_2: 49.0

Input Geographic Transformation:

none


Output Geographic Transformation:

WGS_1984_4_To_NAD_1927 [8073]



Output Coordinate System:

Name: Custom
POSC: -1
Unit: Meter
Geographic CSYS: GCS_North_American_1927
Datum: D_North_American_1927
Prime Meridian: Greenwich
False Easting: 600000
False Northing: 0
Base Projection: Albers
Central_Meridian: -109.5
Central_Parallel: 44.25
Standard_Parallel_1: 46.0
Standard_Parallel_2: 48.0



Figures




**ArcView Shape File of
Mile Markers along MDT Routes**





**ArcView Shape File of
Mile Markers with 10 mi Buffers**




**Intersection of Buffered Mile Markers and
Predicted Species Distribution**




**ArcView Grid File of
Predicted Species Distribution**



**ArcView Shape File of
Predicted Species Distribution**





Shape	Point
Buff id	1558
Route	P00099N
Milemarker	55
X_coord	722055
Y_coord	529133
Richness	46
Maba0101	0
Maba0103	0
Maba0107	0
Maba0108	0
Maba0113	0
Maba0115	0
Maba0123	0
Maba0125	0
Maba0128	1
Macc0101	1
Macc0102	0
Macc0107	1
Macc0109	0
Macc0111	1
Macc0112	0
Macc0114	1
Macc0115	0
Macc0201	1
Macc0401	1
Macc0503	1
Macc0701	0
Macc0801	1
Macc1001	0
Maja0101	1
Maja0103	0
Maja0301	1
Maja0303	1
Majb0101	0
Majb0102	0
Maje0201	1
Majf0101	0
Majf0102	0

Montana Natural Heritage Program Species of Special Concern



P.O. Box 201800
1515 East Sixth Avenue
Helena, Montana 59620-1800
406-444-3009
mtnhp@state.mt.us

Animal Species of Special Concern

September, 1999

Introduction

The Montana Natural Heritage Program (MTNHP) maintains an inventory of the elements of biological diversity in Montana. The inventory focuses on plant species, animal species, plant communities, and biological features that are rare, endemic, disjunct, threatened, or endangered throughout their range or in Montana, vulnerable to extirpation from Montana, or in need of further research.

The attached list of animal species of special concern in Montana is used to direct data acquisition by MTNHP, and to provide information to others on the current status of these species. It has been developed largely from information in the scientific literature, unpublished reports, agency databases, field research, and field inventories. This information comes from a variety of cooperating local, state and federal agencies, private organizations and businesses, academic researchers, and interested individuals.

Background information was obtained from sources such as *Vertebrate Species of Special Interest or Concern* (Flath 1984, 1995), *P. D. Skaar's Montana Bird Distribution* (Montana Bird Distribution Committee 1996), *Fishes of Montana* (Brown 1971), the USFWS's *Animal Candidate Review for Listing as Endangered or Threatened Species* (61 FR 7596, Feb. 28, 1996), and from scientific literature, personal contacts, museum specimens, and MTNHP staff research.

This list is dynamic and is periodically revised as new data becomes available. Changes to the list since the previous edition (Reichel 1997) are highlighted for quick reference. Printed revisions are produced annually, while regularly-updated versions are available from our office or via our home page at www.nris.state.mt.gov/mtnhp. Comments and suggestions are continually needed on this list, particularly regarding recommended additions or deletions. We also welcome new or updated location information for any of these taxa.

Heritage Program Ranks

Taxa are evaluated and ranked by the Heritage Program on the basis of their global (range-wide) status, and their state-wide status. These ranks are used to determine protection and data collection priorities, and are revised as new information becomes available.

A scale of 1 (critically imperiled) to 5 (demonstrably secure) is used for these ranks. Each species is assigned the appropriate combination of global (G) and state (S) ranks.

Example: Merriam's shrew = G5 / S3 (i.e., species is demonstrably secure globally; in Montana is found within a restricted range).

Global and state ranks are assigned according to a standardized procedure used by all Natural Heritage Programs (The Nature Conservancy 1992), and are briefly defined below.

Rank	Definition
G1 S1	Critically imperiled because of extreme rarity (5 or fewer occurrences, or very few remaining individuals), or because of some factor of its biology making it especially vulnerable to extinction.
G2 S2	Imperiled because of rarity (6 to 20 occurrences), or because of other factors demonstrably making it very vulnerable to extinction throughout its range.
G3 S3	Either very rare and local throughout its range, or found locally (even abundantly at some of its locations) in a restricted range, or vulnerable to extinction throughout its range because of other factors; in the range of 21 to 100 occurrences.
G4 S4	Apparently secure, though it may be quite rare in parts of its range, especially at the periphery.
G5 S5	Demonstrably secure, though it may be quite rare in parts of its range, especially at the periphery.
GU SU	Possibly in peril, but status uncertain; more information needed.
GH SH	Historically known; may be rediscovered.
GX SX	Believed to be extinct; historical records only, continue search.

Other Heritage Codes

A	Accidental in the state; including species (usually birds or butterflies) recorded very infrequently, hundreds or thousands of miles outside their usual range.
B	A state rank modifier indicating breeding status for a migratory species. Example: S1B,SZN - breeding occurrences for the species are ranked S1 (critically imperiled) in the state, nonbreeding occurrences are not ranked in the state.

E	An exotic established in the state; may be native in nearby regions.
HYB	Element represents a hybrid of species
N	A state rank modifier indicating non-breeding status for a migratory species. Example: S1B,SZN - breeding occurrences for the species are ranked S1 (critically imperiled) in the state, nonbreeding occurrences are not ranked in the state.
P	Indicates the element may potentially occur in the state.
Q	Taxonomic questions or problems involved, more information needed; appended to the global rank.
R	Reported in the state; but lacking documentation which would provide a basis for either accepting or rejecting the report.
T	Rank for subspecific taxon (subspecies, variety, or population); appended to the global rank for the full species.
Z	Ranking not applicable.
C	Captive or Cultivated: extant only in captivity or cultivation, or as a reintroduced population not yet established.

Fish and Wildlife Service Status

The abbreviations in this column denote the categories defined in the U.S. Fish and Wildlife Service Notice of Review (1980, 1983, 1985, 1990, 1993, 1996) and indicate the status of a taxon in Montana under the federal Endangered Species Act of 1973 (16 U.S.C.A. § 1531-1543 (Supp. 1996)). Categories are listed below:

LE	listed endangered
LT	listed threatened
PE	proposed endangered
PT	proposed threatened
C	candidate--substantial information exists in U.S. Fish and Wildlife Service files on biological vulnerability to support proposals to list as endangered or threatened.
XN	non-essential experimental population

Forest Service Status

This column indicates the status of species on Forest Service lands as defined by the U.S. Forest Service manual (2670.22). These taxa are listed as such by the Regional Forester (Northern Region) on National Forests in Montana. The Forest Service lists species as either:

E	edangered: federally listed as endangered
T	threatened: federally listed as threatened
PT	proposed threatened: federally-proposed as threatened
S	sensitive: animal species identified by the Regional Forester for which population viability is a concern as evidenced by significant downward trend in population or a significant downward trend in habitat capacity.

Bureau of Land Management Status

This column indicates the status of species on Bureau of Land Management lands as defined by the BLM 6840 Manual.

SS Special Status: federally-listed Endangered, Threatened, or Candidate species or other rare or endemic species that occur on BLM lands.

Changes from the June, 1997 List

The Eastern Red Bat has been added to the Species of Special Concern List after first-time observations were reported for Montana. The Boreal Toad and Northern Leopard Frog have been moved from the Watch List to the Species of Special Concern List following recent marked declines in numbers reported. And the Wood Frog has been moved from the Species of Special Concern list to the Watch List due to a lack of confirmed observations of the species in Montana.

The Deepwater Sculpin has been added to the Watch List as potentially occurring in Montana following a report of the species being collected in Waterton Lake on the Canadian side. The Short-horned Lizard and Common Garter Snake have been added to the watch list due to the few numbers reported.

Global and state ranks have changed for a number of species since the 1997 list. These changes are highlighted in the following pages.

Selected References

- Brown, C. J. D. 1971. Fishes of Montana. Montana State University, Bozeman. 207 pp.
- Flath, D. L. 1984. Vertebrate species of special interest or concern. Montana Department of Fish, Wildlife and Parks. Helena. 76 pp.
- Flath, D. L. 1995. Species of special interest or concern. [Species list]. 7 pp. Montana Department of Fish, Wildlife and Parks, Helena, Montana.
- Frest, T. J. and E. J. Johannes. 1995. Interior Columbia Basin mollusk species of special concern. Final report to the Interior Columbia Basin Ecosystem Management Project, Walla Walla, WA. Contract #43-0E00-4-9112. 274 pp. plus appendices.
- Holton, G. D., and H. E. Johnson. 1996. A field guide to Montana fishes. Second Edition. Mont. Dept. Fish, Wildl. Parks, Helena. 104 pp.

- Miller, K. B. and D. L. Gustafson. 1996. Distribution of the Odonata of Montana. *Bulletin of American Odonatology* 3(4):75-88.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana bird distribution. Fifth edition. Special Publication No. 3, Montana Natural Heritage Program, Helena. 129 pp.
- Reichel, J. D. 1997. Montana animal species of special concern. [Unpublished list.] Montana Natural Heritage Program, Helena. 9 pp.
- Reichel, J. D. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34. The Nature Conservancy, Conservation Science Division, in association with the Network of Natural Heritage Programs and Conservation Data Centers. October 1992. Biological and Conservation Data System (Supplement 2+, released March, 1994). Arlington, Virginia.
- Thompson, L. S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Montana Audubon Council, Helena. 24 pp.
- U.S. Fish and Wildlife Service. 1996. Animal Candidate Review for Listing as Endangered and Threatened Species. 61 FR 7596, February 28, 1996.
- U.S. Fish and Wildlife Service. 1993. Endangered and threatened wildlife and plants. 50 CFR 17.11 & 17.12, August 23, 1993. U. S. Govt. Printing Office: 1993-342-477/94033. 40 pp.

More Information

Extensive additional information, including location data, is available on all species listed here. MTNHP serves as an information clearinghouse, and each year responds to hundreds of data requests regarding plants, animals, and biological communities of special concern.

To place a data request, contact us via phone (406-444-3009), email (mtnhp@state.mt.us), or use our on-line request form (www.nris.state.mt.us/mtnhp).

MTNHP's website provides direct access to species data, photographs, search tools, and links to related sites. Visit us at www.nris.state.mt.us/mtnhp.

The preferred citation for this document is:

Roedel, M.D. 1999. Montana animal species of special concern. [Unpublished list.] Montana Natural Heritage Program, Helena. 8 pp.

Watch List Species

These species are not currently known to meet the criteria of the Species of Special Concern. They are on the Watch List for one or more of the following reasons:

- (1) there are indications that they may be less common than currently thought;
- (2) they are currently declining in Montana or across much of their range;
- (3) there is so little information available that we cannot adequately rank them; and

(4) they are not known to have established breeding populations in Montana, but are known from states or provinces near their border with Montana and if they occur in Montana they would be Species of Special Concern.

While not actively soliciting information for these species, we are accumulating data as it is received for possible re-ranking of these species. Modifications to global and state ranks are the result of changes in either our knowledge of the species or actual changes in population status.

Fish

Myoxocephalus thompsoni
Deepwater Sculpin
Hybognathus argyritis
Western Silvery
Minnow
Hybognathus placitus
Plains Minnow
Hybopsis gracilis
Flathead Chub

Amphibians

Ascaphus truei
Tailed Frog
Bufo cognatus
Great Plains Toad
Spea bombifrons
Plains Spadefoot
Spea intermontana
Great Basin Spadefoot
Rana sylvatica
Wood Frog

Reptiles

Phrynosoma hernandesi
Short-horned Lizard
Sceloporus graciosus
Sagebrush Lizard
Eumeces skiltonianus
Western Skink
Thamnophis sirtalis
Common Garter Snake

Birds

Ardea herodias
Great Blue Heron
Lagopus leucurus
White-tailed Ptarmigan
Centrocercus urophasianus
Sage Grouse
Tyto alba
Barn Owl
Otus asio
Eastern Screech-owl
Otus kennicottii
Western Screech-owl
Surnia ulula
Northern Hawk Owl
Selasphorus platycercus
Broad-tailed Hummingbird
Lanius ludovicianus
Loggerhead Shrike
Vermivora peregrina
Tennessee Warbler
Mniotilta varia
Black-and-white Warbler
Amphispiza belli
Sage Sparrow

Mammals

Myotis yumanensis
Yuma Myotis
Sylvilagus floridanus
Eastern Cottontail
Spilogale putorius
Eastern Spotted Skunk
Spilogale gracilis
Western Spotted Skunk

Insects

Tanypteryx hageni
Black Petaltail
Cordulegaster dorsalis
Pacific Spiketail
Gomphus externus
Plains Clubtail
Ophiogomphus occidentis
Sinuous Snaketail
Aeshna canadensis
Canada Darner
Aeshna constricta
Lance-tipped Darner
Aeshna multicolor
Blue-eyed Darner
Aeshna sitchensis
Zigzag Darner
Aeshna tuberculifera
Black-tipped Darner
Macromia illinoiensis
Illinois River Cruiser
Somatochlora minor
Ocellated Emerald

Libellula composita
Bleached Skimmer
Libellula nodisticta
Hoary Skimmer
Pantala flavescens
Wandering Glider
Sympetrum madidum
Red-veined Meadowhawk
Lestes forcipatus
Sweetflag Spreadwing
Argia fumipennis violacea
Variable Dancer
Argia alberta
Paiute Dancer
Coenagrion angulatum
Prairie Bluet
Enallagma civile
Familiar Bluet
Enallagma clausum
Alkali Bluet
Argiomphus cornutus
Horned Clubtail

Mollusks

Radiodiscus abietum
Fir Pinwheel
Polygyrella polygyrella
Humped Coin
Udosarx lyrata lyrata
Lyre Mantleslug
Protenetus exacuus megas
Prairie Sprite

Species of Special Concern

Fish

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Cottus confusus</i>	Shorthead Sculpin	G5	S3			
<i>Cottus rhotheus</i>	Torrent Sculpin	G5	S2		S	
<i>Cottus ricei</i>	Spoonhead Sculpin	G5	S1			
<i>Acipenser transmontanus pop 1</i>	White Sturgeon (Kootenai River Population)	G4T1Q	S1	LE	E	
<i>Scaphirhynchus albus</i>	Pallid Sturgeon	G1G2	S1	LE		SS
<i>Polyodon spathula</i>	Paddlefish	G4	S1S2			SS
<i>Lepisosteus platostomus</i>	Shortnose Gar	G5	S1			SS
<i>Oncorhynchus clarki bouvieri</i>	Yellowstone Cutthroat Trout	G4T2	S2		S	SS
<i>Oncorhynchus clarki lewisi</i>	Westslope Cutthroat Trout	G4T3	S3		S	SS
<i>Oncorhynchus mykiss gairdneri</i>	Interior Redband Trout	S3	S2		S	
<i>Salvelinus confluentus</i>	Bull Trout	G3	S3	LT	T	SS
<i>Thymallus arcticus montanus</i>	Montana Arctic Grayling	G5T2Q	S1	C	S	SS
<i>Phoxinus eos x phoxinus neogaeus</i>	Northern Redbelly X Finescale Dace	HYB	S3			SS
<i>Hybopsis gelida</i>	Sturgeon Chub	G2	S2	C		SS
<i>Hybopsis meeki</i>	Sicklefin Chub	G3	S1	C		SS
<i>Semotilus margarita</i>	Pearl Dace	G5	S2			SS
<i>Cycleptus elongatus</i>	Blue Sucker	G4	S3?			SS
<i>Percopsis omiscomaycus</i>	Trout-perch	G5	S1			

Amphibians

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Plethodon idahoensis</i>	Coeur d'Alene Salamander	G3	S2		S	SS
<i>Dicamptodon aterrimus</i>	Idaho Giant Salamander	G3	S4			
<i>Bufo boreas</i>	Boreal Toad	G4	S3S4		S	
<i>Bufo hemiophrys</i>	Canadian Toad	G4	S1			SS
<i>Rana pipiens</i>	Northern Leopard Frog	G5	S3S4		S	

Reptiles

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Chelydra serpentina</i>	Snapping Turtle	G5	S3			SS
<i>Trionyx spiniferus</i>	Spiny Softshell	G5	S3			SS
<i>Heterodon nasicus</i>	Western Hognose Snake	G5	S			
<i>Lampropeltis triangulum</i>	Milk Snake	G5	S2			
<i>Liochlorophis vernalis</i>	Smooth Green Snake	G5	S2S3			

Birds

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Gavia immer</i>	Common Loon	G5	S2B,SZN		S	SS
<i>Aechmophorus clarkii</i>	Clark's Grebe	G5	S2S4B,SZN			

Birds (continued)

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Pelecanus erythrorhynchos</i>	American White Pelican	G3	S2B,SZN			
<i>Nycticorax nycticorax</i>	Black-crowned Night-heron	G5	S2S3B,SZN			
<i>Cygnus buccinator</i>	Trumpeter Swan	G4	S2B,S2N		S	SS
<i>Histrionicus histrionicus</i>	Harlequin Duck	G4	S2B,SZN		S	SS
<i>Haliaeetus leucocephalus</i>	Bald Eagle	G4	S3B,S3N	LT	T	SS
<i>Accipiter gentilis</i>	Northern Goshawk	G5	S3S4		S	SS
<i>Buteo regalis</i>	Ferruginous Hawk	G4	S3B,SZN			SS
<i>Falco peregrinus</i>	Peregrine Falcon	G4	S1S2B,SZN		pending	SS
<i>Tympanuchus phasianellus columbianus</i>	Columbian Sharp-tailed Grouse	G4T3	S1		S	SS
<i>Coturnicops noveboracensis</i>	Yellow Rail	G4	S1B,SZN			
<i>Grus americana</i>	Whooping Crane	G1	SZN	LE	E	SS
<i>Charadrius melodus</i>	Piping Plover	G3	S2B,SZN	LT		SS
<i>Charadrius montanus</i>	Mountain Plover	G2	S2B,SZN	PT	PT	SS
<i>Himantopus mexicanus</i>	Black-necked Stilt	G5	S2B,SZN			
<i>Larus pipixcan</i>	Franklin's Gull	G4G5	S3B,SZN			
<i>Sterna caspia</i>	Caspian Tern	G5	S2B,SZN			
<i>Sterna hirundo</i>	Common Tern	G5	S3B,SZN			
<i>Sterna forsteri</i>	Forster's Tern	G5	S2B,SZN			
<i>Sterna antillarum athalassos</i>	Interior Least Tern	G4T2Q	S1B,SZN	LE		SS
<i>Chlidonias niger</i>	Black Tern	G4	S3B,SZN			SS
<i>Coccyzus americanus</i>	Yellow-billed Cuckoo	G5	S3B,SZN			
<i>Otus flammeolus</i>	Flammulated Owl	G4	S3B,SZN		S	SS
<i>Athene cunicularia</i>	Burrowing Owl	G4	S3B,SZN		S	SS
<i>Strix nebulosa</i>	Great Gray Owl	G5	S3			SS
<i>Aegolius funereus</i>	Boreal Owl	G5	S4			SS
<i>Cypseloides niger</i>	Black Swift	G4	S3B,SZN			
<i>Picoides arcticus</i>	Black-backed Woodpecker	G5	S3		S	SS
<i>Empidonax alnorum</i>	Alder Flycatcher	G5	S1B,SZN			
<i>Tyrannus vociferans</i>	Cassin's Kingbird	G5	S1S3B,SZN			
<i>Poliophtila caerulea</i>	Blue-gray Gnatcatcher	G5	S1B,SAN			
<i>Spiza americana</i>	Dickcissel	G5	S1S2B,SZN			SS
<i>Ammodramus bairdii</i>	Baird's Sparrow	G4	S3S4B,SZN		S	
<i>Ammodramus leconteii</i>	Le Conte's Sparrow	G4	S1S2B,SZN			SS
<i>Ammodramus nelsoni</i>	Nelson's Sharp-tailed Sparrow	G5	S1B,SZN			

Mammals

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Sorex nanus</i>	Dwarf Shrew	G4	S3			
<i>Sorex merriami</i>	Merriam's Shrew	G5	S3			SS
<i>Myotis thysanodes</i>	Fringed Myotis	G5	S3			
<i>Myotis septentrionalis</i>	Northern Myotis	G4	S1			
<i>Lasturus borealis</i>	Eastern Red Bat	G5	S1			

Mammals (continued)

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Euderma maculatum</i>	Spotted Bat	G4	S1		S	SS
<i>Corynorhinus townsendii</i>	Townsend's Big-eared Bat	G4	S2S3		S	SS
<i>Lepus californicus</i>	Black-tailed Jack Rabbit	G5	S2S3			
<i>Brachylagus idahoensis</i>	Pygmy Rabbit	G4	S2S3		S	SS
<i>Tamias umbrinus</i>	Uinta Chipmunk	G5	S3?			
<i>Cynomys ludovicianus</i>	Black-tailed Prairie Dog	G4	S3S4		S	SS
<i>Cynomys leucurus</i>	White-tailed Prairie Dog	G4			S	SS
<i>Perognathus parvus</i>	Great Basin Pocket Mouse	G5	S2S4			
<i>Chaetodipus hispidus</i>	Hispid Pocket Mouse	G5	S1			
<i>Synaptomys borealis</i>	Northern Bog Lemming	G4	S2		S	SS
<i>Zapus hudsonius</i>	Meadow Jumping Mouse	G5	S2S3			SS
<i>Canis lupus</i>	Gray Wolf	G4	S1	LE	E	SS
<i>Vulpes velox</i>	Swift Fox	G3	S1	C	S	SS
<i>Ursus arctos horribilis</i>	Grizzly Bear	G4T3	S1S2	LT	T	SS
<i>Martes pennanti</i>	Fisher	G5	S2		S	SS
<i>Mustela nigripes</i>	Black-footed Ferret	G1	SH#	E	E	SS
<i>Gulo gulo luscus</i>	North American Wolverine	G5T4	S2		S	SS
<i>Lynx canadensis</i>	Lynx (US Lower 48 Population)	G5T?Q	S2	PT	PT	SS
<i>Rangifer tarandus caribou</i>	Woodland Caribou	G5T4	SX	LE	E	

Mollusks

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Discus brunsoni</i>	Mission Range Disc	G1	S1			
<i>Hemphillia danielsi</i>	Marbled Jumping-slug	G1G3	S1S3			
<i>Magnipelta mycophaga</i>	Spotted Slug	G2G3	S1S3			
<i>Zacoleus idahoensis</i>	Sheathed Slug	G3G4	S2S3			
<i>Oreohelix alpina</i>	Alpine Mountainsnail	G1	S1			
<i>Oreohelix amariradix</i>	Bitterroot Mountainsnail	G1G2	S1S2			
<i>Oreohelix carinifera</i>	Keeled Mountainsnail	G1	S1			
<i>Oreohelix elrodi</i>	Carinate Mountainsnail	G1	S1			
<i>Oreohelix strigosa berryi</i>	Berry's Mountainsnail	G5T2	S1S2			
<i>Oreohelix yavapai mariae</i>	Gallatin Mountainsnail	G4?T1	S1			
<i>Oreohelix sp 3</i>	Bearmouth Mountainsnail	G1G2	S1S2			
<i>Oreohelix sp 4</i>	Drummond Mountainsnail	G1	S1			
<i>Oreohelix sp 5</i>	Brunson Mountainsnail	G1G2	S1S2			
<i>Oreohelix sp 6</i>	Kintla Lake Mountainsnail	G1	S1			
<i>Oreohelix sp 7</i>	Kitchen Creek Mountainsnail	G1G2	S1S2			
<i>Oreohelix sp 10</i>	Missoula Mountainsnail	G1G3	S1S3			
<i>Oreohelix sp 11</i>	Subcarinate Mountainsnail	G1	S1			
<i>Oreohelix sp 31</i>	Byrne Resort Mountainsnail	G1G2	S1S2			
<i>Udosarx lyrata russelli</i>	Russell Mantleslug	G1	S1			
<i>Lyogyrus greggi</i>	Rocky Mountain Dusksnail	G3G4	S1			
<i>Amnicola sp 2</i>	Washington Dusksnail	G1	S1			

Mollusks (continued)

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Fluminicola fuscus</i>	Columbia Pebblesnail	G2G3	SX			
<i>Acroloxus coloradensis</i>	Rocky Mountain Capshell	G?	S1			
<i>Stagnicola elrodianus</i>	Largemouth Pondsail	G1	S1			
<i>Stagnicola montanensis</i>	Mountain Marshsnail	G3	S1S3			
<i>Fisherola nuttalli</i>	Shortface Lanx	G2?	S1S3			
<i>Physa megalochlamys</i>	Large-mantle Physa	G3	S1			

Insects

Scientific Name	Common Name	Global Rank	State Rank	ESA	USFS	BLM
<i>Microcylloepus browni</i>	Brown's Microcylloepus Riffle Beetle	G1	S1			
<i>Caenis youngi</i>	A Mayfly	G3	S2			
<i>Phyciodes batesii</i>	Tawny Crescent	G4	S2S3			
<i>Euphydrys gillettii</i>	Gillette's Checkerspot	G3	S3			
<i>Erpetogomphus designatus</i>	Eastern Ringtail	G5	S1			
<i>Aeshna subarctica</i>	Subarctic Darner	G5	S1S2			
<i>Somatochlora albicincta</i>	Ringed Emerald	G5	S1S3			
<i>Somatochlora walshii</i>	Brush-tipped Emerald	G5	S1S2			
<i>Erythemis collocata</i>	Western Pondhawk	G5	S1S2			
<i>Leucorrhinia borealis</i>	Boreal Whiteface	G5	S1			
<i>Coenagrion interrogatum</i>	Subarctic Bluet	G5	S1S2			
<i>Enallagma optimolocus</i>	Last Best Place Damselfly	G1G3G	S1S3			
<i>Isocapnia crinita</i>	A Stonefly	G4	S2			
<i>Isocapnia integra</i>	A Stonefly	G4	S2			
<i>Utacapnia columbiana</i>	A Stonefly	G4	S2			
<i>Lednia tumana</i>	Meltwater Lednian Stonefly	G1	S1			
<i>Zapada cordillera</i>	A Stonefly	G3	S2			
<i>Zapada glacier</i>	Western Glacier Stonefly	G3	S1			
<i>Isoperla petersoni</i>	A Stonefly	G4	S2?			
<i>Rhyacophila alexanderi</i>	Alexander's Rhyacophilan Caddisfly	G2	S2			
<i>Rhyacophila ebria</i>	A Caddisfly	G1?	S1?			
<i>Rhyacophila newelli</i>	A Rhyacophilan Caddisfly	G2?	S2?			
<i>Rhyacophila glaciali</i>	A Rhyacophilan Caddisfly	G1	S1			
<i>Agapetus montanus</i>	An Agapetus Caddisfly	G2?	S2?			

List of Montana Vertebrate Species with Predicted GAP Distributions

AAAA0108	AAAAA01080	LONG-TOED SALAMANDER	<i>Ambystoma macrodactylum</i>	
AAAA0114	AAAAA01140	TIGER SALAMANDER	<i>Ambystoma tigrinum</i>	
AAAD1227	AAAAD12270	COEUR D'ALENE SALAMANDER	<i>Plethodon idahoensis</i>	special concern
AAAH0103	AAAAH01030	IDAHO GIANT SALAMANDER	<i>Dicamptodon aterrimus</i>	special concern
AABA0101	AAABA01010	TAILED FROG	<i>Ascaphus truei</i>	watch list
AABB0103	AAABB01030	WESTERN TOAD	<i>Bufo boreas</i>	special concern
AABB0105	AAABB01050	GREAT PLAINS TOAD	<i>Bufo cognatus</i>	watch list
AABB0108	AAABB01080	CANADIAN TOAD	<i>Bufo hemiophrys</i>	special concern
AABB0118	AAABB01180	WOODHOUSE'S TOAD	<i>Bufo woodhousii</i>	
AABC0507	AAABC05070	WESTERN CHORUS FROG	<i>Pseudacris triseriata</i>	
AABC0510	AAABC05100	PACIFIC CHORUS FROG	<i>Pseudacris regilla</i>	
AABF0201	AAABF02010	PLAINS SPADEFOOT	<i>Spea bombifrons</i>	watch list
AABH0107	AAABH01070	BULLFROG	<i>Rana catesbeiana</i>	
AABH0117	AAABH01170	NORTHERN LEOPARD FROG	<i>Rana pipiens</i>	special concern
AABH0120	AAABH01200	WOOD FROG	<i>Rana sylvatica</i>	watch list
AABH0129	AAABH01290	COLUMBIA SPOTTED FROG	<i>Rana luteiventris</i>	
BNBA0103	ABNBA01030	COMMON LOON	<i>Gavia immer</i>	special concern
BNCA0201	ABNCA02010	PIED-BILLED GREBE	<i>Podilymbus podiceps</i>	
BNCA0301	ABNCA03010	HORNED GREBE	<i>Podiceps auritus</i>	
BNCA0302	ABNCA03020	RED-NECKED GREBE	<i>Podiceps grisegena</i>	
BNCA0303	ABNCA03030	EARED GREBE	<i>Podiceps nigricollis</i>	
BNCA0401	ABNCA04010	WESTERN GREBE	<i>Aechmophorus occidentalis</i>	
BNCA0402	ABNCA04020	CLARK'S GREBE	<i>Aechmophorus clarkii</i>	special concern
BNFC0101	ABNFC01010	AMERICAN WHITE PELICAN	<i>Pelecanus erythrorhynchos</i>	special concern
BNFD0102	ABNFD01020	DOUBLE-CRESTED CORMORANT	<i>Phalacrocorax auritus</i>	
BNGA0102	ABNGA01020	AMERICAN BITTERN	<i>Botaurus lentiginosus</i>	
BNGA0401	ABNGA04010	GREAT BLUE HERON	<i>Ardea herodias</i>	watch list
BNGA1101	ABNGA11010	BLACK-CROWNED NIGHT-HERON	<i>Nycticorax nycticorax</i>	special concern
BNGE0202	ABNGE02020	WHITE-FACED IBIS	<i>Plegadis chihi</i>	special concern
BNJB0201	ABNJB02010	TUNDRA SWAN	<i>Cygnus columbianus</i>	
BNJB0203	ABNJB02030	TRUMPETER SWAN	<i>Cygnus buccinator</i>	special concern
BNJB0304	ABNJB03040	GREATER WHITE-FRONTED GOOSE	<i>Anser albifrons</i>	
BNJB0401	ABNJB04010	SNOW GOOSE	<i>Chen caerulescens</i>	
BNJB0402	ABNJB04020	ROSS'S GOOSE	<i>Chen rossii</i>	
BNJB0503	ABNJB05030	CANADA GOOSE	<i>Branta canadensis</i>	
BNJB0901	ABNJB09010	WOOD DUCK	<i>Aix sponsa</i>	
BNJB1001	ABNJB10010	GREEN-WINGED TEAL	<i>Anas crecca</i>	
BNJB1006	ABNJB10060	MALLARD	<i>Anas platyrhynchos</i>	
BNJB1011	ABNJB10110	NORTHERN PINTAIL	<i>Anas acuta</i>	
BNJB1013	ABNJB10130	BLUE-WINGED TEAL	<i>Anas discors</i>	
BNJB1014	ABNJB10140	CINNAMON TEAL	<i>Anas cyanoptera</i>	
BNJB1015	ABNJB10150	NORTHERN SHOVELER	<i>Anas clypeata</i>	
BNJB1016	ABNJB10160	GADWALL	<i>Anas strepera</i>	
BNJB1018	ABNJB10180	AMERICAN WIGEON	<i>Anas americana</i>	
BNJB1102	ABNJB11020	CANVASBACK	<i>Aythya valisineria</i>	
BNJB1103	ABNJB11030	REDHEAD	<i>Aythya americana</i>	
BNJB1104	ABNJB11040	RING-NECKED DUCK	<i>Aythya collaris</i>	
BNJB1106	ABNJB11060	GREATER SCAUP	<i>Aythya marila</i>	
BNJB1107	ABNJB11070	LESSER SCAUP	<i>Aythya affinis</i>	
BNJB1501	ABNJB15010	HARLEQUIN DUCK	<i>Histrionicus histrionicus</i>	special concern
BNJB1801	ABNJB18010	COMMON GOLDENEYE	<i>Bucephala clangula</i>	
BNJB1802	ABNJB18020	BARROW'S GOLDENEYE	<i>Bucephala islandica</i>	
BNJB1803	ABNJB18030	BUFFLEHEAD	<i>Bucephala albeola</i>	
BNJB2001	ABNJB20010	HOODED MERGANSER	<i>Lophodytes cucullatus</i>	
BNJB2101	ABNJB21010	COMMON MERGANSER	<i>Mergus merganser</i>	
BNJB2102	ABNJB21020	RED-BREASTED MERGANSER	<i>Mergus serrator</i>	
BNJB2201	ABNJB22010	RUDDY DUCK	<i>Oxyura jamaicensis</i>	
BNKA0201	ABNKA02010	TURKEY VULTURE	<i>Cathartes aura</i>	
BNKC0101	ABNKC01010	OSPREY	<i>Pandion haliaetus</i>	
BNKC1001	ABNKC10010	BALD EAGLE	<i>Haliaeetus leucocephalus</i>	special concern
BNKC1101	ABNKC11010	NORTHERN HARRIER	<i>Circus cyaneus</i>	

BNKC1202	ABNKC12020	SHARP-SHINNED HAWK	Accipiter striatus	
BNKC1204	ABNKC12040	COOPER'S HAWK	Accipiter cooperii	
BNKC1206	ABNKC12060	NORTHERN GOSHAWK	Accipiter gentilis	special concern
BNKC1905	ABNKC19050	BROAD-WINGED HAWK	Buteo platypterus	
BNKC1907	ABNKC19070	SWAINSON'S HAWK	Buteo swainsoni	
BNKC1911	ABNKC19110	RED-TAILED HAWK	Buteo jamaicensis	
BNKC1912	ABNKC19120	FERRUGINOUS HAWK	Buteo regalis	special concern
BNKC1913	ABNKC19130	ROUGH-LEGGED HAWK	Buteo lagopus	
BNKC2201	ABNKC22010	GOLDEN EAGLE	Aquila chrysaetos	
BNKD0602	ABNKD06020	AMERICAN KESTREL	Falco sparverius	
BNKD0603	ABNKD06030	MERLIN	Falco columbarius	
BNKD0607	ABNKD06070	PEREGRINE FALCON	Falco peregrinus	special concern
BNKD0608	ABNKD06080	GYRFALCON	Falco rusticolus	
BNKD0609	ABNKD06090	PRAIRIE FALCON	Falco mexicanus	
BNLC0101	ABNLC01010	GRAY PARTRIDGE	Perdix perdix	
BNLC0301	ABNLC03010	CHUKAR	Alectoris chukar	
BNLC0701	ABNLC07010	RING-NECKED PHEASANT	Phasianus colchicus	
BNLC0901	ABNLC09010	SPRUCE GROUSE	Falcapennis canadensis	
BNLC0902	ABNLC09020	BLUE GROUSE	Dendragapus obscurus	
BNLC1003	ABNLC10030	WHITE-TAILED PTARMIGAN	Lagopus leucurus	watch list
BNLC1101	ABNLC11010	RUFFED GROUSE	Bonasa umbellus	
BNLC1201	ABNLC12010	SAGE GROUSE	Centrocercus urophasianus	watch list
BNLC1303	ABNLC13030	SHARP-TAILED GROUSE	Tympanuchus phasianellus	
BNLC1304	ABNLC13033	COLUMBIAN SHARP-TAILED GROUSE	Tympanuchus phasianellus columbianus	special concern
BNLC1401	ABNLC14010	WILD TURKEY	Meleagris gallopavo	
BNLC2102	ABNLC21020	NORTHERN BOBWHITE	Colinus virginianus	
BNME0101	ABNME01010	YELLOW RAIL	Coturnicops noveboracensis	special concern
BNME0503	ABNME05030	VIRGINIA RAIL	Rallus limicola	
BNME0802	ABNME08020	SORA	Porzana carolina	
BNME1402	ABNME14020	AMERICAN COOT	Fulica americana	
BNMK0101	ABNMK01010	SANDHILL CRANE	Grus canadensis	
BNMK0103	ABNMK01030	WHOOPIING CRANE	Grus americana	special concern
BNNB0201	ABNNB02010	BLACK-BELLIED PLOVER	Pluvialis squatarola	
BNNB0306	ABNNB03060	SEMIPALMATED PLOVER	Charadrius semipalmatus	
BNNB0307	ABNNB03070	PIPING PLOVER	Charadrius melodus	special concern
BNNB0309	ABNNB03090	KILLDEER	Charadrius vociferus	
BNNB0310	ABNNB03100	MOUNTAIN PLOVER	Charadrius montanus	special concern
BNND0101	ABNND01010	BLACK-NECKED STILT	Himantopus mexicanus	special concern
BNND0201	ABNND02010	AMERICAN AVOCET	Recurvirostra americana	
BNNF0102	ABNNF01020	GREATER YELLOWLEGS	Tringa melanoleuca	
BNNF0103	ABNNF01030	LESSER YELLOWLEGS	Tringa flavipes	
BNNF0107	ABNNF01070	SOLITARY SANDPIPER	Tringa solitaria	
BNNF0201	ABNNF02010	WILLET	Catoptrophorus semipalmatus	
BNNF0402	ABNNF04020	SPOTTED SANDPIPER	Actitis macularia	
BNNF0601	ABNNF06010	UPLAND SANDPIPER	Bartramia longicauda	
BNNF0707	ABNNF07070	LONG-BILLED CURLEW	Numenius americanus	
BNNF0804	ABNNF08040	MARbled GODWIT	Limosa fedoa	
BNNF1104	ABNNF11040	SEMIPALMATED SANDPIPER	Calidris pusilla	
BNNF1105	ABNNF11050	WESTERN SANDPIPER	Calidris mauri	
BNNF1110	ABNNF11100	LEAST SANDPIPER	Calidris minutilla	
BNNF1111	ABNNF11110	WHITE-RUMPED SANDPIPER	Calidris fuscicollis	
BNNF1112	ABNNF11120	BAIRD'S SANDPIPER	Calidris bairdii	
BNNF1113	ABNNF11130	PECTORAL SANDPIPER	Calidris melanotos	
BNNF1119	ABNNF11190	STILT SANDPIPER	Calidris himantopus	
BNNF1601	ABNNF16010	SHORT-BILLED DOWITCHER	Limnodromus griseus	
BNNF1602	ABNNF16020	LONG-BILLED DOWITCHER	Limnodromus scolopaceus	
BNNF1801	ABNNF18010	COMMON SNIPE	Gallinago gallinago	
BNNF2001	ABNNF20010	WILSON'S PHALAROPE	Phalaropus tricolor	
BNNF2002	ABNNF20020	RED-NECKED PHALAROPE	Phalaropus lobatus	
BNNM0302	ABNNM03020	FRANKLIN'S GULL	Larus pipixcan	special concern
BNNM0305	ABNNM03050	BONAPARTE'S GULL	Larus philadelphia	

BNNM0310	ABNNM03100	RING-BILLED GULL	<i>Larus delawarensis</i>	
BNNM0311	ABNNM03110	CALIFORNIA GULL	<i>Larus californicus</i>	
BNNM0802	ABNNM08020	CASPIAN TERN	<i>Sterna caspia</i>	special concern
BNNM0807	ABNNM08070	COMMON TERN	<i>Sterna hirundo</i>	special concern
BNNM0809	ABNNM08090	FORSTER'S TERN	<i>Sterna forsteri</i>	special concern
BNNM0810	ABNNM08100	LEAST TERN	<i>Sterna antillarum</i>	special concern
BNNM1002	ABNNM10020	BLACK TERN	<i>Chlidonias niger</i>	special concern
BNPB0101	ABNPB01010	ROCK DOVE	<i>Columbia livia</i>	
BNPB0404	ABNPB04040	MOURNING DOVE	<i>Zenaida macroura</i>	
BNRB0201	ABNRB02010	BLACK-BILLED CUCKOO	<i>Coccyzus erythrophthalmus</i>	
BNRB0202	ABNRB02020	YELLOW-BILLED CUCKOO	<i>Coccyzus americanus</i>	special concern
BNSA0101	ABNSA01010	BARN OWL	<i>Tyto alba</i>	watch list
BNSB0102	ABNSB01020	FLAMMULATED OWL	<i>Otus flammeolus</i>	special concern
BNSB0103	ABNSB01030	EASTERN SCREECH-OWL	<i>Otus asio</i>	watch list
BNSB0104	ABNSB01040	WESTERN SCREECH-OWL	<i>Otus kennicottii</i>	watch list
BNSB0501	ABNSB05010	GREAT HORNED OWL	<i>Bubo virginianus</i>	
BNSB0601	ABNSB06010	SNOWY OWL	<i>Nyctea scandiaca</i>	
BNSB0701	ABNSB07010	NORTHERN HAWK OWL	<i>Surnia ulula</i>	watch list
BNSB0801	ABNSB08010	NORTHERN PYGMY-OWL	<i>Glauclidium gnoma</i>	
BNSB1001	ABNSB10010	BURROWING OWL	<i>Athene cucularia</i>	special concern
BNSB1202	ABNSB12020	BARRED OWL	<i>Strix varia</i>	
BNSB1204	ABNSB12040	GREAT GRAY OWL	<i>Strix nebulosa</i>	special concern
BNSB1301	ABNSB13010	LONG-EARED OWL	<i>Asio otus</i>	
BNSB1304	ABNSB13040	SHORT-EARED OWL	<i>Asio flammeus</i>	
BNSB1501	ABNSB15010	BOREAL OWL	<i>Aegolius funereus</i>	special concern
BNSB1502	ABNSB15020	NORTHERN SAW-WHET OWL	<i>Aegolius acadicus</i>	
BNTA0202	ABNTA02020	COMMON NIGHTHAWK	<i>Chordeiles minor</i>	
BNTA0401	ABNTA04010	COMMON POORWILL	<i>Phalaenoptilus nuttallii</i>	
BNUA0101	ABNUA01010	BLACK SWIFT	<i>Cypseloides niger</i>	special concern
BNUA0301	ABNUA03010	CHIMNEY SWIFT	<i>Chaetura pelagica</i>	
BNUA0302	ABNUA03020	VAUX'S SWIFT	<i>Chaetura vauxi</i>	
BNUA0601	ABNUA06010	WHITE-THROATED SWIFT	<i>Aeronautes saxatalis</i>	
BNUC4501	ABNUC45010	RUBY-THROATED HUMMINGBIRD	<i>Archilochus colubris</i>	
BNUC4502	ABNUC45020	BLACK-CHINNED HUMMINGBIRD	<i>Archilochus alexandri</i>	
BNUC4801	ABNUC48010	CALLIOPE HUMMINGBIRD	<i>Stellula calliope</i>	
BNUC5102	ABNUC51020	RUFOUS HUMMINGBIRD	<i>Selasphorus rufus</i>	
BNXD0102	ABNXD01020	BELTED KINGFISHER	<i>Ceryle alcyon</i>	
BNYF0401	ABNYF04010	LEWIS'S WOODPECKER	<i>Melanerpes lewis</i>	
BNYF0404	ABNYF04040	RED-HEADED WOODPECKER	<i>Melanerpes erythrocephalus</i>	
BNYF0503	ABNYF05030	WILLIAMSON'S SAPSUCKER	<i>Sphyrapicus thyroideus</i>	
BNYF0504	ABNYF05040	RED-NAPED SAPSUCKER	<i>Sphyrapicus nuchalis</i>	
BNYF0703	ABNYF07030	DOWNY WOODPECKER	<i>Picoides pubescens</i>	
BNYF0704	ABNYF07040	HAIRY WOODPECKER	<i>Picoides villosus</i>	
BNYF0708	ABNYF07080	THREE-TOED WOODPECKER	<i>Picoides tridactylus</i>	
BNYF0709	ABNYF07090	BLACK-BACKED WOODPECKER	<i>Picoides arcticus</i>	special concern
BNYF1002	ABNYF10020	NORTHERN FLICKER	<i>Colaptes auratus</i>	
BNYF1202	ABNYF12020	PILEATED WOODPECKER	<i>Dryocopus pileatus</i>	
BPAE3201	ABPAE32010	OLIVE-SIDED FLYCATCHER	<i>Contopus cooperi</i>	
BPAE3205	ABPAE32050	WESTERN WOOD-PEWEE	<i>Contopus sordidulus</i>	
BPAE3303	ABPAE33030	ALDER FLYCATCHER	<i>Empidonax alnorum</i>	special concern
BPAE3304	ABPAE33040	WILLOW FLYCATCHER	<i>Empidonax traillii</i>	
BPAE3307	ABPAE33070	LEAST FLYCATCHER	<i>Empidonax minimus</i>	
BPAE3308	ABPAE33080	HAMMOND'S FLYCATCHER	<i>Empidonax hammondii</i>	
BPAE3309	ABPAE33090	DUSKY FLYCATCHER	<i>Empidonax oberholseri</i>	
BPAE3316	ABPAE33160	CORDILLERAN FLYCATCHER	<i>Empidonax occidentalis</i>	
BPAE3503	ABPAE35030	SAY'S PHOEBE	<i>Sayornis saya</i>	
BPAE5203	ABPAE52030	CASSIN'S KINGBIRD	<i>Tyrannus vociferans</i>	special concern
BPAE5205	ABPAE52050	WESTERN KINGBIRD	<i>Tyrannus verticalis</i>	
BPAE5206	ABPAE52060	EASTERN KINGBIRD	<i>Tyrannus tyrannus</i>	
BPAT0201	ABPAT02010	HORNED LARK	<i>Eremophila alpestris</i>	
BPAT0301	ABPAU03010	TREE SWALLOW	<i>Tachycineta bicolor</i>	

BPAU0304	ABPAU03040	VIOLET-GREEN SWALLOW	Tachycineta thalassina	
BPAU0701	ABPAU07010	NORTHERN ROUGH-WINGED SWALLOW	Stelgidopteryx serripennis	
BPAU0801	ABPAU08010	BANK SWALLOW	Riparia riparia	
BPAU0901	ABPAU09010	CLIFF SWALLOW	Petrochelidon pyrrhonota	
BPAU0903	ABPAU09030	BARN SWALLOW	Hirundo rustica	
BPAV0101	ABPAV01010	GRAY JAY	Perisoreus canadensis	
BPAV0201	ABPAV02010	STELLER'S JAY	Cyanocitta stelleri	
BPAV0202	ABPAV02020	BLUE JAY	Cyanocitta cristata	
BPAV0701	ABPAV07010	PINYON JAY	Gymnorhinus cyanocephalus	
BPAV0801	ABPAV08010	CLARK'S NUTCRACKER	Nucifraga columbiana	
BPAV0901	ABPAV09010	BLACK-BILLED MAGPIE	Pica pica	
BPAV1001	ABPAV10010	AMERICAN CROW	Corvus brachyrhynchos	
BPAV1011	ABPAV10110	COMMON RAVEN	Corvus corax	
BPAW0101	ABPAW01010	BLACK-CAPPED CHICKADEE	Poecile atricapillus	
BPAW0104	ABPAW01040	MOUNTAIN CHICKADEE	Poecile gambeli	
BPAW0106	ABPAW01060	BOREAL CHICKADEE	Poecile hudsonicus	
BPAW0107	ABPAW01070	CHESTNUT-BACKED CHICKADEE	Poecile rufescens	
BPAZ0101	ABPAZ01010	RED-BREASTED NUTHATCH	Sitta canadensis	
BPAZ0102	ABPAZ01020	WHITE-BREASTED NUTHATCH	Sitta carolinensis	
BPAZ0103	ABPAZ01030	PYGMY NUTHATCH	Sitta pygmaea	
BPBA0101	ABPBA01010	BROWN CREEPER	Certhia americana	
BPBG0301	ABPBG03010	ROCK WREN	Salpinctes obsoletus	
BPBG0401	ABPBG04010	CANYON WREN	Catherpes mexicanus	
BPBG0901	ABPBG09010	HOUSE WREN	Troglodytes aedon	
BPBG0905	ABPBG09050	WINTER WREN	Troglodytes troglodytes	
BPBG1002	ABPBG10020	MARSH WREN	Cistothorus palustris	
BPBH0101	ABPBH01010	AMERICAN DIPPER	Cinclus mexicanus	
BPBJ0501	ABPBJ05010	GOLDEN-CROWNED KINGLET	Regulus satrapa	
BPBJ0502	ABPBJ05020	RUBY-CROWNED KINGLET	Regulus calendula	
BPBJ0801	ABPBJ08010	BLUE-GRAY GNATCATCHER	Poliptila caerulea	special concern
BPBJ1501	ABPBJ15010	EASTERN BLUEBIRD	Sialia sialis	
BPBJ1502	ABPBJ15020	WESTERN BLUEBIRD	Sialia mexicana	
BPBJ1503	ABPBJ15030	MOUNTAIN BLUEBIRD	Sialia currucoides	
BPBJ1601	ABPBJ16010	TOWNSEND'S SOLITAIRE	Myadestes townsendi	
BPBJ1808	ABPBJ18080	VEERY	Catharus fuscescens	
BPBJ1809	ABPBJ18090	GRAY-CHEEKED THRUSH	Catharus minimus	
BPBJ1810	ABPBJ18100	SWAINSON'S THRUSH	Catharus ustulatus	
BPBJ1811	ABPBJ18110	HERMIT THRUSH	Catharus guttatus	
BPBJ2017	ABPBJ20170	AMERICAN ROBIN	Turdus migratorius	
BPBJ2201	ABPBJ22010	VARIED THRUSH	Ixoreus naevius	
BPBK0101	ABPBK01010	GRAY CATBIRD	Dumetella carolinensis	
BPBK0401	ABPBK04010	SAGE THRASHER	Oreoscoptes montanus	
BPBK0601	ABPBK06010	BROWN THRASHER	Toxostoma rufum	
BPBM0205	ABPBM02050	AMERICAN PIPIT	Anthus rubescens	
BPBM0206	ABPBM02060	SPRAGUE'S PIPIT	Anthus spragueii	
BPBN0101	ABPBN01010	BOHEMIAN WAXWING	Bombycilla garrulus	
BPBN0102	ABPBN01020	CEDAR WAXWING	Bombycilla cedrorum	
BPBR0102	ABPBR01020	NORTHERN SHRIKE	Lanius excubitor	
BPBR0103	ABPBR01030	LOGGERHEAD SHRIKE	Lanius ludovicianus	watch list
BPBT0101	ABPBT01010	EUROPEAN STARLING	Sturnus vulgaris	
BPBW0116	ABPBW01160	SOLITARY VIREO	Vireo solitarius	
BPBW0121	ABPBW01210	WARBLING VIREO	Vireo gilvus	
BPBW0124	ABPBW01240	RED-EYED VIREO	Vireo olivaceus	
BPBX0104	ABPBX01040	TENNESSEE WARBLER	Vermivora peregrina	watch list
BPBX0105	ABPBX01050	ORANGE-CROWNED WARBLER	Vermivora celata	
BPBX0106	ABPBX01060	NASHVILLE WARBLER	Vermivora ruficapilla	
BPBX0301	ABPBX03010	YELLOW WARBLER	Dendroica petechia	
BPBX0306	ABPBX03060	YELLOW-RUMPED WARBLER	Dendroica coronata	
BPBX0308	ABPBX03080	TOWNSEND'S WARBLER	Dendroica townsendi	
BPBX0323	ABPBX03230	BLACKPOLL WARBLER	Dendroica striata	
BPBX0501	ABPBX05010	BLACK-AND-WHITE WARBLER	Mniotilta varia	watch list

BPBX0601	ABPBX06010	AMERICAN REDSTART	<i>Setophaga ruticilla</i>	
BPBX1001	ABPBX10010	OVENBIRD	<i>Seiurus aurocapillus</i>	
BPBX1002	ABPBX10020	NORTHERN WATERTHRUSH	<i>Seiurus noveboracensis</i>	
BPBX1104	ABPBX11040	MACGILLIVRAY'S WARBLER	<i>Oporornis tolmiei</i>	
BPBX1201	ABPBX12010	COMMON YELLOWTHROAT	<i>Geothlypis trichas</i>	
BPBX1602	ABPBX16020	WILSON'S WARBLER	<i>Wilsonia pusilla</i>	
BPBX2401	ABPBX24010	YELLOW-BREASTED CHAT	<i>Icteria virens</i>	
BPBX4505	ABPBX45050	WESTERN Tanager	<i>Piranga ludoviciana</i>	
BPBX6104	ABPBX61040	BLACK-HEADED GROSBEAK	<i>Pheucticus melanocephalus</i>	
BPBX6402	ABPBX64020	LAZULI BUNTING	<i>Passerina amoena</i>	
BPBX6403	ABPBX64030	INDIGO BUNTING	<i>Passerina cyanea</i>	
BPBX6501	ABPBX65010	DICKCISSEL	<i>Spiza americana</i>	special concern
BPBX7401	ABPBX74010	GREEN-TAILED TOWHEE	<i>Pipilo chlorurus</i>	
BPBX7408	ABPBX74080	SPOTTED TOWHEE	<i>Pipilo maculatus</i>	
BPBX9401	ABPBX94010	AMERICAN TREE SPARROW	<i>Spizella arborea</i>	
BPBX9402	ABPBX94020	CHIPPING SPARROW	<i>Spizella passerina</i>	
BPBX9403	ABPBX94030	CLAY-COLORED SPARROW	<i>Spizella pallida</i>	
BPBX9404	ABPBX94040	BREWER'S SPARROW	<i>Spizella breweri</i>	
BPBX9405	ABPBX94050	FIELD SPARROW	<i>Spizella pusilla</i>	
BPBX9501	ABPBX95010	VESPER SPARROW	<i>Poocetes gramineus</i>	
BPBX9601	ABPBX96010	LARK SPARROW	<i>Chondestes grammacus</i>	
BPBX9801	ABPBX98010	LARK BUNTING	<i>Calamospiza melanocorys</i>	
BPBX9901	ABPBX99010	SAVANNAH SPARROW	<i>Passerculus sandwichensis</i>	
BPBXA001	ABPBXA0010	BAIRD'S SPARROW	<i>Ammodramus bairdii</i>	special concern
BPBXA002	ABPBXA0020	GRASSHOPPER SPARROW	<i>Ammodramus savannarum</i>	
BPBXA004	ABPBXA0040	LE CONTE'S SPARROW	<i>Ammodramus leconteii</i>	special concern
BPBXA007	ABPBXA0070	NELSON'S SHARP-TAILED SPARROW	<i>Ammodramus nelsoni</i>	special concern
BPBXA201	ABPBXA2010	FOX SPARROW	<i>Passerella iliaca</i>	
BPBXA301	ABPBXA3010	SONG SPARROW	<i>Melospiza melodia</i>	
BPBXA302	ABPBXA3020	LINCOLN'S SPARROW	<i>Melospiza lincolni</i>	
BPBXA402	ABPBXA4020	WHITE-THROATED SPARROW	<i>Zonotrichia albicollis</i>	
BPBXA404	ABPBXA4040	WHITE-CROWNED SPARROW	<i>Zonotrichia leucophrys</i>	
BPBXA405	ABPBXA4050	HARRIS'S SPARROW	<i>Zonotrichia querula</i>	
BPBXA502	ABPBXA5020	DARK-EYED JUNCO	<i>Junco hyemalis</i>	
BPBXA601	ABPBXA6010	MCCOWN'S LONGSPUR	<i>Calcarius mccownii</i>	
BPBXA602	ABPBXA6020	LAPLAND LONGSPUR	<i>Calcarius lapponicus</i>	
BPBXA604	ABPBXA6040	CHESTNUT-COLLARED LONGSPUR	<i>Calcarius ornatus</i>	
BPBXA801	ABPBXA8010	SNOW BUNTING	<i>Plectrophenax nivalis</i>	
BPBXA901	ABPBXA9010	BOBOLINK	<i>Dolichonyx oryzivorus</i>	
BPBXB001	ABPBXB0010	RED-WINGED BLACKBIRD	<i>Agelaius phoeniceus</i>	
BPBXB203	ABPBXB2030	WESTERN MEADOWLARK	<i>Sturnella neglecta</i>	
BPBXB301	ABPBXB3010	YELLOW-HEADED BLACKBIRD	<i>Xanthocephalus xanthocephalus</i>	
BPBXB501	ABPBXB5010	RUSTY BLACKBIRD	<i>Euphagus carolinus</i>	
BPBXB502	ABPBXB5020	BREWER'S BLACKBIRD	<i>Euphagus cyanocephalus</i>	
BPBXB607	ABPBXB6070	COMMON GRACKLE	<i>Quiscalus quiscula</i>	
BPBXB703	ABPBXB7030	BROWN-HEADED COWBIRD	<i>Molothrus ater</i>	
BPBXB907	ABPBXB9070	ORCHARD ORIOLE	<i>Icterus spurius</i>	
BPBXB922	ABPBXB9220	BULLOCK'S ORIOLE	<i>Icterus bullockii</i>	
BPBY0201	ABPBY02010	BLACK ROSY-FINCH	<i>Leucosticte atrata</i>	
BPBY0203	ABPBY02030	GRAY-CROWNED ROSY-FINCH	<i>Leucosticte tephrocotis</i>	
BPBY0301	ABPBY03010	PINE GROSBEAK	<i>Pinicola enucleator</i>	
BPBY0402	ABPBY04020	PURPLE FINCH	<i>Carpodacus purpureus</i>	
BPBY0403	ABPBY04030	CASSIN'S FINCH	<i>Carpodacus cassinii</i>	
BPBY0404	ABPBY04040	HOUSE FINCH	<i>Carpodacus mexicanus</i>	
BPBY0501	ABPBY05010	RED CROSSBILL	<i>Loxia curvirostra</i>	
BPBY0502	ABPBY05020	WHITE-WINGED CROSSBILL	<i>Loxia leucoptera</i>	
BPBY0601	ABPBY06010	COMMON REDPOLL	<i>Carduelis flammæa</i>	
BPBY0602	ABPBY06020	HOARY REDPOLL	<i>Carduelis hornemanni</i>	
BPBY0603	ABPBY06030	PINE SISKIN	<i>Carduelis pinus</i>	
BPBY0611	ABPBY06110	AMERICAN GOLDFINCH	<i>Carduelis tristis</i>	
BPBY0902	ABPBY09020	EVENING GROSBEAK	<i>Coccothraustes vespertinus</i>	

BPBZ0101	ABPBZ01010	HOUSE SPARROW	<i>Passer domesticus</i>	
MABA0101	AMABA01010	MASKED SHREW	<i>Sorex cinereus</i>	
MABA0103	AMABA01030	PREBLE'S SHREW	<i>Sorex preblei</i>	
MABA0107	AMABA01070	VAGRANT SHREW	<i>Sorex vagrans</i>	
MABA0108	AMABA01080	DUSKY OR MONTANE SHREW	<i>Sorex monticolus</i>	
MABA0113	AMABA01130	DWARF SHREW	<i>Sorex nanus</i>	special concern
MABA0115	AMABA01150	WATER SHREW	<i>Sorex palustris</i>	
MABA0123	AMABA01230	MERRIAM'S SHREW	<i>Sorex merriami</i>	special concern
MABA0125	AMABA01250	PYGMY SHREW	<i>Sorex hoyi</i>	
MABA0128	AMABA01280	HAYDEN'S SHREW	<i>Sorex haydeni</i>	
MACC0101	AMACC01010	LITTLE BROWN MYOTIS	<i>Myotis lucifugus</i>	
MACC0102	AMACC01020	YUMA MYOTIS	<i>Myotis yumanensis</i>	watch list
MACC0107	AMACC01070	LONG-EARED MYOTIS	<i>Myotis evotis</i>	
MACC0109	AMACC01090	FRINGED MYOTIS	<i>Myotis thysanodes</i>	special concern
MACC0111	AMACC01110	LONG-LEGGED MYOTIS	<i>Myotis volans</i>	
MACC0112	AMACC01120	CALIFORNIA MYOTIS	<i>Myotis californicus</i>	
MACC0114	AMACC01140	WESTERN SMALL-FOOTED MYOTIS	<i>Myotis ciliolabrum</i>	
MACC0115	AMACC01150	NORTHERN MYOTIS	<i>Myotis septentrionalis</i>	special concern
MACC0201	AMACC02010	SILVER-HAIRED BAT	<i>Lasionycteris noctivagans</i>	
MACC0401	AMACC04010	BIG BROWN BAT	<i>Eptesicus fuscus</i>	
MACC0503	AMACC05030	HOARY BAT	<i>Lasiurus cinereus</i>	
MACC0701	AMACC07010	SPOTTED BAT	<i>Euderma maculatum</i>	special concern
MACC0801	AMACC08010	TOWNSEND'S BIG-EARED BAT	<i>Corynorhinus townsendii</i>	special concern
MACC1001	AMACC10010	PALLID BAT	<i>Antrozous pallidus</i>	
MAEA0102	AMAEA01020	AMERICAN PIKA	<i>Ochotona princeps</i>	
MAEB0104	AMAEB01040	EASTERN COTTONTAIL	<i>Sylvilagus floridanus</i>	watch list
MAEB0106	AMAEB01060	MOUNTAIN COTTONTAIL	<i>Sylvilagus nuttallii</i>	
MAEB0107	AMAEB01070	DESERT COTTONTAIL	<i>Sylvilagus audubonii</i>	
MAEB0301	AMAEB03010	SNOWSHOE HARE	<i>Lepus americanus</i>	
MAEB0304	AMAEB03040	WHITE-TAILED JACKRABBIT	<i>Lepus townsendii</i>	
MAEB0305	AMAEB03050	BLACK-TAILED JACKRABBIT	<i>Lepus californicus</i>	special concern
MAEB0401	AMAEB04010	PYGMY RABBIT	<i>Brachylagus idahoensis</i>	special concern
MAFB0202	AMAFB02020	LEAST CHIPMUNK	<i>Tamias minimus</i>	
MAFB0203	AMAFB02030	YELLOW-PINE CHIPMUNK	<i>Tamias amoenus</i>	
MAFB0213	AMAFB02130	RED-TAILED CHIPMUNK	<i>Tamias ruficaudus</i>	
MAFB0219	AMAFB02190	UINTA CHIPMUNK	<i>Tamias umbrinus</i>	special concern
MAFB0302	AMAFB03020	YELLOW-BELLIED MARMOT	<i>Marmota flaviventris</i>	
MAFB0304	AMAFB03040	HOARY MARMOT	<i>Marmota caligata</i>	
MAFB0504	AMAFB05040	RICHARDSON'S GROUND SQUIRREL	<i>Spermophilus richardsonii</i>	
MAFB0505	AMAFB05050	UINTA GROUND SQUIRREL	<i>Spermophilus armatus</i>	
MAFB0507	AMAFB05070	COLUMBIAN GROUND SQUIRREL	<i>Spermophilus columbianus</i>	
MAFB0509	AMAFB05090	THIRTEEN-LINED GROUND SQUIRREL	<i>Spermophilus tridecemlineatus</i>	
MAFB0517	AMAFB05170	GOLDEN-MANTLED GROUND SQUIRREL	<i>Spermophilus lateralis</i>	
MAFB0519	AMAFB05190	WYOMING GROUND SQUIRREL	<i>Spermophilus elegans</i>	
MAFB0601	AMAFB06010	BLACK-TAILED PRAIRIE DOG	<i>Cynomys ludovicianus</i>	special concern
MAFB0602	AMAFB06020	WHITE-TAILED PRAIRIE DOG	<i>Cynomys leucurus</i>	special concern
MAFB0801	AMAFB08010	RED SQUIRREL	<i>Tamiasciurus hudsonicus</i>	
MAFB0902	AMAFB09020	NORTHERN FLYING SQUIRREL	<i>Glaucomys sabrinus</i>	
MAFC0104	AMAF01040	NORTHERN POCKET GOPHER	<i>Thomomys talpoides</i>	
MAFC0107	AMAF01070	IDAHO POCKET GOPHER	<i>Thomomys idahoensis</i>	
MAFD0101	AMAFD01010	OLIVE-BACKED POCKET MOUSE	<i>Perognathus fasciatus</i>	
MAFD0107	AMAFD01070	GREAT BASIN POCKET MOUSE	<i>Perognathus parvus</i>	special concern
MAFD0301	AMAFD03010	ORD'S KANGAROO RAT	<i>Dipodomys ordii</i>	
MAFD0505	AMAFD05050	HISPID POCKET MOUSE	<i>Chaetodipus hispidus</i>	special concern
MAFE0101	AMAFE01010	AMERICAN BEAVER	<i>Castor canadensis</i>	
MAFF0203	AMAFF02030	WESTERN HARVEST MOUSE	<i>Reithrodontomys megalotis</i>	
MAFF0304	AMAFF03040	DEER MOUSE	<i>Peromyscus maniculatus</i>	
MAFF0307	AMAFF03070	WHITE-FOOTED MOUSE	<i>Peromyscus leucopus</i>	
MAFF0601	AMAFF06010	NORTHERN GRASSHOPPER MOUSE	<i>Onychomys leucogaster</i>	
MAFF0809	AMAFF08090	BUSHY-TAILED WOODRAT	<i>Neotoma cinerea</i>	
MAFF0902	AMAFF09020	SOUTHERN RED-BACKED VOLE	<i>Clethrionomys gapperi</i>	


MAFF1001	AMAFF10010	HEATHER VOLE	<i>Phenacomys intermedius</i>	
MAFF1101	AMAFF11010	MEADOW VOLE	<i>Microtus pennsylvanicus</i>	
MAFF1102	AMAFF11020	MONTANE VOLE	<i>Microtus montanus</i>	
MAFF1106	AMAFF11060	LONG-TAILED VOLE	<i>Microtus longicaudus</i>	
MAFF1114	AMAFF11140	PRAIRIE VOLE	<i>Microtus ochrogaster</i>	
MAFF1119	AMAFF11190	WATER VOLE	<i>Microtus richardsoni</i>	
MAFF1301	AMAFF13010	SAGEBRUSH VOLE	<i>Lemmiscus curtatus</i>	
MAFF1501	AMAFF15010	MUSKRAT	<i>Ondatra zibethicus</i>	
MAFF1702	AMAFF17020	NORTHERN BOG LEMMING	<i>Synaptomys borealis</i>	special concern
MAFH0101	AMAFH01010	MEADOW JUMPING MOUSE	<i>Zapus hudsonius</i>	special concern
MAFH0102	AMAFH01020	WESTERN JUMPING MOUSE	<i>Zapus princeps</i>	
MAFJ0101	AMAFJ01010	COMMON PORCUPINE	<i>Erethizon dorsatum</i>	
MAJA0101	AMAJA01010	COYOTE	<i>Canis latrans</i>	
MAJA0103	AMAJA01030	GRAY WOLF	<i>Canis lupus</i>	special concern
MAJA0301	AMAJA03010	RED FOX	<i>Vulpes vulpes</i>	
MAJA0303	AMAJA03030	SWIFT FOX	<i>Vulpes velox</i>	special concern
MAJB0101	AMAJB01010	BLACK BEAR	<i>Ursus americanus</i>	
MAJB0102	AMAJB01020	GRIZZLY OR BROWN BEAR	<i>Ursus arctos</i>	special concern
MAJE0201	AMAJE02010	COMMON RACCOON	<i>Procyon lotor</i>	
MAJF0101	AMAJF01010	AMERICAN MARTEN	<i>Martes americana</i>	
MAJF0102	AMAJF01020	FISHER	<i>Martes pennanti</i>	special concern
MAJF0201	AMAJF02010	ERMINE	<i>Mustela erminea</i>	
MAJF0202	AMAJF02020	LEAST WEASEL	<i>Mustela nivalis</i>	
MAJF0203	AMAJF02030	LONG-TAILED WEASEL	<i>Mustela frenata</i>	
MAJF0204	AMAJF02040	BLACK-FOOTED FERRET	<i>Mustela nigripes</i>	special concern
MAJF0205	AMAJF02050	MINK	<i>Mustela vison</i>	
MAJF0301	AMAJF03010	WOLVERINE	<i>Gulo gulo</i>	special concern
MAJF0401	AMAJF04010	AMERICAN BADGER	<i>Taxidea taxus</i>	
MAJF0502	AMAJF05020	WESTERN SPOTTED SKUNK	<i>Spilogale gracilis</i>	watch list
MAJF0601	AMAJF06010	STRIPED SKUNK	<i>Mephitis mephitis</i>	
MAJF0801	AMAJF08010	NORTHERN RIVER OTTER	<i>Lutra canadensis</i>	
MAJH0102	AMAJH01020	MOUNTAIN LION	<i>Felis concolor</i>	
MAJH0301	AMAJH03010	LYNX	<i>Lynx canadensis</i>	special concern
MAJH0302	AMAJH03020	BOBCAT	<i>Lynx rufus</i>	
MALC0101	AMALC01010	WAPITI OR ELK	<i>Cervus elaphus</i>	
MALC0201	AMALC02010	MULE DEER	<i>Odocoileus hemionus</i>	
MALC0202	AMALC02020	WHITE-TAILED DEER	<i>Odocoileus virginianus</i>	
MALC0301	AMALC03010	MOOSE	<i>Alces alces</i>	
MALD0101	AMALD01010	PRONGHORN	<i>Antilocapra americana</i>	
MALE0101	AMALE01010	AMERICAN BISON	<i>Bos bison</i>	
MALE0201	AMALE02010	MOUNTAIN GOAT	<i>Oreamnos americanus</i>	
MALE0401	AMALE04010	MOUNTAIN SHEEP	<i>Ovis canadensis</i>	
RAAB0101	ARAAB01010	SNAPPING TURTLE	<i>Chelydra serpentina</i>	special concern
RAAD0101	ARAAD01010	PAINTED TURTLE	<i>Chrysemys picta</i>	
RAAG0103	ARAAG01030	SPINY SOFTSHELL	<i>Apalone spinifera</i>	special concern
RACB0101	ARACB01010	NORTHERN ALLIGATOR LIZARD	<i>Elgaria coerulea</i>	
RACF1203	ARACF12030	SHORT-HORNED LIZARD	<i>Phrynosoma douglasii</i>	watch list
RACF1403	ARACF14030	SAGEBRUSH LIZARD	<i>Sceloporus graciosus</i>	watch list
RACH0111	ARACH01110	WESTERN SKINK	<i>Eumeces skiltonianus</i>	watch list
RADA0101	ARADA01010	RUBBER BOA	<i>Charina bottae</i>	
RADB0701	ARADB07010	RACER	<i>Coluber constrictor</i>	
RADB1701	ARADB17010	WESTERN HOGNOSE SNAKE	<i>Heterodon nasicus</i>	special concern
RADB1905	ARADB19050	MILK SNAKE	<i>Lampropeltis triangulum</i>	special concern
RADB2601	ARADB26010	PINE OR GOPHER SNAKE	<i>Pituophis melanoleucus</i>	
RADB3605	ARADB36050	WESTERN TERRESTRIAL GARTER SNAKE	<i>Thamnophis elegans</i>	
RADB3610	ARADB36100	PLAINS GARTER SNAKE	<i>Thamnophis radix</i>	
RADB3613	ARADB36130	COMMON GARTER SNAKE	<i>Thamnophis sirtalis</i>	watch list
RADB4701	ARADB47010	SMOOTH GREEN SNAKE	<i>Liochlorophis vernalis</i>	special concern
RADE0212	ARADE02120	WESTERN RATTLESNAKE	<i>Crotalus viridis</i>	

**Predicted Distributions for Species of Special Concern and Watch List
Species and Potential Conflicts with Montana Highways**

COEUR D'ALENE SALAMANDER

Plethodon idahoensis

State rank	S2	Element code	AAAAD12270
Global rank	G3	Modeled by	M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell
MTNHP status	special concern		



State range

Rare and locally distributed in the Clark Fork, Kootenai and Bitterroot River drainages in northwestern Montana from Lake Kooconusa at the Canadian border south to the Bitterroot Valley. Range from 500 to 1,550 meters elevation (Wilson et al. 1997).

Habitat description

Coeur d'Alene salamanders respire through their skin and lose water to the environment through evaporation. They are therefore restricted to cool, damp environments (Groves et al. 1996). Primary habitats include springs or seeps, waterfall spray zones and damp streambanks in talus or fractured rock sites with conifer forest nearby (Wilson et al. 1997). However, habitats also include leaf litter, bark and rotting logs in damp conifer forests, north-facing talus slopes with abundant moist soil, and mesic forest types containing moist microhabitats (Groves et al. 1996). The species is found in conjunction with both persistent and intermittent surface waters, but depends on the presence of stable subterranean water flows (Groves et al. 1996). Adults are usually above ground only at night during moist weather; surface activity is negatively correlated with high daytime temperatures and days since last rain (Wilson and Larsen 1988). Adults breed terrestrially in late summer, fall and, to a lesser extent, in the spring. Females deposit eggs in April or May, presumably in underground rock crevices, although no nest sites have been found in Montana. Juveniles emerge directly from the eggs in mid-September (Lynch 1984).


Model assumptions & caveats

Model based on species' dependence on hydric stability of subterranean habitats immediately adjacent to springs, seeps, waterfalls and streamsides with canopy cover (Groves et al. 1996). An elevation limit of 1,650 meters was selected to eliminate areas 100 meters above the species known elevational limits (Wilson et al. 1997). Springs were not included in the model because an adequate GIS layer was not available. Buffers were not used because the species is not known to inhabit areas far from stable hydrographic features. Initial review of the model showed intermediate levels of accuracy in

predicting the species presence in areas where it is known to occur. The model is not likely to accurately represent the species' distribution because it cannot adequately identify specific microsites that support the species.

References


- Cassirer, E.F., C.R. Groves, and D. Genter. 1994. Coeur d'Alene salamander conservation assessment; report to U.S.D.A. Forest Service Region 1. Idaho Department of Fish and Game, Boise, and Montana Natural Heritage Program, Helena, MT. 55 pp.
- Groves, C.R., E.F. Cassirer, D.L. Genter and J.D. Reichel. 1996. Element Stewardship Abstract for the Coeur d'Alene Salamander (*Plethodon idahoensis*). *Natural Areas Journal* 16: 238-247.
- Howard, J. 1993. Genetic variation and population divergence in the *Plethodon vandykei* species group. *Herpetologica* 49: 238-247.
- Lynch, J.E. 1984. Reproductive ecology of *Plethodon idahoensis*. M.S. thesis, University of Idaho, Moscow, ID. 59 pp.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Werner, J.K. and J.D. Reichel. 1996. Amphibian and reptile monitoring/survey of the Kootenai National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 115 pp.
- Wilson, A.G. Jr. and J.H. Larsen Jr. 1988. Activity and diet in seepage-dwelling Coeur d'Alene salamanders (*Plethodon vandykei idahoensis*). *Northwest Science* 62(5): 211-217.
- Wilson, A.G. Jr., E.M. Wilson, C.R. Groves, and R.L. Wallace. 1997. U.S. distribution of the Coeur d'Alene Salamander (*Plethodon idahoensis* Slater and Slipp). *Great Basin Nat.* 57(4): 359-362.



IDAHO GIANT SALAMANDER

Dicamptodon aterrimus

State rank	SR	Element code	AAAAH01030
Global rank	G3	Modeled by	M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell
MTNHP status	special concern		



State range

Range unclear. Reported, but not confirmed, at only two locations near the town of Saltese in western Montana; however, found in Idaho adjacent to the state border from the Clark Fork River to the southern end of the Bitterroot Valley (Reichel and Flath 1995; Peterson and Fabian 1997). Ranges up to 2,165 meters elevation in Idaho (Nussbaum et al. 1983).

Habitat description

Although seldom seen, adults are found terrestrially in moist coniferous forests under rocks, bark and logs and aquatically under stones in mountain streams or lakes. Adults are active terrestrially on warm, rainy nights (Nussbaum et al. 1983). Adults breed in the spring or fall in hidden water-filled nest chambers beneath logs and stones or in crevices in mountain streams or lakes. Females subsequently deposit eggs in these chambers and guard the eggs throughout the incubation period (Nussbaum et al. 1983). Larvae are usually found in swift, cold mountain streams, but may occasionally be found in lakes or ponds. Larvae usually metamorphose in 18-24 months, but may become sexually mature (paedogenesis) and reproduce as larvae (Nussbaum et al. 1983).


Model assumptions & caveats

The species' range and status is unclear in the state. Model mostly developed for survey purposes; not to be included in final MT-GAP analyses.

Model based on species dependence on mountain streams, lakes and ponds for breeding and larval rearing and moist conifer forests for terrestrial adults. An elevation limit was not included because the species is known to occur at elevations exceeding those containing suitable habitat within its range in Montana (D. Pilliod, pers. comm.). Lacking observations, model accuracy could not be assessed. However, the model undoubtedly is an overestimate of the species' distribution.

References

- Cassirer, E.F. 1995. Wildlife inventory, Craig Mountain, Idaho. Idaho Department of Fish and Game, Lewiston Idaho. 182 pp.
- Daugherty, C.H., F.W. Allendorf, W.W. Dunlap and K.L. Knudsen. 1983. Systematic implications of geographic patterns of genetic variation in the genus *Dicamptodon*. *Copeia* 1983: 679-691.
- Good, D.A. 1989. Hybridization and cryptic species in *Dicamptodon* Caudata *Dicamptodontidae*. *Evolution* 43(4): 728-744.
- Maughan, O.E., M.G. Wickham, P. Laumeyer and R.L. Wallace. 1976. Records of the Pacific Giant Salamander, *Dicamptodon ensatus*, (Amphibia, Urodela, Ambystomatidae) from the Rocky Mountains in Idaho. *Journal of Herpetology* 10(3): 249-251.
- Nussbaum, R. A., E. D. Brodie, Jr., and R. M. Storm. 1983. Amphibians and reptiles of the Pacific Northwest. University of Idaho Press. Moscow, ID. 332 pp.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Thompson, L.S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Bozeman: Montana Audubon Council. 24 pp.



TAILED FROG

Ascaphus truei


State rank S4

Global rank G4

MTNHP status watch list

Element code AAABA01010

Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell



Predicted habitat: 492,122 ha, 1.29 % of state.

State range

Ranges west of the continental divide and in suitable habitats in the eastern front ranges immediately adjacent to the continental divide. Ranges to 2,590 meters elevation in Idaho (David Pilliod, pers. comm.).

Habitat description

Found in fast, small, permanent forest streams with clear, cold water, cobble or boulder substrates, and little silt (Franz 1971, Welsh 1990). In Montana, adults usually remain underwater hidden by rocks or debris and emerge at night or during humid weather from May to September to feed terrestrially along stream edges (Daugherty and Sheldon 1982a). Adults are highly philopatric, but are known to forage up to 25 or more meters away from water (Daugherty and Sheldon 1982b, Nussbaum et al. 1983). In Montana adults breed via internal fertilization in streams in August or September and females deposit eggs in long double strings under large stones in areas with slight current the following June or July (Franz 1970, Daugherty and Sheldon 1982a). Eggs hatch in August or September, and tadpoles cling to the undersides or tops of smooth rocks which lack periphyton or silt (Nussbaum et al. 1983). Tadpoles usually metamorphose in the third summer after hatching and adults reproduce for the first time four or five years after metamorphosis; females reproduce in alternate years thereafter (Daugherty and Sheldon 1982a).


Model assumptions & caveats

Model based on species' dependence on streams, riparian areas and moist conifer forests. Major rivers and lakes were removed because the species is not normally found in these habitats. An elevation limit was not included because the species is known to occur at elevations exceeding those containing suitable habitat within its range in Montana (D. Pilliod, pers. comm.). Buffers were not used because the species is not known to inhabit areas far from stable hydrographic features in Montana. Initial review of the model showed high levels of accuracy in predicting the species presence in areas where it is known to occur. The model is probably an accurate representation of the species' distribution, with

the exception of fragments of habitat propagated beyond the known range in southwestern Montana and on the East Front.

References

- Brunson, R. B., and H. A. Demaree, Jr. 1951. The herpetology of the Mission Mountains, Montana. *Copeia* 1951:306-308.
- Daugherty, C.H. 1979. Population ecology and genetics of *Ascaphus truei*: an examination of gene flow and natural selection. Ph.D. dissertation, University of Montana. Missoula, MT. 143 pp.
- Daugherty, C.H. and A.L. Sheldon. 1982a. Age-determination, growth, and life history of a Montana population of the tailed frog (*Ascaphus truei*). *Herpetologica* 38(4): 461-468.
- Daugherty, C.H. and A.L. Sheldon. 1982b. Age specific movement patterns of the frog *Ascaphus truei*. *Herpetologica* 38(4): 468-474.
- Franz, R. 1970. Egg development of the tailed frog under natural conditions. *Bulletin of the Maryland Herpetological Society* 6(2): 27-30.
- Franz, R. 1971. Notes on the distribution and ecology of the herpetofauna of northwestern Montana. *Bulletin of the Maryland Herpetological Society* 7: 1-10.
- Franz, R. and D.S. Lee. 1970. The ecological and biogeographical distribution of the tailed frog, *Ascaphus truei*, in the Flathead River drainage of northwestern Montana. *Bulletin of the Maryland Herpetological Society* 6: 62-73.
- Hendricks, P., and J. D. Reichel. 1996. Amphibian and reptile survey of the Bitterroot National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 95 pp.
- Landreth, H.F. and D.E. Ferguson. 1967. Movements and orientation of the tailed frog, *Ascaphus truei*. *Herpetologica* 23: 81-93.
- Marnell, L. E. 1997. Herpetofauna of Glacier National Park. *Northwest. Nat.* 78:17-33.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Reichel, J. D. 1995. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.
- Reichel, J.D. 1996. Preliminary amphibian and reptile survey of the Helena National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 87 pp.
- Welsh, H. H., Jr. 1990. Relictual amphibians and old-growth forests. *Conservation Biology* 4: 309-319.
- Werner, J.K. and J.D. Reichel. 1996. Amphibian and reptile monitoring/survey of the Kootenai National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 115 pp.
- Werner, J.K. and T. Plummer. 1995. Amphibian and reptile survey of the Flathead Indian Reservation 1993-1994. Salish Kootenai College, Pablo, MT. 55 pp.



WESTERN TOAD

Bufo boreas


State rank S3S4

Global rank G4

MTNHP status watch list

Element code AAABB01030

Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell



State range

Ranges throughout the region west of the continental divide. Also present east of the continental divide in the eastern front ranges and in the mountainous areas of the southwest and central parts of the state. Known to range up to approximately 2,900 meters in Montana (Black 1970).

Habitat description

Found in a wide variety of habitats including mountain wetlands, forests, woodlands, sagebrush, meadows and deserts (Nussbaum et al. 1983); probably a coniferous forest species in Montana (S. Corn, pers. comm.). Adult toads hibernate and shelter in self-made burrows and rodent burrows or under logs and rocks. Adults may move more than two kilometers away from water after breeding, and can remain away from surface water for relatively long periods of time (Koch and Peterson 1995). Breeding and subsequent egg deposition is dependent on temperature and snowmelt, and takes place from May to July in shallow areas of large and small lakes, beaver ponds, temporary ponds, slow-moving streams, and backwater channels of rivers (Black 1970). In the Yellowstone area, water chemistry at most breeding sites generally has a high pH (>8.0), high conductivity, and high acid-neutralizing capacity (Koch and Peterson 1995). Tadpoles commonly form dense aggregations in shallow warmer waters and metamorphose in mass at various times during their first summer (Nussbaum et al. 1983, Koch and Peterson 1995).

Model assumptions & caveats

Original intent was to base model on a 2 km maximum migration distance from terrestrial burrows to breeding sites. Although a 2 km buffer on hydrography is biologically reasonable, when applied to the state, nearly all areas are within the buffer, except for 1:100,000 DLG tiles where hydrography was mapped less densely (thus highlighting limitations of the layer). Also, the buffer became quite blocky, rather than curving with the hydrographic features, because it could only be built in raster format. To reduce such buffering artifacts and to minimize processing time, the buffer was eliminated. An elevation limit of 3,450 meters was selected to eliminate areas 250 meters above those the species is

known to inhabit in Wyoming (Baxter and Stone 1985). Initial review of the model showed high levels of accuracy in predicting the species' presence in areas where it is known to occur. However, the model probably overestimates the species' distribution. Recent declines make modeling problematic. Propagation of predicted habitat onto the prairies east of the continental divide is also a problem, but may be informative in some instances.

References

Baxter, G. T., and M. D. Stone. 1985. *Amphibians and reptiles of Wyoming*, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Black, J.H. 1970. Some aspects of the distribution, natural history and zoogeography of the toad genus *Bufo* in Montana. M.S. thesis, University of Montana. Missoula, MT. 70 pp.

Black, J.H. 1971. The toad genus *Bufo* in Montana. *Northwest Science* 45: 156-162.

Brunson, R.B. 1952. Recent collections of *Bufo boreas* from western Montana. *Proceedings of Montana Academy of Sciences* 11: 17-19.

Brunson, R. B., and H. A. Demaree, Jr. 1951. The herpetology of the Mission Mountains, Montana. *Copeia* 1951:306-308.

Corn, P.S., M.L. Jennings, and E. Muths. 1997. Survey and assessment of amphibian populations in Rocky Mountain National Park. *Northwest. Nat.* 78:34-55.

Hendricks, P., and J. D. Reichel. 1996. Amphibian and reptile survey of the Bitterroot National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 95 pp.

Koch, E. D., and C. R. Peterson. 1995. *Amphibians & reptiles of Yellowstone and Grand Teton national parks*. University of Utah Press. Salt Lake City, UT. 188 pp.

Marnell, L. E. 1997. Herpetofauna of Glacier National Park. *Northwest. Nat.* 78:17-33.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Reichel, J. D. 1995a. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.


Reichel, J. D. 1995b. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.

Reichel, J.D. 1996. Preliminary amphibian and reptile survey of the Helena National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 87 pp.

Russell, A. P., and A. M. Bauer. 1993. *The amphibians and reptiles of Alberta*. University of Calgary Press. Calgary, Alberta. 264 pp.

Werner, J.K. and J.D. Reichel. 1996. Amphibian and reptile monitoring/survey of the Kootenai National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 115 pp.

Werner, J.K. and T. Plummer. 1995. Amphibian and reptile survey of the Flathead Indian Reservation 1993-1994. Salish Kootenai College, Pablo, MT. 55 pp.



GREAT PLAINS TOAD

Bufo cognatus

State rank S3S4


Element code AAABB01050

Global rank G5

Modeled by M. Hart, B. Maxell, P. Hendricks, C.

MTNHP status watch list

Peterson, and J. Reichel; writeup by
B. Maxell



Predicted habitat: 17,566,226 ha, 46.13 % of state.

State range

Range unclear. Recorded from southeastern Alberta by Russell and Bauer (1993), north-central and southwest North Dakota by Wheeler and Wheeler (1966), and northeastern Wyoming by Baxter and Stone (1985). Range in Montana is likely to include much of the eastern plains. Known to range up to 1,220 meters in elevation in Montana (Black 1970).

Habitat description

Found in deserts, semidesert shrublands, grasslands, open floodplains and agricultural areas in stream valleys. In Montana they seem to prefer the higher elevations of short-grass prairies or undifferentiated grasslands, meadows within open stands of ponderosa pine, and areas near streams and irrigated lands (Black 1970, Bragg 1940). Adults hibernate and aestivate in self-excavated burrows during the long periods when terrestrial conditions aren't favorable (Wheeler and Wheeler 1966). Breeding takes place from May to July in clear, shallow temporary pools of flooded grasslands, ponds and reservoirs after suitable rains. Eggs are deposited in gelatinous strings on the pond bottom and are not attached to vegetation. Tadpoles metamorphose in about six weeks (Baxter and Stone 1985, Wheeler and Wheeler 1966).


Model assumptions & caveats

Original intent was to base model on a 2 km maximum migration distance from terrestrial burrows to breeding sites. Although a 2 km buffer on hydrography is biologically reasonable, when applied to the state, nearly all areas are within the buffer, except for 1:100,000 DLG tiles where hydrography was mapped less densely (thus highlighting limitations of the layer). Also, the buffer became quite blocky, rather than curving with the hydrographic features, because it could only be built in raster format. To reduce such buffering artifacts and to minimize processing time, the buffer was eliminated. An elevation limit of 1,500 meters was selected to eliminate areas 280 meters above that reported in Montana by Black (1970). The species' range and status is very unclear in the state. Not enough data points exist to assess model accuracy. However, the model probably overestimates the species'

distribution.

References


- Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.
- Black, J.H. 1970. Some aspects of the distribution, natural history and zoogeography of the toad genus *Bufo* in Montana. M.S. thesis, University of Montana. Missoula, MT. 70 pp.
- Black, J.H. 1971. The toad genus *Bufo* in Montana. *Northwest Science* 45: 156-162.
- Bragg, A.N. 1940. Observations on the ecology and natural history of Anura. I. Habits, habitat and breeding of *Bufo cognatus* Say. *Am. Nat.* 74: 322-349, 424-438.
- Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.
- Mosimann, J.E. and G.B. Rabb. 1952. The herpetology of Tiber Reservoir Area, Montana. *Copeia* 1952: 23-27.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Reichel, J. D. 1995a. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.
- Reichel, J. D. 1995b. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.
- Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.
- Wheeler, G.C. and J. Wheeler. 1966. The amphibians and reptiles of North Dakota. University of North Dakota, Grand Forks, ND. 104 pp.



CANADIAN TOAD

Bufo hemiophrys

State rank	S1	Element code	AAABB01080
Global rank	G4	Modeled by	M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell
MTNHP status	special concern		



State range

Range unclear. Recorded in eastern Alberta down to the Saskatchewan River by Russell and Bauer (1993) and in north-central North Dakota by Wheeler and Wheeler (1966). A subspecies of *Bufo hemiophrys*, the Wyoming toad, *Bufo hemiophrys baxteri*, exists as a small isolated population in the Laramie Basin of Wyoming (Baxter and Stone 1985). Range in Montana is likely to be limited to isolated remnant populations east of the Milk River and north of the Missouri River. Ranges up to 1,200 meters in Alberta (Russell and Bauer 1993).

Habitat description

Found in short-grass prairies, aspen parkland, boreal forest and floodplains (Baxter and Stone 1985, Russell and Bauer 1993). In Montana they seem to prefer ponds and marshes with relatively stable water levels surrounded by sedges and bulrushes in short-grass prairies and undifferentiated grasslands (Black 1970). Adults aestivate by burrowing into nearby friable soils and hibernate by burrowing into slight rises in the surrounding terrain (Tester and Breckenridge 1964). Breeding probably occurs with the first good rains from late April to early June and eggs are deposited in shallow waters of lakes, ponds, ditches, marshes, and temporary water bodies (Black 1970, Russell and Bauer 1993). Tadpoles usually transform by August (Baxter and Stone 1985).

Model assumptions & caveats

Model was based on a 500 m migration distance from terrestrial burrows to breeding sites (S. Corn, pers. comm.). An elevation limit was not included because the species is known to occupy habitats exceeding the elevations present within its range in Montana. The species' range and status is very unclear in the state. Not enough data points exist to assess model accuracy. However, the model probably overestimates the species' distribution.

References

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Black, J.H. 1970. Some aspects of the distribution, natural history and zoogeography of the toad genus *Bufo* in Montana. M.S. thesis, University of Montana. Missoula, MT. 70 pp.

Black, J.H. 1971. The toad genus *Bufo* in Montana. *Northwest Science* 45: 156-162.


Black, J.H., and A.N. Bragg. 1968. New additions to the herpetofauna of Montana. *Herpetologica* 24: 247.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.

Tester, J. R. and W. J. Breckenridge. 1964. Population dynamics of the Manitoba toad, *Bufo hemiophrys*, in northwestern Minnesota. *Ecology* 45: 592-601.

Wheeler, G.C. and J. Wheeler. 1966. The amphibians and reptiles of North Dakota. University of North Dakota, Grand Forks, ND. 104 pp.



PLAINS SPADEFOOT

Spea bombifrons


State rank S4?

Global rank G5

MTNHP status --

Element code AAABF02010

Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell



Predicted habitat: 26,537,934 ha, 69.69 % of state.

State range

Spea bombifrons = *Scaphiopus bombifrons*. Range in Montana is likely to include most of the eastern plains wherever burrowing is possible. May also be present in isolated localities west of the continental divide where observations have been reported, but no specimens collected. Although only documented by a few specimens and observations throughout eastern Montana, the species' range is fairly certain in the state due to records in surrounding states and provinces. Recorded from southeastern Alberta by Russell and Bauer (1993), all along the North Dakota border by Wheeler and Wheeler (1966), and in northeastern Wyoming by Baxter and Stone (1985). Known to range up to 1,830 meters in elevation in Wyoming (Baxter and Stone 1985).

Habitat description

Found in arid grasslands and sagebrush with sandy or loose soils; usually near permanent or temporary water bodies (Baxter and Stone 1985, Russell and Bauer 1993). Adults hibernate and aestivate in burrows excavated to depths of three feet in damp loose soils during the long periods when terrestrial conditions aren't favorable. Adults are only present on the surface on warm nights when the surface is damp (Wheeler and Wheeler 1966). Breeding takes place in warm, often muddy, temporary water bodies formed by substantial spring or summer rains. Eggs are deposited on submerged vegetation and hatch in about 48 hours. Tadpoles are carnivorous and frequently become cannibalistic before metamorphoses approximately 40 days after hatching (Wheeler and Wheeler 1966, Russell and Bauer 1993).


Model assumptions & caveats

Model was intended to be based on the availability of sandy, loose soils for burrowing (i.e. the exclusion of clay soils). However, the resolution of the soils layer was too coarse and was, therefore, excluded from the model. Instead, model was based on suitable cover types and elevation. An elevation limit of 1,950 meters was selected to eliminate areas 120 meters above those the species is known to inhabit in Wyoming (Baxter and Stone 1985). Not enough data points exist to assess model

accuracy. However, the model probably greatly overestimates the species' distribution due to its inability to exclude improper soil types. Propagation of predicted habitat west of the continental divide is problematic in several instances.

References

- Baxter, G. T., and M. D. Stone. 1985. *Amphibians and reptiles of Wyoming*, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.
- Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.
- Kellog, R. 1932. Notes on the spadefoot of the western plains (*Scaphiopus hammondi*). *Copeia* 1932(1): 36.
- Mosimann, J.E. and G.B. Rabb. 1952. The herpetology of Tiber Reservoir Area, Montana. *Copeia* 1952: 23-27.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Reichel, J. D. 1995a. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.
- Reichel, J. D. 1995b. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.
- Reichel, J.D. 1996. Preliminary amphibian and reptile survey of the Helena National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 87 pp.
- Russell, A. P., and A. M. Bauer. 1993. *The amphibians and reptiles of Alberta*. University of Calgary Press. Calgary, Alberta. 264 pp.
- Wheeler, G.C. and J. Wheeler. 1966. *The amphibians and reptiles of North Dakota*. University of North Dakota, Grand Forks, ND. 104 pp.



NORTHERN LEOPARD FROG

Rana pipiens


State rank S3S4

Global rank G5

MTNHP status special concern

Element code AAABH01170

Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell



State range

Occurred historically throughout Montana except for high mountain regions where species was mostly limited to lower valleys. Currently, nearly extirpated in western Montana, very localized in central Montana and abundant only in southeastern Montana (Reichel and Flath 1995). Ranges up to 2,743 meters in elevation in Wyoming (Baxter and Stone 1985).

Habitat description


Typically found in and adjacent to permanent slow-moving or standing water bodies with considerable vegetation. However, they may range widely into moist meadows, grassy woodlands and even agricultural areas (Nussbaum et al. 1983). In Montana adults are found primarily in riparian habitats or on the prairies near permanent waters without tall dense vegetation (Mosimann and Rabb 1952, Black 1969, Miller 1978). Adults hibernate underwater in permanent water bodies (Nussbaum et al. 1983). Breeding occurs from April to July, after individuals have migrated to preferred shallow water breeding sites from hibernation sites. Egg masses usually are deposited in March and April on vegetation or warm shallow pond bottoms exposed to sunlight (Corn and Livo 1989, Mosimann and Rabb 1952, Russell and Bauer 1993). Eggs hatch in 10-20 days and tadpoles metamorphose approximately two months later (Nussbaum et al. 1983).

Model assumptions & caveats

Model was based on the species' close dependence on permanent waters. An elevational limit of 2,250 meters was used to exclude high elevation areas (Chuck Peterson, pers. observ.). Initial review of the model showed high levels of accuracy in predicting the species presence in areas where it is currently known to occur. The species' distribution in western Montana is very much overestimated due to declines. Because the hydrography layer (assembled from USGS 1:100,000 digital line graphs) contains mapping inconsistencies among tiles, "squares" can be seen in the model output, especially in eastern Montana.

References


- Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.
- Black, J.H. 1969. The frog genus *Rana* in Montana. *Northwest Science* 43(4): 191-195.
- Brunson, R. B., and H. A. Demaree, Jr. 1951. The herpetology of the Mission Mountains, Montana. *Copeia* 1951:306-308.
- Corn, P.S. and J.C. Fogleman. 1984. Extinction of montane populations of the northern leopard frog (*Rana pipiens*) in Colorado. *J. Herpetol.* 18(2):147-152.
- Corn, P.S. and L.J. Livo. 1989. Leopard frog and wood frog reproduction in Colorado and Wyoming. *Northwest Nat.* 70:1-9.
- Corn, P.S., M.L. Jennings, and E. Muths. 1997. Survey and assessment of amphibian populations in Rocky Mountain National Park. *Northwest Nat.* 78:34-55.
- Dumas, P. C. 1964. Species-pair allopatry in the genera *Rana* and *Phrynosoma*. *Ecology* 45: 178-181.
- Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.
- Miller, J.D. 1978. Observations on the diets of *Rana pretiosa*, *Rana pipiens*, and *Bufo boreas* from western Montana. *Northwest Sci.* 52(3): 243-249.
- Mosimann, J.E. and G.B. Rabb. 1952. The herpetology of Tiber Reservoir Area, Montana. *Copeia* 1952: 23-27.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Reichel, J. D. 1995a. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.
- Reichel, J. D. 1995b. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.
- Reichel, J.D. 1996. Preliminary amphibian and reptile survey of the Helena National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 87 pp.
- Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.
- Werner, J.K. and J.D. Reichel. 1996. Amphibian and reptile monitoring/survey of the Kootenai National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 115 pp.
- Werner, J.K. and T. Plummer. 1995. Amphibian and reptile survey of the Flathead Indian Reservation 1993-1994. Salish Kootenai College, Pablo, MT. 55 pp.
- Wheeler, G.C. and J. Wheeler. 1966. The amphibians and reptiles of North Dakota. University of North Dakota, Grand Forks, ND. 104 pp.



WOOD FROG

Rana sylvatica

State rank	SR	Element code	AAABH01200
Global rank	G5	Modeled by	M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by B. Maxell
MTNHP status	special concern		



State range

No specimens recorded in Montana. However, the species has been reported in northern Idaho close to the Canadian and Montana border by Nussbaum et al. (1983), just north of Glacier National Park in Alberta by Russell and Bauer (1995), and in the Bighorn Mountains just south of the Montana border in Wyoming by Baxter and Stone (1985). The species may be present in the Bighorn Mountains in south-central part of the state, or in the northwest corner of the state. Ranges up to 8,200 meters elevation in Alberta (Russell and Bauer 1995).

Habitat description

Found along temporary ponds, lakes and stream shores, but adults also move into shaded portions of adjacent forests or brush where there is damp ground litter. Adults are largely terrestrial during the non-breeding season, but are usually not found far from water (Nussbaum et al. 1983). In Wyoming they were found only in shallow glacial kettle ponds without fish and most frogs were found in areas with extensive shallows and dense emergent sedges on the north side of the ponds (Garber 1992). Adults hibernate terrestrially in burrows, root channels and crevices. Adults may migrate up to 0.5 km to small pools, backwaters and beaver ponds which are used for breeding. Breeding takes place from March to June, and eggs are often deposited communally on emergent or submerged vegetation. Eggs hatch at different times depending on water temperatures and tadpoles metamorphose in about two months (Nussbaum et al. 1983).

Model assumptions & caveats


Modeled, but will not be included in final MT-GAP reporting. State rank SR -- reported, but lacking documentation that would provide a basis for accepting or rejecting observation.

Model based on species' dependence on hydrographic features and riparian areas. An elevation limit was not included because the species is known to occupy habitats exceeding the elevations present within its possible range in Montana. The model greatly overestimates the species' distribution.

Limiting the model to higher elevations (or to cover types at higher elevations), at least in south-central Montana, should be considered for the future.

References

- Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.
- Corkran, C.C., and C. Thoms. 1996. Amphibians of Oregon, Washington, and British Columbia. Lone Pine Publishing, Edmonton, Alberta. 175 pp.
- Corn, P.S. and L.J. Livo. 1989. Leopard frog and wood frog reproduction in Colorado and Wyoming. Northwest Nat. 70:1-9.
- Corn, P.S., M.L. Jennings, and E. Muths. 1997. Survey and assessment of amphibian populations in Rocky Mountain National Park. Northwest Nat. 78:34-55.
- Dunlap, D.G. 1977. Wood and western spotted frogs (Amphibia, Anura, Ranidae) in the Big Horn Mountains of Wyoming. Journal of Herpetology 11: 85-87.
- Garber, C. S. 1992. A survey for spotted frogs (*Rana pretiosa*), wood frogs (*Rana sylvatica*), and boreal toads (*Bufo boreas*) in Wyoming. Wyoming Natural Diversity Database, Laramie. 15 pp. + appendix.
- Nussbaum, R. A., E. D. Brodie, Jr., and R. M. Storm. 1983. Amphibians and reptiles of the Pacific Northwest. University of Idaho Press. Moscow, ID. 332 pp.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. Montana Outdoors 26(3):15-34.
- Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.
- Wheeler, G.C. and J. Wheeler. 1966. The amphibians and reptiles of North Dakota. University of North Dakota, Grand Forks, ND. 104 pp.




SNAPPING TURTLE

Chelydra serpentina

State rank S3
Global rank G5
MTNHP status special concern

Element code ARAAB01010
Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



State range

Throughout southeastern and far eastern Montana, although not common. Occurs in the Yellowstone River and major tributaries below Billings; probably in Missouri River and major tributaries below Fort Benton, although poorly documented in this drainage (Reichel and Flath 1995). Introduced locally west of the Continental Divide; status at these locations unknown.

Habitat description

Permanent freshwater bodies of nearly all kinds (ponds, lakes, streams, rivers), usually in shallow water; sometimes enters brackish coastal waterways and marshes. Typically in quiet or sluggish water with an abundance of aquatic plants and a soft mud or sand bottom or an abundance of submerged tree trunks and brush (Ernst et al. 1994). Hibernates in bottom mud or under submerged vegetation. Nests in soft soil, sometimes > 1.5 km from home range and overland > 0.5 km from water (Ernst et al. 1994).

In Montana: Almost any permanent and shallow fresh water with aquatic vegetation and muddy or sandy bottom and banks, although this is poorly documented. Nest habitat not described.

Model assumptions & caveats

Model based on hydrographic features and adjacent riparian habitats. An elevational limit was used to restrict predicted range around mountain ranges. Model may overestimate range by including lower order streams; would prefer to restrict model to \geq 3rd order streams.

References

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Ernst, C. H., R. W. Barbour, and J. E. Lovich. 1994. Turtles of the United States and Canada. Smithsonian Institution Press. Washington, D.C. 578 pp.


Hammer, D.A. 1969. Parameters of a marsh snapping turtle population, Lacreek Refuge, South Dakota. *J. Wildl. Manage.* 33(4):995-1005.

Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Reichel, J. D. 1995. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.

Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.




SPINY SOFTSHELL

Apalone spinifera

State rank S3
Global rank G5
MTNHP status special concern

Element code ARAAG01030
Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



State range

Apalone spinifera = *Trionyx spiniferus*. Eastern plains, mainly in the Missouri and Yellowstone rivers and their major tributaries (Reichel and Flath 1995); not yet documented below Fort Peck.

Habitat description

Primarily a riverine species of shallow well-oxygenated water, also inhabiting creeks, bayous, oxbows, lakes and reservoirs. Typically in water with a soft bottom and some aquatic vegetation; sandbars and mud flats often present, fallen trees with underwater limbs frequented (Ernst et al. 1994). Hibernates submerged in mud in the northern parts of its range. A highly aquatic turtle, with most nests located close to water in adjacent sand or gravel bars; rarely wanders as far as 100 m from water to nest (Ernst et al. 1994).

In Montana: Primarily in major rivers; in slower moving water with nearby mud flats and sandbars, sometimes in backwater sloughs (P. Hendricks, pers. observ.). Nest habitat not described.

Model assumptions & caveats

Model based on major rivers. An elevational limit was used to restrict predicted range around mountain ranges. Model may underestimate range slightly by excluding some lower stream orders.

References


- Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.
- Ernst, C. H., R. W. Barbour, and J. E. Lovich. 1994. Turtles of the United States and Canada. Smithsonian Institution Press. Washington, D.C. 578 pp.

Hammerson, G. A. 1982. Amphibians and reptiles in Colorado. Colorado Division of Wildlife. Denver, CO. 131 pp.

Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. Montana Outdoors 26(3):15-34.

Reichel, J. D. 1995. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.



SHORT-HORNED LIZARD

Phrynosoma douglasii


State rank S4

Global rank G5

MTNHP status --

Element code ARACF12030

Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



Predicted habitat: 17,966,952 ha, 47.18 % of state.

State range

Phrynosoma douglasii = *Phrynosoma hernandezii*. Plains east of the Continental Divide (Reichel and Flath 1995); most common in the southeast quarter of the state. Zamudio and others (1997) split short-horned lizards into two species, *P. hernandezii* and *P. douglasii*. Only *P. hernandezii* has been documented in Montana.

Habitat description

Occurs in a variety of habitats, including sagebrush, prairie grassland, open juniper and pine forest; up to 3300 m elevation. Substrate can vary from hardpan to sandy to rocky, but patches of loose soil for burrowing always present (Hammerson 1982, Nussbaum et al 1983). Gives birth to live young.

In Montana: Occurs in habitats with loose soils, including grasslands, sagebrush, open forests of ponderosa pine, rocky mountain juniper, limber pine / Utah juniper, up to at least 1600 m elevation (Hendricks and Reichel 1996, P. Hendricks pers. observ.).

Model assumptions & caveats

Model based on cover types; unable to use soil layer. An elevational limit of 2100 m, consistent with range in Wyoming (Baxter and Stone 1985), was used to restrict predicted range upward into mountains. Model overestimates range because of inability to use loose, sandy soils in model.

References

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Dumas, P. C. 1964. Species-pair allopatry in the genera *Rana* and *Phrynosoma*. *Ecology* 45: 178-181.


- Hammerson, G. A. 1982. Amphibians and reptiles in Colorado. Colorado Division of Wildlife. Denver, CO. 131 pp.
- Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.
- Mosimann, J.E. and G.B. Rabb. 1952. The herpetology of Tiber Reservoir Area, Montana. *Copeia* 1952: 23-27.
- Nussbaum, R. A., E. D. Brodie, Jr., and R. M. Storm. 1983. Amphibians and reptiles of the Pacific Northwest. University of Idaho Press. Moscow, ID. 332 pp.
- Powell, G. L., and A. P. Russell. 1991. Distribution of the eastern short-horned lizard (*Phrynosoma douglasi brevirostre*) in Alberta, Canada. *Northwest. Nat.* 72:21-26.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.
- Werner, J.K. 1974. *Phrynosoma douglasi brevirostre*. *Herpetological Review* 5(1):20.
- Zamudio, K.R., K.B. Jones, and R.H. Ward. 1997. Molecular systematics of short-horned lizards: biogeography and taxonomy of a widespread species complex. *Systematic Biology* 46:284-305.

SAGEBRUSH LIZARD

Sceloporus graciosus

State rank S3S4
Global rank G5
MTNHP status watch list

Element code ARACF14030
Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



Predicted habitat: 12,434,258 ha, 32.65 % of state.



State range

East of Continental Divide, mostly in southern half of eastern Montana (Reichel and Flath 1995).

Habitat description

Sagebrush plains, greasewood and other desert shrubs, open juniper and pine woodlands with brushy cover, up to 3200 m elevation (Nussbaum et al. 1983), also geothermal areas in Yellowstone National Park (Koch and Peterson 1995). Eggs (typically 3-4) buried in clutches in loose soil, usually at the base of a shrub (Nussbaum et al. 1983).

In Montana: Sagebrush plains, open pine/juniper woodland with brushy cover and small rock outcrops (Koch and Peterson 1995, Hendricks and Reichel 1996); up to 2200 m in Yellowstone National Park, but usually below 1700 m elevation elsewhere (P. Hendricks, pers. observ.). Nest sites include loose soil at base of shrubs (Algard 1968).

Model assumptions & caveats

Model based on cover types; unable to use soil layer. An elevational limit of 2250 m, consistent with range in Wyoming (Baxter and Stone 1985), was used to restrict predicted range upward into mountains. Model overestimates range because of inability to use loose, sandy soils in model.

References

Algard, G. A. 1968. Distribution, temperature and population studies of *Sceloporus graciosus* in Yellowstone National Park. M.S. thesis, Montana State University. Bozeman, MT. 34 pp.

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.

Koch, E. D., and C. R. Peterson. 1995. Amphibians & reptiles of Yellowstone and Grand Teton national parks. University of Utah Press. Salt Lake City, UT. 188 pp.

Mueller, C.F. 1967. Temperature and energy characteristics of the sagebrush lizard in Yellowstone National Park. Ph.D. dissertation, Montana State University, Bozeman. 38 pp.


Mueller, C.F. 1969. Temperature and energy characteristics of the sagebrush lizard (*Sceloporus graciosus*) in Yellowstone National Park. *Copeia* 1969:153-160.

Mueller, C. F., and R. E. Moore. 1969. Growth of the sagebrush lizard, *Sceloporus graciosus*, in Yellowstone National Park. *Herpetologica* 25:35-38.

Nussbaum, R. A., E. D. Brodie, Jr., and R. M. Storm. 1983. Amphibians and reptiles of the Pacific Northwest. University of Idaho Press. Moscow, ID. 332 pp.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Reichel, J. D. 1995. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.



WESTERN SKINK

Eumeces skiltonianus

State rank S3S4


Element code ARACH01110

Global rank G5

Modeled by M. Hart, B. Maxell, P. Hendricks, C.

MTNHP status watch list

Peterson, and J. Reichel; writeup by P. Hendricks



State range

West of the Continental Divide; near the Idaho border in the Clark Fork River and Kootenai River drainages in northwestern Montana (Reichel and Flath 1995).

Habitat description

Grassy slopes in oak woodland and mesic conifer forest, pine and juniper woodland, rocky canyons and rimrock up to 2150 m elevation; under rotting wood, surface litter and large flat stones (Nussbaum et al. 1983, Storm and Leonard 1995). Eggs (2-6) are laid in clutches in burrows or cavities under logs and stones (Nussbaum et al. 1983, Storm and Leonard 1995).

In Montana: Grassland low shrub-covered slopes in open ponderosa pine, Douglas-fir and juniper woodlands, rocky outcrops and talus; known up to 1750 m elevation (Hendricks and Reichel 1996, P. Hendricks pers. observ.). Nest habitat not described.

Model assumptions & caveats

Model based on cover types. An elevational limit of 2100 m, consistent with other areas in the Pacific Northwest (Nussbaum et al 1983), was used to restrict predicted range upward into mountains. Model overestimates range due to a lack of good information on habitat associations.

References

Hendricks, P., and J. D. Reichel. 1996. Amphibian and reptile survey of the Bitterroot National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 95 pp.


Nussbaum, R. A., E. D. Brodie, Jr., and R. M. Storm. 1983. Amphibians and reptiles of the Pacific Northwest. University of Idaho Press. Moscow, ID. 332 pp.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Rodgers, T. L., and W. L. Jellison. 1942. A collection of amphibians and reptiles from western Montana. *Copeia* 1942:10-13.

Storm, R. M., and W. P. Leonard (eds.). 1995. *Reptiles of Washington and Oregon*. Seattle Audubon Society. Seattle, WA. 176 pp.

Werner, J.K. and J.D. Reichel. 1996. *Amphibian and reptile monitoring/survey of the Kootenai National Forest: 1995*. Montana Natural Heritage Program. Helena, MT. 115 pp.



WESTERN HOGNOSE SNAKE

Heterodon nasicus


State rank S3?

Element code ARADB17010

Global rank G5

Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks

MTNHP status special concern



Predicted habitat: 19,836,672 ha, 52.09 % of state.

State range

Plains of eastern Montana; western limits not well-known (Reichel and Flath 1995).

Habitat description

Sandy or gravelly soils in prairie, sandhills, open woodland, floodplains, semidesert, mountain canyon bottoms, badlands; near water (Hammerson 1982, Baxter and Stone 1985, Russell and Bauer 1993). Eggs laid in sandy soil.

In Montana: Arid areas, prairie grasslands and shrublands, floodplains with gravelly or sandy soils (Reichel and Flath 1995). Nest and den sites not described.

Model assumptions & caveats

Model based on cover types and riparian habitat. An upper elevational limit of 1650 m was used to prevent predicted range upward into mountains; this limit may be too high. The model overestimates habitat because a loose, sandy soil layer was not available at sufficiently fine resolution.

References

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Hammerson, G. A. 1982. Amphibians and reptiles in Colorado. Colorado Division of Wildlife. Denver, CO. 131 pp.

Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.


Mosimann, J.É. and G.B. Rabb. 1952. The herpetology of Tiber Reservoir Area, Montana. *Copeia* 1952: 23-27.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Reichel, J. D. 1995a. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.

Reichel, J. D. 1995b. Preliminary amphibian and reptile survey of the Sioux District of the Custer National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 75 pp.

Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.




MILK SNAKE

Lampropeltis triangulum

State rank S2
Global rank G5
MTNHP status special concern

Element code ARADB19050
Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



State range

Southeastern Montana and along the Missouri River; may be localized (Reichel and Flath 1995).

Habitat description

Grasslands, sandhills, open ponderosa pine and juniper woodlands, arid river valleys, often associated with rock outcrops and scarps; usually below 2450 m elevation (Hammerson 1982, Baxter and Stone 1985). Dens in rock outcrops (Hammerson 1982).

In Montana: Sandstone bluffs, rock outcrops, grasslands, open ponderosa pine and juniper stands (Hendricks and Reichel 1996); den sites undescribed, habitat associations poorly known because of few observations.

Model assumptions & caveats

Model based on cover types. An upper elevational limit of 1950 m, based on range in Wyoming (Baxter and Stone 1985), was used to restrict predicted range upward into mountains. Model grossly overestimates suitable habitat; would have preferred to base model on sandstone outcrops and hogback rims. Predicted habitat extrapolated from known range into southwestern Montana may be problematic.

References

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Hammerson, G. A. 1982. Amphibians and reptiles in Colorado. Colorado Division of Wildlife. Denver, CO. 131 pp.


Hendricks, P., and J. D. Reichel. 1996. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.

Nelson, D.J. 1948. *Lampropeltis triangulum gentilis* in Montana. *Herpetologica* 4:170.

Nelson, D.J. 1950. *Lampropeltis triangulum gentilis* in Montana. *Herpetologica* 6:41.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.

Williams, K.L. 1989. Systematics and natural history of the American milksnake, *Lampropeltis triangulum*. Second revised edition. 192 pp.




COMMON GARTER SNAKE

Thamnophis sirtalis

State rank S4
Global rank G5
MTNHP status --

Element code ARADB36130
Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



State range

Throughout Montana (Reichel and Flath 1995).

Habitat description

Prairie grassland, shrubland, coniferous woodland, farms and urban areas, usually associated with permanent water (lakes, streams, marshes, wet meadows); below 2100 m elevation (Russell and Bauer 1993, Koch and Peterson 1995, Storm and Leonard 1995). Dens in rocky sites near water; give birth to live young. May disperse many miles from dens (Storm and Leonard 1995).

In Montana: Coniferous (often ponderosa pine and Douglas-fir) forest, broadleaf woodlands, shrubland and grassland, urban areas, usually near water; recorded to 1700 m elevation (Werner and Reichel 1994, Reichel 1995a, Hendricks and Reichel 1996a, Hendricks and Reichel 1996b). Den sites not described.


Model assumptions & caveats

Model based on cover types and hydrography, using a 500 m buffer into selected cover types. An upper elevational limit of 2250 m was used, based on Koch and Peterson (1995). Model may underestimate suitable habitat; there is some biological justification for increasing the buffer distance, but most occurrences are near water. Elevational limit may be too high in the northwestern mountains. Because the hydrography layer (assembled from USGS 1:100,000 digital line graphs) contains mapping inconsistencies among tiles, "squares" can be seen in the model output, especially in eastern Montana.

References

Anderson, M.E. 1977. Aspects of the ecology of two sympatric species of *Thamnophis* and heavy metal accumulation with the species. M.S. thesis, University of Montana, Missoula. 147 pp.

- Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.
- Brunson, R. B., and H. A. Demaree, Jr. 1951. The herpetology of the Mission Mountains, Montana. *Copeia* 1951:306-308.
- Hendricks, P., and J. D. Reichel. 1996a. Preliminary amphibian and reptile survey of the Ashland District, Custer National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 79 pp.
- Hendricks, P., and J. D. Reichel. 1996b. Amphibian and reptile survey of the Bitterroot National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 95 pp.
- Koch, E. D., and C. R. Peterson. 1995. Amphibians & reptiles of Yellowstone and Grand Teton national parks. University of Utah Press. Salt Lake City, UT. 188 pp.
- Marnell, L. E. 1997. Herpetofauna of Glacier National Park. *Northwest. Nat.* 78:17-33.
- Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. *Montana Outdoors* 26(3):15-34.
- Reichel, J. D. 1995. Preliminary amphibian and reptile survey of the Lewis and Clark National Forest: 1994. Montana Natural Heritage Program. Helena, MT. 92 pp.
- Reichel, J.D. 1996. Preliminary amphibian and reptile survey of the Helena National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 87 pp.
- Rodgers, T. L., and W. L. Jellison. 1942. A collection of amphibians and reptiles from western Montana. *Copeia* 1942:10-13.
- Russell, A. P., and A. M. Bauer. 1993. The amphibians and reptiles of Alberta. University of Calgary Press. Calgary, Alberta. 264 pp.
- Storm, R. M., and W. P. Leonard (eds.). 1995. Reptiles of Washington and Oregon. Seattle Audubon Society. Seattle, WA. 176 pp.
- Werner, J.K. and J.D. Reichel. 1996. Amphibian and reptile monitoring/survey of the Kootenai National Forest: 1995. Montana Natural Heritage Program. Helena, MT. 115 pp.
- Werner, J.K. and T. Plummer. 1995. Amphibian and reptile survey of the Flathead Indian Reservation 1993-1994. Salish Kootenai College, Pablo, MT. 55 pp.




SMOOTH GREEN SNAKE

Liochlorophis vernalis

State rank S2S3
Global rank G5
MTNHP status special concern

Element code ARADB47010
Modeled by M. Hart, B. Maxell, P. Hendricks, C. Peterson, and J. Reichel; writeup by P. Hendricks



State range

Liochlorophis vernalis = *Opheodrys vernalis*. Far northeastern corner of the state (Sheridan and Roosevelt counties); distribution based on very few observations (Reichel and Flath 1995).

Habitat description

Riparian vegetation, often with dense herbaceous cover, mountain shrubland, woodland meadows, drier habitats adjacent to riparian areas; usually below 2750 m elevation (Hammerson 1982, Baxter and Stone 1985). Dens under rocks or wood, or in rotting logs.

In Montana: Riparian vegetation, irrigated lawns in rural communities (T. Nordhagen, pers. comm. to P. Hendricks); habitat associations poorly known (Reichel and Flath 1995). Dens undescribed.

Model assumptions & caveats

Model based on cover types, riparian habitats and hydrography; hydrography with a 500 m buffer into selected cover types to accommodate dispersal. No elevational limit seemed necessary. Model appears reasonable, although may overestimate range because of propagation of predicted habitat along major rivers.

References

Baxter, G. T., and M. D. Stone. 1985. Amphibians and reptiles of Wyoming, second edition. Wyoming Game and Fish Department. Cheyenne, WY. 137 pp.

Black, J.H., and A.N. Bragg. 1968. New additions to the herpetofauna of Montana. *Herpetologica* 24: 247.

Grobman, A. 1992. On races, clines, and common names in *Opheodrys*. *Herpetological Review* 23:14-15.


Hammerson, G. A. 1982. Amphibians and reptiles in Colorado. Colorado Division of Wildlife. Denver, CO. 131 pp.

Livo, L.J., D. Chiszar, and H.M. Smith. 1996. *Liochlorophis* (=Opheodrys) *vernalis* (smooth green snake). Herpetol. Rev. 27(3):154.

Reichel, J. and D. Flath. 1995. Identification of Montana's amphibians and reptiles. Montana Outdoors 26(3):15-34.

Stebbins, R.C. 1985. A field guide to western reptiles and amphibians. Houghton Mifflin Company, Boston. 336 pp.

Wheeler, G.C. and J. Wheeler. 1966. The amphibians and reptiles of North Dakota. University of North Dakota, Grand Forks, ND. 104 pp.



COMMON LOON

Gavia immer


State rank S1S2B,SZN

Element code ABNBA01030

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Breeding range primarily restricted to low elevation (below ~1600m) glacial lakes (morainal, kettle) in the northwest corner of the state; smaller populations possibly breed locally in southcentral and eastern (Nelson Reservoir) Montana, but most records are from the early 1900's (Dolan 1994, Montana Bird Distribution Committee 1996). Transient sightings occur throughout the state; occasional overwintering also in northwest and possibly in southcentral and eastern MT breeding areas (Montana Bird Distribution Committee 1996).

Habitat description

Breeding habitat generally limited to glacial lakes within tundra and coniferous regions, although occasionally breed on northern prairies (Palmer 1962, Fitch *in* Clark et al. 1989, Dolan 1994). Larger, clear lakes with partially forested shorelines, varying depths, numerous islands, highly irregular shorelines that provide areas protected from winds and rough water, and that are free from human disturbance represent optimal breeding habitat (Palmer 1962, Johnsgard 1979, McIntyre 1988, Fitch *in* Clark et al. 1989, Heglund et al. 1994, McIntyre 1994). Avoid lakes with fluctuating water levels (dammed), moderate to high turbidity, unprotected bays and inlets, and entirely barren shorelines (Fitch *in* Clark et al. 1989, Heglund et al. 1994, McIntyre 1994). Water clarity is critical; in Ontario, abandoned breeding on lakes after industrial pollution caused high turbidity levels (Barr 1986). Typically, a 40-80ha lake is required to support one breeding pair, but a 20ha lake with irregular shoreline will suffice (McIntyre 1988, Kerekes et al. 1994). Nests built of mud and vegetation; always within 1-2m of water, preferably on islands, but muskrat houses, floating vegetation mats, rocks and shorelines also used (McIntyre 1983, Ehrlich et al. 1988, Fitch *in* Clark et al. 1989, Dolan 1994, McIntyre 1994, Ruggles 1994). In Alberta, preferentially nest on islands or muskrat houses (Salt and Salt 1976). Brood areas in shallower, protected bays or inlets of lakes where emergent vegetation provides cover and foraging habitat (McIntyre 1988, Ehrlich et al. 1988, McIntyre 1994). In the Turtle Mountains of ND, breeding restricted to larger, permanent, freshwater lakes with abundant fish

populations, bordered by extensive stands of woodland (Quaking Aspen), and with peripheral zones of deep-marsh emergent vegetation (Stewart 1975). In MT, lakes must be greater than 10ha, deep enough to avoid freezing solid in winter (to sustain fish food base), and be below ~1600m in elevation (Skaar 1989, Kelley 1992a). Small peat islands in marshy areas and shallow bays are typical nest sites on MT lakes (Kelley 1992a, Dolan 1994). Subsist primarily on fish (80% of the diet); also crustaceans, amphibians, insects, vegetation (DeGraaf et al. 1991). During migration, inland and coastal waters are used; may gather on larger lakes than used for breeding before migrating to wintering areas (Palmer 1962, Dolan 1994). Inland migration disperses widely throughout the U.S., on relatively large, obstruction free expanses of water (Palmer 1962, DeGraaf et al. 1991); Twin Lakes Reservoir, ID is an important staging area for the Rocky Mountain population (Fitch in Clark et al. 1989). Unknown where Northern Rocky Mountain populations winter - breeders of the east side of the Divide likely migrate to Gulf of Mexico; others winter on the Pacific Coast (Dolan 1994). Winter habitat primarily marine - coastal waters, bays, estuaries; also large freshwater lakes and reservoirs that remain ice free, usually near coasts (Palmer 1962, Dolan 1994). On nesting and staging lakes, sensitive to human disturbances (especially from boating), and water level fluctuations (Fitch in Clark et al. 1989). Globally, populations are secure, however unregulated hunting, chemical contamination, effects of acid rain, oil spills on wintering grounds, and increased human development and activity on/near breeding lakes continue to reduce the breeding range (Fitch in Clark et al. 1989, Dolan 1994). In Alberta, have been driven out of most deepwater breeding lakes by human recreation; only adult nonbreeders seen on these lakes, as breeders are locally restricted to more northern, remote areas (Salt and Salt 1976). In the Greater Yellowstone Ecosystem, many lakes ideal for nesting are not used because human disturbance is high (Fitch in Clark et al. 1989). Additionally, at higher elevations, the shorter ice-free season makes timing of nesting and chick rearing critical; therefore are particularly vulnerable to human disturbances (Fitch in Clark et al. 1989). Local declines are suspected within the MT breeding range due to habitat loss (shoreline development, dam construction), and lowered reproductive success induced by human disturbance/recreation (Dolan 1994). The Common Loon is a species of special concern in ID and MT, is considered "protected non-game" by MT Fish Wildlife & Parks, and is a priority 1 species in WY (Fitch in Clark et al. 1989, Dolan 1994).


Model assumptions & caveats

Breeding habitat modeled. Rare winter resident; primary winter range/habitat is coastal, therefore not included in model. Common migrant; preferred migration habitat used in MT assumed defined within breeding habitat parameters. Potential breeding habitat in prairie regions of MT not specified in model, but somewhat defined within breeding habitat parameters. Lake depth, clarity, island characteristics, and degree of human disturbance information not available to include in model. Assume adequate nest sites (lake islands, peat islands, muskrat houses) and brood rearing areas available within suitable habitat. Hydrography coverage may not be consistently accurate/true to mapping scale used. Most MT lakes are simple in shape, lacking highly irregular shorelines that provide marshy inlets/bays preferred for nesting - could not accurately resolve shoreline morphology at mapping scale used. Habitat associations (i.e., lakes with forest and emergent vegetation), could not be modeled at mapping scale used (must query lakes with forest or emergent vegetation). Potential for human disturbance in predicted habitat areas cannot be assessed at mapping scale used. Suitable breeding habitat overestimated.

References

- Barr, J.F. 1986. Population dynamics of the Common Loon (*Gavia immer*) associated with mercury contaminated waters in northwestern Ontario. Canadian Wildl. Serv. Occ. Paper no. 56. 25 pp.
- Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.

- Dolan, P. M. 1994. The Common loon (*Gavia immer*) in the northern region: biology and management recommendations. Unpublished report. U.S.D.A. Forest Service Region 1. 76 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Heglund, P.S., J.R. Jones, L.H. Fredrickson, and M.S. Kaiser. 1994. Use of boreal forested wetlands by Pacific Loons (*Gavia immer*, Lawrence) and Horned Grebes (*Podiceps auritus*, L.): relations with limnological characteristics. *Hydrobiologia* 279/280: 171-183.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Kelley, L.M. 1992a. The effects of human disturbance on Common Loon productivity in northwestern Montana. M.S. thesis, Montana State University, Bozeman. 65 pp.
- Kerekes, J., R. Tordon, A. Nieuwburg, and L. Risk. 1994. Fish eating bird abundance in oligotrophic lakes in Kejimikujik National Park, Nova Scotia, Canada. *Hydrobiologia* 279/280: 57-61.
- McIntyre, J. W. 1983. Nurseries: A Consideration of Habitat Requirements During the Early Chick-Rearing Period in Common Loons. *Journal of Field Ornithology* 54(3):247-253.
- McIntyre, J. W. 1988. The common loon: spirit of northern lakes. Univ. Minnesota Press, Minneapolis. x + 200 pp.
- McIntyre, J.W. 1994. Loons in freshwater lakes. *Hydrobiologia* 279/280:393-413.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Palmer, R. S. 1962. Handbook of North American birds. Vol. 1. Loons through flamingos. Yale University Press, New Haven. 567 pp.
- Ruggles, A.K. 1994. Habitat selection by loons in southcentral Alaska. *Hydrobiologia* 279/280: 421-430.
- Salt, W.R. and J.R. Salt. 1976. The birds of Alberta. Hurtig Publishers, Edmonton, Alberta. xv + 498 pp.
- Skaar, D. 1989. Montana Common Loon Management Plan. USDA-Forest Service, Region 1, Missoula, MT.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



CLARK'S GREBE

Aechmophorus clarkii


State rank S2S4B,SZN

Element code ABNCA04020

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Breed very locally in Montana; known or suspected to breed in Ninepipe NWR in w Montana, and Lake Helena, Canyon Ferry Reservoir, Freezout Lake WMA, and Bowdoin NWR in e Montana (Montana Bird Distribution Committee 1996). Transient sightings recorded for nw/cent, scen, and ne Montana; no confirmed or suspected overwintering recorded (Montana Bird Distribution Committee 1996).

Habitat description

Please see Western Grebe habitat description. Similar habitat requirements as for Western Grebes, differing only in distribution - Clark's and Western Grebes sympatric over entire range; Clark's more common in southern part of range, but numbers increasing in the north; has uncommon status in MT (Ehrlich et al. 1988, Wassink 1991, Wright 1996). Differences in breeding ecology between Western and Clark's Grebes not well studied/defined. All breeding reported for MT is on very large lakes and reservoirs - nesting on Lake Helena is near mid-lake on subsurface, aquatic vegetation mat (MNHP-VCA 1996). Nest colonially in large marshes and lakes near emergent vegetation; tend to forage farther from shore and in deeper water than Western Grebes (Ehrlich et al. 1988, Storer and Nuechterlain 1992). Winter on coastal waters, bays, and less frequently inland (Storer and Nuechterlain 1992).

Model assumptions & caveats

Breeding habitat modeled. Rare breeder, uncommon migrant; does not overwinter in the state. Migration habitat similar to that of breeding and is assumed is defined within breeding habitat parameters. Please see assumptions/caveats for Western Grebe. Since only known breeding sites within the state are on very large lakes and reservoirs, these hydrographic limitations were included in model. Suitable breeding habitat may be overestimated, however, limited information regarding breeding requirements, and difficulty in distinguishing this species from closely related Western

Grebe, may have underestimated state range, making habitat modeling difficult.

References


Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds*. Simon and Shuster, Inc., New York. xxx + 785 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Storer, R.W. and G.L. Nuechterlein. 1992. Western Grebe (*Aechmophorus occidentalis*), in *The Birds of North America* No. 26. A. Poole and F. Gill, eds. The Academy of Natural Sciences, Philadelphia, and The American Ornithologists Union, Washington, D.C.

Wassink, J. 1991. *Birds of the Central Rockies*. Mountain Press Publishing Co., Missoula, MT.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



AMERICAN WHITE PELICAN

Pelecanus erythrorhynchos


State rank S2B,SZN

Element code ABNFC01010

Global rank G3

Modeled by W. Williams

MTNHP status special concern



Predicted habitat: 32,451 ha, 0.09 % of state.

State range

Only four breeding sites known, in eastern Montana, but breeding birds may move 100 miles to forage, so are widely seen (Montana Bird Distribution Committee 1996). Successful colonies breed at Medicine Lake NWR, Bowdoin NWR, Canyon Ferry Reservoir, Arrod Lake, and Charles M. Russell NWR (Sidle et al. 1985, MNHP-VCA 1996). Widely distributed in migration; transient sightings recorded throughout the state, save extreme nw; overwintering does not occur in Montana (Palmer 1962, Montana Bird Distribution Committee 1996).

Habitat description

Breed mainly on isolated, sparsely vegetated islands in large freshwater lakes and reservoirs, inland rivers, and bays (Stewart 1975, Johnsgard 1979, Ehrlich et al. 1988, Evans and Knopf 1993). Prefer flat to gently sloped, loosely earthed islands free of obstructions that may interfere with flight access or visibility (Palmer 1962, Johnsgard 1979, Evans and Knopf 1993). Does nest at mainland (peninsula) sites but success is low as nests are vulnerable to mammalian predators (Findholt and Diem 1988, Hart in Clark et al. 1989). In 1993, low water levels at Alkali Lake, MT allowed cattle and coyote access to a breeding island; to date the island still has not been used (MNHP-VCA 1996). Breeding islands commonly 50+km from feeding areas; adults may travel 600km round trip to forage (Hart in Clark et al. 1989, Wassink 1991, Evans and Knopf 1993). Colonies often associated with Double-crested Cormorants; in the Prairie Pothole Region of ND Ring-billed Gulls and California Gulls are breeding associates as well (Sidle and Ferguson 1984). Nest on the ground; prefer loose soil or sand mounded with shells/stones to create a shallow depression with a low rim (Johnsgard 1979, Sidle and Ferguson 1984, Evans and Knopf 1993). Will nest adjacent to but not in available herbaceous or shrubby vegetation, or along lines of driftwood; avoid nesting on gravel substrates (Knopf 1979, Sidle and Ferguson 1984, Evans and Knopf 1993). In forested regions, occasionally nest under trees (poplars, spruce, fir, willows), but must then walk from nest to the forest edge to take flight (Evans and Knopf 1993). Diet consists mainly of fish, crayfish, salamanders; taken from shallow water (0.3-2.5m deep), open areas within marshes, along lakes, along river edges, or on or below rapids (Ehrlich et al. 1988,

Evans and Knopf 1993). Migration is inland, along river valleys, over deserts and over mountains, however the Rocky Mountains are generally not crossed; populations remain e or w of Continental Divide and migrate primarily south (Evans and Knopf 1993). Foraging/loafing stopovers in areas similar to breeding habitat; also including brackish and marine shallow bays and inlets (Palmer 1962, Evans and Knopf 1993). The American White Pelican western population has been identified by the U.S. Fish and Wildlife Service as "sensitive" and is listed as a "threatened, endangered, or sensitive species" in MT (Findholt and Diem 1988, Montana Bird Distribution Committee 1996). Highly sensitive to human disturbance; intrusions into the breeding colony will cause desertion, consequent nest destruction and increased predation by gulls and eagles (Findholt and Diem 1988, Evans and Knopf 1993). A 400m buffer around nest islands recommended to reduce human perturbations; foraging lakes should be protected as well (Findholt and Diem 1988, Hart in Clark et al. 1989, Evans and Knopf 1993).

Model assumptions & caveats

Breeding habitat modeled. Does not overwinter in the state. Common migrant; migration/stopover habitat similar to that of breeding, therefore assumed defined within breeding habitat parameters. Breeds on 'large' lakes, but minimum lake size not reported in literature - so that suitable habitat is not grossly overestimated, minimum lake size limitation set to sizes of smaller, known breeding lakes in MT. Island characteristics information not available; Hydrography and island coverage may not be consistently accurate/true to mapping scale used. Potential foraging areas not included in model, therefore model focused on suitable nesting habitat. Potential for/degree of human disturbance information not available. Suitable nesting habitat defined by known colony locations, however still may be overestimated. Foraging habitat, however, is likely underestimated.

References

Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.

Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.

Evans, R. M. and F. L. Knopf. 1993. American White Pelican (*Pelecanus erythrorhynchos*). In: A. Poole and F. Gill, (eds.), *The Birds of North America*, No. 57. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union.

Findholt, S. L. and K. L. Diem. 1988. Status and distribution of American white pelican nesting colonies in Wyoming: an update. *Great Basin Nat.* 48:285-289.

Johnsgard, P.A. 1979. *Birds of the Great Plains: breeding species and their distribution*. Univ. Nebraska Press, Lincoln. 539 pp.

Knopf, F.L. 1979. Spatial and temporal aspects of colonial nesting of white pelicans. *Condor* 81: 353-363.

Montana Bird Distribution Committee. 1996. P. D. Skaar's *Montana Bird Distribution*, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.


Palmer, R. S. 1962. *Handbook of North American birds*. Vol. 1. Loons through flamingos. Yale University Press, New Haven. 567 pp.

Sidle, J.G. and E.L. Ferguson. 1984. White Pelican populations at Chase Lake, North Dakota, evaluated by aerial photography. *Prairie Nat.* 16: 13-26.

Sidle, J.G., W.H. Koonz and K. Roney. 1985. Status of the American White Pelican: an update. *Am. Birds* 39(5): 859-864.

Stewart, R.E. 1975. *Breeding birds of North Dakota*. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.

Wassink, J. 1991. Birds of the Central Rockies. Mountain Press Publishing Co., Missoula, MT.



GREAT BLUE HERON

Ardea herodias


State rank S4B,SZN

Element code ABNGA04010

Global rank G5

Modeled by W. Williams

MTNHP status watch list



Predicted habitat: 461,676 ha, 1.21 % of state.

State range

Breed throughout Montana; transient sightings widely scattered; overwintering documented locally in western Montana (Montana Bird Distribution Committee 1996).

Habitat description

Widespread and adaptable; breed in a variety of aquatic habitats, usually where there are trees: freshwater and brackish marshes, swamps, river and lake edges, mangroves, tidal flats (Palmer 1962, Johnsgard 1979, Hancock and Kushlan 1984, Butler 1992). Nest sites often chosen to avoid mammalian predators and are fairly isolated on islands, trees in swamps, on high branches (Johnsgard 1979, Butler 1992). Nesting usually colonial in deciduous or conifer trees surrounded by water to 1,100m elevation; occasionally in shrubs; rarely ground or cliff (Hancock and Kushlan 1984, Ehrlich et al. 1988, Butler 1992). Colony location directly related to the distribution of foraging habitats (Butler 1992). In ME, colonies commonly form near wetlands as well as centrally within a wetlands complex (Gibbs 1991). In SD, typically nest in riparian woodlands (Dowd and Flake 1985); in eastern OR, cottonwoods frequently used (Henny and Kurtz 1978); in ID prefer willows or cottonwoods (Burleigh 1978). In nw MT, primarily nest in largest available cottonwoods of riparian areas; also conifers in drier sites (MNHP-VCA 1996). Colonies often nest in association with Double-crested Cormorants in the Great Plains, especially where reservoirs or waterfowl management impoundments have flooded areas of tall trees (Stewart 1975, Johnsgard 1979). Flooded tree nesting habitat somewhat temporary, as dead nest trees decay and eventually fall (Stewart 1975). Opportunistic forager: primarily fish; also small mammals, nestlings, amphibians, reptiles, invertebrates (Ehrlich et al. 1988, Butler 1992). Feeding habitat includes breeding areas; also mudflats, sandbars, wet meadows, pastures, dry fields, near emergent vegetation (Palmer 1962, Ehrlich et al. 1988, Butler 1992). Migration habitat is similar to that of breeding; in winter favor natural wetlands, river banks, estuaries, grasslands, coasts (Butler 1992). The Great Blue Heron was Blue-Listed 1980-81 and is considered an "SI" species in Montana, although overall, populations are apparently stable or increasing (Ehrlich et al. 1988, Montana Bird


Distribution Committee 1996). Are sensitive to human disturbance, road building, and logging activity within 0.5km of nesting colonies, and will abandon nests when perturbed (Butler 1992). Wetland loss, however, may have contributed more to population declines than human disturbance (Butler 1992).

Model assumptions & caveats

Breeding habitat modeled. Common migrant; uncommon winter resident. Migration habitat is similar to that of breeding and assumed defined within breeding habitat parameters. Winter habitat in MT not part of typical range and is excluded from model - winter survivability would better define suitable winter habitat in the state, but is unknown. Hydrography coverage not consistently accurate/true to mapping scale used. Cannot model elements of foraging habitat (indicator of suitable nesting habitat). Cannot determine extent of standing water in treed habitat at mapping scale used. Tree age/size class and potential for/degree of human disturbance information not available to include in model. Suitable breeding habitat likely overestimated.

References

- Burleigh, T.D. 1972. Birds of Idaho. The Caxton Printers, Ltd., Caldwell, ID. 467 pp.
- Butler, R. W. 1992. Great Blue Heron (*Ardea herodias*). In: The Birds of North America No. 25. A. Poole and F. Gill, eds. The Academy of Natural Sciences, Philadelphia and the American Ornithologists Union, Washington, D. C.
- Dowd, E. M. and L. D. Flake. 1985. Foraging habitats and movements of nesting Great Blue Herons in a prairie river ecosystem, South Dakota. *J. Field Ornithol.* 56(4): 379-387.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Gibbs, J. P. 1991. Spatial relationships between nesting colonies and foraging areas of great blue herons. *Auk* 108:764-770.
- Hancock, J. and J. Kushlan. 1984. The Herons Handbook. Croom Helm, Ltd., Kent.
- Henny, J.C. AND J.E. Kurtz. 1978. Great Blue Herons respond to habitat loss. *Wildl. Soc. Bull.* 6:35-37.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Palmer, R. S. 1962. Handbook of North American birds. Vol. 1. Loons through flamingos. Yale University Press, New Haven. 567 pp.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



BLACK-CROWNED NIGHT-HERON

Nycticorax nycticorax


State rank S2S3B,SZN

Element code ABNGA11010

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Fairly widely distributed in migration; more common in e Montana (Montana Bird Distribution Committee 1996). Breed mainly in n/ncen Montana, e of the Continental Divide, and in the extreme sw; does not overwinter in the state (Montana Bird Distribution Committee 1996).

Habitat description

Broad habitat preferences, including freshwater, brackish, and marine situations (Davis 1993). Most frequently nest on islands and margins of marshes/swamps; also margins of streams, rivers, pools, ponds, lakes, lagoons, tidal mudflats, tropical mangroves, salt marshes; man-made ditches, canals, ponds, reservoirs; wet agricultural fields and dry grasslands (Hancock and Kushlan 1984, Davis 1993). Nest colonially, commonly in emergent vegetation over water, on islands, or in trees; trees preferred for roosting (Wolford and Boag 1971, Ehrlich et al. 1988, Davis 1993). Inland, colonies usually associated with large wetlands having equal proportions of open water and vegetated habitat (Davis 1993). Use of these wetlands fluctuates accordingly with rainfall conditions/water levels; occasionally may use dry grasslands and may fly up to 24 km to feeding areas (Hancock and Kushlan 1984, Davis 1993). In the Great Plains, commonly nest in bulrush or cattail dominated prairie marshes, or in groves of trees (often cottonwoods) near rivers (Johnsgard 1979, Greenwood 1981). Common throughout ND save areas with few wetlands; breeds in prairie marshes (Lokemoen 1979). In ID, prefer to nest in larger trees (alder), will also use bulrushes, willow (Burleigh 1972). Colony sites free from predators and human disturbance may persist 30-50 years (Davis 1993). Breeding distribution determined by suitable wetland habitat for feeding (aquatic and terrestrial invertebrates, fish, amphibians, reptiles, birds, rodents); breeding range has expanded to irrigated areas and water impoundments (Ehrlich et al. 1988, Davis 1993). Often nest in association with Franklin's Gulls, and may change location of the colony site if these gulls move (Wolford and Boag 1971, Davis 1993). Adequate nest cover, abundant feeding grounds, and security from predators are primary factors in nest/colony site choice (Wolford and Boag 1971, Hancock and Kushlan 1984, Davis 1993). Migration


habitat similar to that of breeding, but centers along coasts and the Mississippi Drainage (Davis 1993). Winter habitat primarily mangrove swamps in tropical areas, and marshes and swamps in FL; secondarily wetland breeding habitats (Davis 1993). Post breeding dispersal may include areas outside of the normal breeding or wintering range (Palmer 1962). Although Blue Listed by the National Audubon Society (1972-81) and listed as threatened or endangered by many states (1970s, 1980s) due to effects of pesticides and habitat loss, most populations stabilized and increasing (Ehrlich et al. 1988, Davis 1993). Trends, however, are difficult to assess because this species is nocturnal/crepuscular in nature and nests are well concealed (Ehrlich et al. 1988, Davis 1993). The Black-crowned Night-Heron is considered an "SI" species in Montana, as breeding ecology is not well known in the state (Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Uncommon migrant; does not overwinter. Migration habitat used in MT is similar to that of breeding and is assumed defined within breeding habitat parameters. Lake coverage not consistently accurate/true to mapping scale used. Assumed suitable wetland feeding habitat defined within breeding habitat parameters. Cannot model interspersed/extent of emergent marsh vegetation with open water.

References

- Burleigh, T.D. 1972. Birds of Idaho. The Caxton Printers, Ltd., Caldwell, ID. 467 pp.
- Davis, W. E., Jr. 1993. Black-crowned Night-Heron (*Nycticorax nycticorax*). In: The Birds of North America, No. 74 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union. 20 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Greedwood, R.J. 1981. Observations on Black-crowned Night Heron breeding success in a North Dakota marsh. *Can Field-Nat.* 95:465-467.
- Hancock, J. and J. Kushlan. 1984. The Herons Handbook. Croom Helm, Ltd., Kent.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Lokemoen, J.T. 1979. The status of herons, egrets, and ibises in North Dakota. *Prairie Nat.* 11(4): 97-110.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Palmer, R. S. 1962. Handbook of North American birds. Vol. 1. Loons through flamingos. Yale University Press, New Haven. 567 pp.
- Wolford, J.W. and D.A. Boag. 1971. Distribution and biology of Black-crowned Night Herons in Alberta. *Can. Field-Nat.* 85: 13-19.



WHITE-FACED IBIS

Plegadis chihi


State rank S1B,SZN

Element code ABNGE02020

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Widely distributed in migration; transient sightings generally scattered around breeding areas (Montana Bird Distribution Committee 1996). Breeding known in only a few locations including Benton Lake NWR (97 nests located in 1994), Red Rock Lakes NWR, Bowdoin NWR, and Freezout Lake WMA; does not overwinter in the state (S. Martin pers. comm., Montana Bird Distribution Committee 1996).

Habitat description

Typically use freshwater wetlands - marshes, ponds, swamps - with "islands" of emergent vegetation, e.g., cattail, bulrush (Johnsgard 1979, Ehrlich et al. 1988, Ryder and Manry 1994). Nest usually in emergent vegetation or shrubs, bushes, low trees in/or surrounded by shallow water; also on deserted muskrat houses; occasionally dredge spoil banks (Ryder 1967, Johnsgard 1979, Schmidt 1980, Ehrlich et al. 1988, Ryder and Manry 1994). Colonies often associated with Black-crowned Night-Herons, Cattle Egrets, Snowy Egrets, Great Blue Herons, Franklin's Gulls (Johnsgard 1979, Ehrlich et al. 1988, Ryder and Manry 1994); in the Rocky Mountain region and Great Basin area, a common nesting associate is the Cinnamon Teal (Ryder 1967). In ND, UT, and s AK, nest commonly within dense stands of bulrushes and/or cattails in large, shallow wetland complexes, near edges of open water (Ryder 1967, Schmidt 1980, Goosen et al. 1995). Are fairly versatile nesters; nest in emergent rushes or willows - primary breeding requirement is adequate isolation of both nesting colony and night roosting sites (Trost in Clark et la. 1989). Colonies established in CO above 7500ft elevation, and as high as 4300m in Bolivia (Ryder 1967, Ryder and Manry 1994). Food taken is primarily aquatic and moist soil invertebrates, insects, fish, small vertebrates; in breeding and migration season, frequently forage in larger, shallowly flooded wetlands of short emergent vegetation, lake and pond edges, irrigated/flooded alfalfa and small grain fields, and hay meadows (Ryder 1967, Ryder and Manry 1994). Drainage of wetlands, development, and intensive overgrazing destroy nesting habitat; many colonies have disappeared - breeding has not occurred in the once important Central Valley, CA since 1954 (Weller et al. 1958, Ryder 1967, Goosen et al. 1995). Entire colonies may move if breeding habitat conditions deteriorate (Trost in Clark et la. 1989); water diversion from wetlands with traditional

colony sites has caused temporary to permanent abandonment - wetland areas subject to periodic severe drought do not serve as consistent colony sites, since emergent vegetation may take up to two years to recover (Ryder and Manry 1994). Presence of Black-crowned Night-Herons may indicate favorable conditions; managed wetlands provide footholds to populations attempting to pioneer new areas (Goosen et al. 1995). Wetlands with relatively consistent water levels are critical to breeding habitat (Ryder and Manry 1994, Goosen et al. 1995). In ND, is casual migrant and summer resident mainly in the se; also breed in Alberta (Lokemoen 1979). Breeding populations have increased in the Great Basin area and expanded into the Great Plains due to pesticide banning, wildlife refuge management, and increased alfalfa planting (Ryder and Manry 1994). Regardless of population recoveries or range expansions, are sensitive to human disturbance, water level fluctuation, and are still vulnerabel to pesticides and insecticides on both breeding and wintering grounds (Trost in Clark et la. 1989). The White-faced Ibis is considered a 'threatened, endangered, or sensitive' species in MT - breeding is uncommon and MT is at the northern limit of range (Montana Bird Distribution Committee 1996, MNHP-VCA 1996) - and a 'species of management concern' and 'category 2 candidate' for listing as a threatened and endangered species by the USF&WS (for the Great Basin and Region 6) because few consistent breeding sites exist and population status is poorly known (Ryder and Manry 1994). Overall, populations are declining throughout North America (Ryder and Manry 1994).

Model assumptions & caveats

Breeding habitat modeled. Uncommon but fairly regular migrant; does not overwinter. Potential migration habitat used in MT assumed defined within breeding habitat parameters. Hydrography coverage not consistently accurate/true to mapping scale used. Although breeding lakes need not be permanent, water levels should be consistent; persistence of intermittent lakes (seasonal vs. semipermanent, etc.) is unknown therefore not included in model. Cannot model interspersions of cover types (islands of emergent vegetation within open water areas) at mapping scale used. Breeding associate relationships not modeled. Foraging habitats included in breeding habitat parameters, but may diminish relavance of model regarding nesting habitat requirements. Assume complex of wetlands inherent with buffer used in model.

References

Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.

Goosen, J.P., D.E. Ealey, H. Judge and D.C. Duncan. 1995. Distribution and breeding status of the White-faced Ibis, *Plegadis chihi*, in Canada. *Can. Field-Nat.* 109(4):391-402.

Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.

Lokemoen, J.T. 1979. The status of herons, egrets, and ibises in North Dakota. *Prairie Nat.* 11(4): 97-110.


Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Ryder, R. A. 1967. Distribution, migration and mortality of the White-faced Ibis (*Plegadis chihi*) in North America. *Bird-Banding* 38: 257-277.

Ryder, R.R. and D.E. Manry. 1994. White-faced Ibis (*Plegadis chihi*). In: The Birds of North America No. 130. A. Poole and F. Gill, eds. The Academy of Natural Sciences, Philadelphia and The American Ornithologists Union, Washington, D.C.

Schmidt, R.A. 1980. First breeding record of the White-faced Ibis in North Dakota. *Prairie Nat.* 12(1): 21-23.

Weller, M.W., B.H. Wingfield and J.B. Low. 1958. Effects of habitat deterioration on bird populations of a small Utah marsh. *Condor* 60: 220-226.



TRUMPETER SWAN

Cygnus buccinator


State rank S2B,S2N

Element code ABNJB02030

Global rank G4

Modeled by W. Williams

MTNHP status special concern



Predicted habitat: 35,011 ha, 0.09 % of state.

State range

Isolated populations currently breed along the Continental Divide (Bean Lake) and in the Red Rock Lakes/Greater Yellowstone area, but old records exist for latilongs 2, 11, 15, 26, 28, & 40 (Hand 1969, Montana Bird Distribution Committee 1996,). Migration is mainly through western Montana and the East Front of the Rockies; overwintering occurs primarily within the Greater Yellowstone Ecosystem and in isolated areas in se and w Montana (T. McEneaney pers. comm., Montana Bird Distribution Committee 1996).

Habitat description

Breed on shallow freshwater marshes, ponds, lakes, and slow moving rivers with stable water levels and adequate area for take of, ~100m (Johnsgard 1979, Hampton 1981, Ehrlich et al. 1988, Mitchell 1994). Preferred nesting areas have abundant and diverse submerged aquatic vegetation for food, dense stands of emergent vegetation for cover, highly irregular shorelines, water depths <1.2m, nest site structures such as muskrat houses or sedge islands, and are protected from human disturbance (Johnsgard, 1979, Ehrlich et al. 1988, Mitchell 1994). Subsist on aquatic vegetation; cygnets feed primarily on insects, other invertebrates, and aquatic vegetation; secluded breeding habitats with abundant invertebrate populations and aquatic plants promote high productivity (Page 1976, Shea 1979, Hampton 1981, Gale et al. 1987, Mitchell 1994). On the Canadian prairie, prefer aspen parkland/pothole habitat, but breeding extends into boreal forest regions (Palmer 1962, Gale et al. 1987). Although the entire state of MT was once part of former breeding range, the Yellowstone/Centennial Valley region of nw WY and sw MT is now the most important breeding area in the contiguous U.S. (Palmer 1962, Bellrose 1978, Gale et al. 1987). Two remnant breeding populations are rare but permanent residents in the area: The Rocky Mountain population, which breeds along the Rocky Mountain corridor; the Tristate population, which breeds in the Red Rock Lakes/Centennial Valley area and includes scattered pairs breeding throughout the e ID/nw WY/YNP area (Johnsgard 1986, Gale et al. 1987). Nesting territories in these areas occurs between 1770-2515m elevation (Shea

1979), and include interconnected, shallow lakes and marshes, extensive wetlands, oxbow/pothole habitat adjacent to rivers, isolated ponds (0.4-14ha) surrounded by rangeland, braided river complexes in forest and sagebrush habitat, and areas of widely scattered ponds and lakes, 1.2-110.8ha in size (Shea 1979, Gale et al. 1987). Nests found very commonly on muskrat houses; also on hummocks, floating bogs, islands, abandoned beaver lodges (Page 1976, Shea 1979, Gale et al. 1987). At higher elevations, select lakes more isolated from human activity; at lower elevations, prefer water areas with stands of emergent vegetation for cover and as a buffer from human disturbance (Gale et al. 1987). Critical to breeding habitat: quiet waters with a highly irregular shoreline, shallow areas for foraging, stable permanent water levels, and isolation from human perturbations (Page 1976, Shea 1979, Hampton 1981, Gale et al. 1987, Mitchell 1994). In migration also use larger rivers and brackish estuaries, larger lakes and reservoirs as staging/foraging areas; in winter also use larger, open lakes, sheltered bays, estuaries (Palmer 1962, Mitchell 1994, MNHP-VCA 1996). In MT, winter habitat consists of larger areas of open, ice free water with adequate aquatic vegetation, sand/gravel areas for loafing, little shrub cover, and little human disturbance (Gale et al. 1987, Mitchell 1994). The geothermal activity of the YNP region keeps water open in subzero temperatures, however, aquatic vegetation that is available in winter is an inadequate food source (Gale et al. 1987). Supplemental grain sustains the resident population throughout the winter, but discourages normal migration patterns - seasonal movements of this population are limited to local flights between breeding and wintering areas (Gale et al. 1987, Palmer 1962). Future Trumpeter Swan management includes cessation of this feeding program, however, making both migration and winter habitat even more critical (A.Perkins, pers. comm.). Breeding range once extended across most of North America, but commercial and sport hunting, and habitat destruction in the 1920s-1930s pushed the Trumpeter Swan to near extinction, and eliminated it from most of its historic range (Ehrlich et al. 1988, Gale in Clark et al. 1989). Yellowstone and Centennial Valley populations have shown past declines; the Trumpeter Swan is considered a threatened, endangered, or sensitive species in MT and is a sensitive species of concern in WY (Gale in Clark et al. 1989, Montana Bird Distribution Committee 1996)

Model assumptions & caveats

Breeding habitat modeled. Uncommon but regular migrant; local year round resident movements limited, hence most suitable winter/migration habitat assumed defined within breeding habitat parameters. Hydrography coverage not consistently accurate/true to mapping scale used. Water body size restrictions local, only minimum size limit included in model. Elevation limits specified in literature lowered to include known breeding lake (Bean Lake), therefore suitable predicted habitat will increase. River flow ('slow') and braiding information/coverage not available, therefore river habitat likely overestimated. Cannot model water body morphology (irregular shoreline) at mapping scale used. Water depth information not available. Principal food source, submerged aquatic vegetation not mapped. Cannot model nest site structures (muskrat houses, sedge islands) at mapping scale used. Potential for human disturbance information not available; cannot model potential human access at mapping scale used. Suitable breeding habitat is overestimated.

References

- Bellrose, F.C. 1978. Ducks, geese, and swans of North America. Wildlife Management Institute, Stockpole Books.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Gale, R.S., E.O. Garten, and I.J. Ball. 1987. The history, ecology, and management of the Rocky Mountain population of Trumpeter Swans. Unpubl. Manuscript. Montana Wildlife Cooperative Unit, University of Montana. 314 pp.
- Hampton, P. D. 1981. The wintering and nesting behavior of the Trumpeter swan. M.S. thesis. Univ. of Montana, Missoula. 185 pp.
- Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.

Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.

Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.


Mitchell, C.D. 1994. Trumpeter Swan (*Cygnus buccinator*). In: The Birds of North America, No. 105. A. Poole and F. Gill, eds. Academy of Natural Sciences, Philadelphia and The American Ornithologists Union, Washington, D. C.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Page, R.D. 1976. The ecology of Trumpeter Swans on Red Rock Lakes National Wildlife Refuge, Montana. Dissertation, University of Montana, Missoula.

Palmer, R. S. 1962. Handbook of North American birds. Vol. 1. Loons through flamingos. Yale University Press, New Haven. 567 pp.

Shea, R. E. 1979. The ecology of Trumpeter Swans in Yellowstone National Park and vicinity. M.S. Thesis. University of Montana, Missoula.



HARLEQUIN DUCK

Histrionicus histrionicus


State rank S2B,SZN

Element code ABNJB15010

Global rank G4

Modeled by W. Williams

MTNHP status special concern



State range

Breed very locally on mountain streams in western Montana (Reichel and Genter 1995, Montana Bird Distribution Committee 1996). Occupied streams primarily tributaries of the Kootenai River, Flathead River, lower Clark Fork River, and upper Blackfoot River; and streams along the Rocky Mountain Front and in the mountains along the northern border of Yellowstone National Park (Reichel and Genter 1995). Transient sightings are rare; overwintering verified in latilong 26 only (Montana Bird Distribution Committee 1996).

Habitat description

Primarily a boreal and subarctic breeder; western breeding population center is in AK and B.C. (Bellrose 1978, Palmer 1962). Use riparian habitats for feeding, nesting, cover; breed near swiftly flowing, clear, forested or well vegetated, undisturbed mountain streams; occasionally on open tundra (Cassirer et al. 1996, Ehrlich et al. 1988, Johnsgard 1975). Adults and broods sited in areas 600m-2000+m elevation (Genter 1992, Cassirer et al. 1991). In the Rocky Mountain area, breeding streams fall into two subprovinces: the n Columbia Basin (nw MT incl. GNP, Rocky Mountain Front, n ID), and the Intermountain Province (ID, sw MT, WY incl. GYE) (Cassirer et al. 1996). Rocky Mountain breeding habitat requirements are fairly rigid: stream size \geq second order; 1%-7% stream gradient with shallow riffle areas; clear water; gravel to boulder sized rocky substrate; forested stream bank vegetation; hiding cover along streams (dense shrub vegetation, undercut banks, logjams, etc.); absence of human access (roads, trails) and disturbance (Cassirer et al. 1996). In WY, occupy stream sections with bank shrub vegetation, gradients \leq 2 degrees, >3 loafing sites per 10m, and braided channels (Wallen 1992). In ID use streams \geq 3m wide with flow velocities > 1 m/sec, gravel substrates, and undisturbed forest along stream banks (Cassirer and Groves 1992). In n ID, occupy streams 600-1200m in elevation with associated riparian habitat dominated by Western Red Cedar or Western Hemlock; in se ID use streams 1770-1890m in elevation with riparian habitat dominated by shrubs and overstory of Douglas-fir (Cassirer et al. 1991). On the Rocky Mountain Front, favor streams 6-10m wide with cobble

substrate, in v-shaped valleys of Lodgepole Pine and Engelmann Spruce stands (Diamond and Finnegan 1992). In nw WA, however, timber stand characteristics are apparently not a factor in habitat use - stream morphology the primary factor in breeding habitat selection (Schirato and Sharpe 1992). Few nest sites have been described for the Pacific population (Cassirer et al. 1993). Nests found within 5m of streams in a variety of locations: on rocks/rock crevices; on the ground in dense vegetation cover; in mossy cliff crevices; in piles of woody debris; in undercut stream banks; in hollow tree cavities near tree base and as high as 1.8m (Cassirer et al. 1996, Cassirer et al. 1993). Brood rearing areas initially in ponds and slow meanders of streams, shifting to slower current areas, then swift water as broods mature (Kuchel 1977). Subsist primarily on aquatic insects and larvae, crustaceans, mollusks (Ehrlich et al. 1988, Bellrose 1978). Migration routes to inland breeding areas are likely along river corridors and ascending tributaries, with intermittent overland flights (Cassirer et al. 1996, Kuchel 1977). Winter almost exclusively on turbulent, rocky coastal waters, 1-2 fathoms deep (A.O.U. 1983, Palmer 1962). Productivity appears limited by human activities/developments in breeding areas, consequently, breeding is habitat restricted to remote, pristine areas of swiftly flowing streams with forested stream banks and calmer oxbow areas (Cassirer and Groves 1992, Kuchel 1977). Breeding populations in Grand Teton National Park and Jasper National Park are declining due to increases in human disturbance in/near breeding areas (Clarkston 1992, Wallen 1992). Stream food supply, high spring runoff, and winter/migration circumstances also limit population size and productivity (Cassirer et al. 1996, Cassirer and Groves 1992). The Harlequin Duck is classified as a C2 candidate for threatened and endangered status in the U.S., and is considered a USFS sensitive species in Regions 1 and 4 (Cassirer et al. 1996, Cassirer et al. 1991).


Model assumptions & caveats

Breeding habitat modeled. Rare migrant, very rare winter resident; potential nonbreeding habitat available in Montana assumed defined within breeding habitat parameters. Hydrography may not be consistently accurate and/or true to mapping scale used. Stream morphology attributes - flow rate, clarity, order number, gradient, substrate characteristics - not available to include in model. Stream bank morphology and potential for human disturbance/access information not available to include in model. Suitable breeding habitat is overestimated.

References

- Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.
- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Bellrose, F.C. 1978. Ducks, geese, and swans of North America. Wildlife Management Institute, Stockpole Books.
- Cassirer, E.F. and C.R. Groves. 1992. Status and breeding of Harlequin Ducks in Idaho. in Proceedings, Harlequin Duck Symposium, April 23-24, 1992. Unpubl. Rep., 45pp. Idaho Dept. Fish and Game, Moscow, ID.
- Cassirer, E.F., C.R. Groves and R.L. Wallen. 1991. Distribution and population status of Harlequin Ducks in Idaho. Wilson Bull. 103(4): 723-725.
- Cassirer, E. F., G. Schirato, F. Sharpe, C. R. Groves, and R. N. Anderson. 1993. Cavity nesting by harlequin ducks in the Pacific Northwest. Wilson Bull. 105:691-694.
- Cassirer, E. F., J. D. Reichel, R. L. Wallen, and E. Atkinson. 1996. Harlequin Duck (*Histrionicus histrionicus*) conservation assessment and strategy for the U.S. Rocky Mountains. Montana Natural Heritage Rep., 53 pp. plus appendices.
- Clarkston, P. 1992. A preliminary investigation into the status and distribution of Harlequin Ducks in Jasper National Park. in Proceedings, Harlequin Duck Symposium, April 23-24, 1992. Unpubl. Rep., 45pp. Idaho Dept. Fish and Game, Moscow, ID.

- Davis, C.V. 1961. A distributional study of the birds of Montana. Ph.D. dissertation. Oregon State University, Corvallis. 462 pp.
- Diamond, S. and P. Finnegan. 1992. Harlequin Duck ecology on Montana's Rocky Mountain Front. in Proceedings, Harlequin Duck Symposium, April 23-24, 1992. Unpubl. Rep., 45pp. Idaho Dept. Fish and Game, Moscow, ID.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Genter, D.L. 1992. Status of the Harlequin Duck (*Histrionicus histrionicus*) in Montana. In Proceedings, Harlequin Duck Symposium, April 23-24, 1992. Unpubl. Rep., 45pp. Idaho Dept. Fish and Game, Moscow, ID.
- Johnsgard, P.A. 1975. Waterfowl of North America. Indiana University Press, Bloomington.
- Kuchel, C. R. 1977. Some aspects of the behavior and ecology of harlequin ducks breeding in Glacier National Park, Montana. M.S. thesis. Univ. of Montana, Missoula. 160 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Palmer, R. S. 1962. Handbook of North American birds. Vol. 1. Loons through flamingos. Yale University Press, New Haven. 567 pp.
- Reichel, J. and D.L. Genter. 1995. Harlequin Duck surveys in western Montana: 1994. Montana Nat. Heritage Prog. 58 pp.
- Schirato, G. and F. Sharpe. 1992. Distribution and habitat of Harlequin Ducks in northwestern Washington. in Proceedings, Harlequin Duck Symposium, April 23-24, 1992. Unpubl. Rep., 45 pp. Idaho Dept. Fish and Game, Moscow, ID.
- Wallen, R. 1992. Annual variation in Harlequin Duck population size, productivity and fidelity in Grand Teton National Park, Wyoming. in Proceedings, Harlequin Duck Symposium, April 23-24, 1992. Unpubl. Rep., 45pp. Idaho Dept. Fish and Game, Moscow, ID.




BALD EAGLE

Haliaeetus leucocephalus

State rank S3B,S3N
Global rank G4
MTNHP status special concern

Element code ABNKC10010
Modeled by C. Tobalske, W. Williams



State range

Breeds and winters widely throughout Montana (Montana Bird Distribution Committee 1996). Nests mostly in the western and central part of the state, especially along the Clark Fork River and tributaries; also on the Missouri River below Ft. Benton and the Greater Yellowstone area — range expansion is accelerating in eastern Montana (Bureau of Land Management 1986, T. McEneaney pers. comm.).

Habitat description

From Montana Bald Eagle Working Group (1991): In general, Bald Eagles prefer solitude, late-successional forests, and shoreline adjacent to open water; proximity of a large prey base and the presence of large, mature trees for nesting are important habitat components. Medium and high habitat quality are characterized by cottonwood, Douglas-fir, Ponderosa Pine or mixed conifer stands of 20 acres minimum, with low to moderate canopy closure, a slope less than 40%, a water body within 1 mile, and no southerly aspect. There should be more than two suitable nest trees, and more than three perch trees, in the stand. Because young are particularly vulnerable to food deprivation, an adequate prey base within the home range is essential (Stalmaster et al. 1985 in Montana Bald Eagle Working Group 1991). Feeding habitat should be greater than 80 acres with shallows, and present some grasslands and meadows. Distance to human activities should be greater than 0.25 miles, and less than 10% of the shoreline should be developed.


In the Greater Yellowstone Ecosystem, breed in riparian or lacustrine habitats, selecting older, larger trees for nesting that are usually within 1.5 km of a river or lake shore; occasionally forage in upland areas (Harmata in Clark et al. 1989). Outside of the breeding season population densities and distributions vary depending on adequate food supply and roosting sites (Johnsgard 1990). Roosting sites are well protected from wind and are usually in the tallest, dominant trees in a stand — in Glacier National Park preferred roost sites were in old growth stands of Western Larch (Crenshaw 1985). Bald Eagles are sensitive to human disturbance and may abandon nest sites or territories depending on its intensity (Montana Bald Eagle Working Group 1991).

Model assumptions & caveats

Hydrography coverage may not be consistently accurate/true to mapping scale used. Focus of model on suitable breeding habitat, however most nonbreeding habitat also defined within breeding habitat parameters. Slope not included in model — if included would diminish availability of wind protected roosting sites within cover types selected; assume suitable, level (<40% slope) nest sites present within hydrography buffer around major rivers/lakes. Human disturbance, prey base and water turbidity information not available to include in model — breeding habitat therefore likely overestimated, as high disturbance and marginal foraging habitat are defined within breeding habitat parameters.

References

- Anthony, R. G., and F. B. Isaacs. 1989. Characteristics of bald eagle nest sites in Oregon. *J. Wildl. Manage.* 53:148-159.
- Bird, D. M., editor. 1983. *Biology and management of bald eagles and ospreys.* MacDonald. 325 pp.
- Bureau of Land Management. 1986. *Montana Bald Eagle Management Plan.* U.S.D.I., Billings, MT.
- Caton, E. L., et al. 1992. Characteristics of foraging perches used by breeding bald eagles in Montana. *Wilson Bull.* 104:136-142.
- Clark, T.W., A.H. Harvey, R.D. Dom, D.L. Genter, and C. Groves, eds. 1989. *Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem.* Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- Crenshaw, J. G. 1985. *Characteristics of bald eagle communal roosts in Glacier National Park, Montana.* M.S. thesis. University of Montana, Missoula. 92 pp.
- Jenkins, A.M. 1979. *Bald eagle essential habitat on and near BLM lands in Wyoming.* Animal Report 79. 34 pp.
- Johnsgard, P.A. 1990. *Hawks, eagles, and falcons of North America.* Smithsonian Institution Press, Washington, D.C. 403 pp.
- Kralovec, M. L., et al. 1992. Nesting productivity, food habits, and nest sites of bald eagles in Colorado and southeastern Wyoming. *Southwest. Nat.* 37:356-361.
- Lincer, J. L., W. S. Clark and M. N. LeFranc, Jr. 1979. *Working bibliography of the bald eagle.* Raptor Information Center, Nat. Wildlife Fed., Washington, D.C. NWF Scientific/technical Series No. 2. 219 pp.
- Mace, R.D., D. Casey and K. Dubois. 1987. *Effects of water level fluctuations on productivity and distribution of Ospreys and Bald Eagles in the northern Flathead Valley.* Montana Fish, Wildl. and Parks, for Montana Power Company, Butte. 90 pp.
- Montana Bald Eagle Working Group. 1991. *habitat management guide for bald eagles in northwestern Montana.* 29 pp.
- Montana Bird Distribution Committee. 1996. *P. D. Skaar's Montana Bird Distribution, Fifth Edition.* Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Swenson, J. E. 1975. *Ecology of the bald eagle and osprey in Yellowstone National Park.* MS thesis. Montana State Univ., Bozeman.
- U.S. Department of the Interior, Bureau of Land Management, Platte River Resource Area. 1992. *Final Bald Eagle Habitat Management Plan.* U.S. Government Printing Office.



NORTHERN GOSHAWK

Accipiter gentilis


State rank S3S4

Element code ABNKC12060

Global rank G5

Modeled by C. Tobalske, W. Williams

MTNHP status special concern



State range

Breeds in mountainous or coniferous regions of the state (primarily in the west) and occasionally winters in the lower valleys of western Montana (Montana Bird Distribution Committee 1996, T. McEneaney pers. comm.).

Habitat description


A raptor usually found in mature, dense stands of mixed or pure forests, 50-100 m from a clearing, often with a permanent or temporary stream or lake nearby (Jones 1979). On the Kaibab Plateau, preferred nesting habitat consists of closed stands of Ponderosa Pine and mixed conifers (canopy closure > 80%), with most nests located on northern aspects; proximity to water important only in more open stands (Crocker-Bedford and Chaney 1988). In Wyoming, often nests in old-growth and mature forests, such as closed Lodgepole Pine and lodgepole-aspen forests composed of large, mature trees (Squires and Ruggiero 1996). In Oregon, nests are located in Douglas-fir, White, Ponderosa and Lodgepole Pines, and mixed conifer stands; prefer mature or old-growth stands on gentle slopes and northern aspects within 600m of water (Reynolds et al. 1982, Moore and Henny 1983). In the Inyo National Forest (California), criteria for primary habitat include: Red Fir, Lodgepole Pine, aspen, or mixed conifer stands; dbh > 61 cm, canopy closure > 40%, slope 0-20%, stand size 20-50 ha, and distance to water < 0.4 km (McCarthy et al. 1987). In Montana, typical nesting habitat consists of mature/overmature coniferous forests with a closed canopy (75-85%) on moderate (15-35%), north-facing slopes, with water or a large clearing within 500 m of the nest — in drier areas, nests can be found in even-aged stands if the canopy is closed (Hayward and Escano 1989, Whitford 1991, Hendricks and Dueholm 1995). A wide variety of habitats are used during the winter in Montana, particularly grass-shrub-steppe habitats which host upland gamebird prey (D. Casey pers. comm.). Data on abundance of breeding Northern Goshawks in the U.S. indicate highly variable densities, both spatially and temporally, but reportedly numbers are stable with no evidence to suggest range contraction (Kennedy 1997).

Model assumptions & caveats

Distance to water not included in model; hydro type (river, lake, etc.) not specified and water not clearly defined as breeding habitat requirement. Interspersion of forest and clearings could not be resolved and included at mapping scale used. High canopy closure was selected in attempt to also select for more mature stands — stand age information not available at mapping scale used.

References

- Anonymous. 1993. Northern goshawk management workshop. Proceedings of March 1993 workshop, Madison, Wisconsin. 60 pp.
- Crocker-Bedford, D. C. 1990. Goshawk reproduction and forest management. *Wildl. Soc. Bull.*, 18:262-269.
- Crocker-Bedford, D. C., and B. Chaney. 1988. Characteristics of goshawk nesting stands. Pages 210-217 in Glinski et al., eds. Proc. Southwest raptor management symposium and workshop. *Natl. Wildl. Fed. Sci. and Tech. Ser. No. 11.*
- Hayward, G. D. 1989. Goshawk Nest-Site Characteristics in Western Montana and Northern Idaho. *Condor* 91:476-479.
- Hayward, G. D., and R. E. Escano. 1989. Goshawk nest-site characteristics in western Montana and northern Idaho. *Condor* 91:476-479.
- Hendricks, P., and K. H. Dueholm. 1995. Cliff-nesting raptor survey of the Sioux District, Custer National Forest: 1994. Unpubl. report to U.S.D.A. For. Serv., Custer N.F., Billings. 20 pp.
- Jones, S. 1979. The accipiters - goshawk, Cooper's hawk, sharp-shinned hawk. US Bureau of Land Management Tech. Rep. No. 335. Denver. 51 pp.
- Kennedy, P.L. 1997. The Northern Goshawk (*Accipiter gentilis atricapillus*): Is there evidence of a population decline? *J. Raptor Res.* 31(2): 95-106.
- McCarthy, C., W.D. Carrier, and W.F. Laudenslayer. 1987. Coordinating timber management activities with raptor nesting habitat requirements. Pages 229-235 in Proceedings of the Western Raptor Management Symposium and Workshop, Boise. National Wildlife Federation Scientific and Technical Series 12.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Moore, K. R., and C. J. Henny. 1983. Nest site characteristics of three coexisting accipiter hawks in northeastern Oregon. *J. Raptor Res.* 17:65-76.
- Reynolds, R. T., E. C. Meslow, and H. M. Wight. 1982. Nesting habitat of coexisting *Accipiter* in Oregon. *J. Wildl. Manage.* 46:124-138.
- Shuster, W. C. 1980. Northern goshawk nest site requirements in the Colorado Rockies. *West. Birds* 11:89-96.
- Squires, J. R., and L. F. Ruggiero. 1996. Nest-site preference of northern goshawks in southcentral Wyoming. *J. Wildl. Manage.* 60(1):170-177.
- Whitford, T. C. 1991. Defining old-growth Douglas-fir forests of central Montana and use of the northern goshawk (*Accipiter gentilis*) as a management indicator species. MS Thesis. Univ. of Montana, Missoula. 62 pp.



FERRUGINOUS HAWK

Buteo regalis


State rank S3B,SZN

Element code ABNKC19120

Global rank G4

Modeled by C. Tobalske, W. Williams

MTNHP status special concern



Predicted habitat: 15,365,144 ha, 40.35 % of state.

State range

Breed east (and very rarely west) of the Continental Divide (Montana Bird Distribution Committee 1996); highest numbers in southwestern (Atkinson 1992) and southeastern Montana (Ensign 1983). Sighted throughout the state during migration (Montana Bird Distribution Committee 1996).

Habitat description

A species found in undisturbed, western plains or shrub-steppe regions; in relatively unbroken terrain with scattered trees, rock outcrops, tall trees along creek bottoms (Evans 1982, Restani *in* Clark et al. 1989). Breed in grassland and sagebrush habitats, saltbush-greasewood shrublands, and forest edges, particularly western pinyon-juniper — avoids high elevations, heavily wooded areas/interiors, narrow canyons, and extensive agricultural lands (Snow 1974, Jasikoff 1982, Olendorff 1993, Bechard and Schmutz 1995). Prefer to nest in isolated trees and small groves of trees in grassland/shrubsteppe areas, sparse riparian forests, and broken bluffs, buttes, cliffs and rock outcrops in canyon/badland habitats (several studies *in* Bechard and Schmutz 1995). Several studies have described nest sites in Montana: in the southeast (Carter County), most nests were found on the ground (Wittenhagen 1992); in the Centennial Valley (south central), most nests were in willows along streams (Restani *in* Clark et al. 1989); in southwestern Montana, Atkinson (1992) located 50 nests upon rocky outcrops, on southern aspects; habitat within 100 m of the nests consisted of grasslands and shrublands (sagebrush steppe, foothill prairie, mountain mahogany associations); within 1600 m grassland was the dominant cover type. Other nest sites include trees, buttes, hay stacks, power poles, and other manmade structures (Hendricks and Dueholm 1995); elevated nest sites are strongly favored (Bechard and Schmutz 1995). Primary prey taken are rabbits, ground squirrels, prairie dogs; in many areas nesting pair densities are correlated with ground squirrel or jackrabbit abundance (Bechard and Schmutz 1995, Hendricks and Dueholm 1995). Winter in open terrain/grasslands, where prey is abundant — commonly around prairie dog towns; also near cultivated fields with pocket gophers (Olendorff 1993, Bechard and Schmutz 1995). Widescale conversion of native prairie into agricultural lands greatly diminished breeding habitat in North Dakota and Manitoba (Stewart 1975, Salt and Salt 1976). Currently,

numbers have increased in North Dakota, South Dakota, Montana, Alberta and Manitoba; populations reportedly stable in these areas (Olendorff 1993, Dobkin 1994), however, human disturbance, overgrazing (esp. in riparian areas), and continued agricultural activities threaten local breeding populations (Restani in Clark et al. 1989).

Model assumptions & caveats

Use of elevation in model not appropriate due to profound differences in habitats with latitude and elevation across the state — slope ($\leq 20\%$) was used in attempt to filter out higher, more mountainous terrain. Versatile nester — assume suitable nest site defined within breeding habitat parameters. Interspersion of forested with grassland cover types could not be resolved at mapping scale used; suitable habitat for this fairly restricted breeder is likely overestimated.

References

- Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.
- Atkinson, E. C. 1992. Ferruginous hawk (*Buteo regalis*) inventories on the Dillon Resource Area of southwest Montana: 1992. Montana Natural Heritage Program for Bureau of Land Management, Dillon Resource Area. 34 pp.
- Atkinson, E. C. 1995. Survey and monitoring guidelines for Ferruginous Hawks in Montana. Bureau of Land Management and Montana Dept. Fish, Wildlife and Parks. 42 pp.
- Bechard, M. J., and J. K. Schmutz. 1995. Ferruginous Hawk (*Buteo regalis*). In: *The Birds of North America*, No. 172 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, and The American Ornithologists' Union, Washington, D.C. 20 pp.
- Black, A. 1992. Ferruginous hawk reproduction and habitat survey. Unpublished report. N. Rockies Cons. Coop., Jackson, WY. 13 pp.
- Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ensign, J. T. 1983. Nest site selection, productivity, and food habits of ferruginous hawks in southeastern Montana. M.S. thesis. Montana State University. 85 pp.
- Evans, D. L. 1982. Status reports on twelve raptors. U.S. Department of the Interior, Fish and Wildlife Service, Special Scientific Report No. 238. 68 pp.
- Hendricks, P., and K. H. Dueholm. 1995. Cliff-nesting raptor survey of the Sioux District, Custer National Forest: 1994. Unpubl. report to U.S.D.A. For. Serv., Custer N.F., Billings. 20 pp.
- Jasikoff, T.M. 1982. Habitat suitability index models: ferruginous hawk. U.S.D.I. Fish Wildl. Serv. FSW/OBS-82/10.10. 18 pp.
- Kantrud, H. A. 1981. Grazing intensity effects on the breeding avifauna of North Dakota native grasslands. *Can. Field-Nat.* 95:404-417.
- Kantrud, H.A. and R.L. Kologiski. 1982. Effects of soils and grazing on breeding birds of uncultivated upland grasslands of the northern Great Plains. U.S.D.I., Fish and Wildl. Serv., Wildl. Res. Rep. 15. 33 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Olendorff, R.R. 1993. Status, biology, and management of Ferruginous Hawks: a review. Raptor Res. and Tech. Asst. Cen., Spec. Rep. U.S. Dept. Interior, Bur. Land Manage., Boise, ID. 84 pp.

Salt, W.R. and J.R. Salt. 1976. The birds of Alberta. Hurtig Publishers, Edmonton, Alberta. xv + 498 pp.

Skaar, P.D. 1969. Birds of the Bozeman latilong. P.D. Skaar, Publ. Bozeman. 132 pp.

Snow, C. 1974. Habitat management series for unique or endangered species: ferruginous hawk, *Buteo regalis*. Report 13, Bureau of Land Management, U.S. Dept. of Int., Technical Note, T/N-255, Denver. 23 pp.

Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.


Wittenhagen, K.W. 1992. Final report on the Ferruginous Hawk in southeastern Montana. Unpub. Report for Bur. Land Manage. Powder River and Big Dry Resource Area, Miles City, Montana. 31 pp.

PEREGRINE FALCON

Falco peregrinus

State rank S1S2B,SZN
Global rank G4
MTNHP status special concern

Element code ABNKD06070
Modeled by C. Tobalske, W. Williams



State range

Breeding scattered widely but rare and localized in Montana; wintering also widespread throughout the state (Bergeron et al. 1992, Montana Bird Distribution Committee 1996).

Habitat description

A bird typically associated with water and tall cliffs; found in diverse, open country on every continent except Antarctica (Evans 1982, Johnsgard 1990). Peregrines nest almost exclusively in high cliffs, favoring those which are near to or overlooking water (Snow 1972, Hendricks and Dueholm 1995). In Utah, all nests found were within 1 mile from water and in cliffs at least 45 m tall; densely forested areas apparently avoided (Herman and Willard 1978). In the Greater Yellowstone Ecosystem, river gorges and mountain cliffs are typical nest sites — cliffs are at least 60 m tall (Langelier *in* Clark et al. 1986). Eyries are usually located below 9500', within 1 mile from water and within 10 miles of hunting habitat: wetland and riparian areas, meadows and parklands, croplands, gorges, mountain valleys, and lakes (U.S. Fish and Wildlife Service 1984). In the Rocky Mountains, most eyries are located near Ponderosa Pine forests or pinyon-juniper woodlands (U.S. Fish and Wildlife Service 1984). Preferred nest locations are in undisturbed areas with a wide view, near water and abundant prey; acceptability of a nesting cliff, however, is a complex combination of factors, including availability and demand for nest sites, geological characteristics of the cliff, changing environmental conditions, increasing human disturbance, changes in prey density, individual behavioral variations, and longer term population shifts (Snow 1972). The diet consists primarily of small birds; hunting has been documented as far as 18 mi from nest sites, although a 10-mile hunting radius is more common (Langelier *in* Clark et al. 1986). Chlorinated hydrocarbon use contributed to dramatic past population declines; by 1970 breeding Peregrine Falcons were extirpated from the northern Rocky Mountain states (Langelier *in* Clark et al. 1986). Reintroduction efforts have been successful; in Montana numbers of breeding pairs increased from 1 in 1984 to 23 in 1997 — from inception of reintroduction efforts in Montana, 546 birds have been released into the wild from hack sites (T.

McEneaney pers. comm.). Subspecies breeding in the state is *F. p. anatum* — extent of migratory behavior is apparently within the interior U.S.; in the more northerly latitudes *anatum* will move southward if winter prey supply is inadequate (Snow 1972, MNHP-VCA 1996). Migratory behavior is largely restricted to *F. p. tundrius* populations, likely the widespread migrant in Montana (Snow 1972, MNHP-VCA 1996).

Model assumptions & caveats

Cliff cover type could not be resolved at mapping scale used — attempted to model potential cliff habitat by intersecting cover type classified as rock with slope greater than or equal to 40%. Potential cliff areas were then buffered by 10 miles (typical hunting radius from nest sites), and foraging cover types selected within the buffer. Assume suitable nesting and foraging habitats (water, Ponderosa Pine stands, etc.), are defined within these breeding habitat parameters. Although water is important component of breeding habitat, cliff sites were considered more limiting; buffering major water courses would have resulted in gross overestimation of habitat. Nest location records not currently available to include in model. Model focus was conservative but in current form likely overestimates suitable breeding habitat in known breeding areas, as local cliff geology / morphology and human disturbance information is not available. Suitable habitat in isolated, historical breeding areas in northeast Montana is not represented in current predicted habitat distribution.

References

Cade, T. J., et al. editors. 1988. Peregrine falcon populations: their management and recovery. The Peregrine Fund, Inc., Boise, Idaho. 949 pp.

Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.

Evans, D. L. 1982. Status reports on twelve raptors. U.S. Department of the Interior, Fish and Wildlife Service, Special Scientific Report No. 238. 68 pp.

Grebence, B. L. and C. M. White. 1989. Physiographic characteristics of peregrine falcon nesting habitat along the Colorado River system in Utah. *Great Basin Nat.* 49:408-418.

Hendricks, P., and K. H. Dueholm. 1995. Cliff-nesting raptor survey of the Sioux District, Custer National Forest: 1994. Unpubl. report to U.S.D.A. For. Serv., Custer N.F., Billings. 20 pp.


Herman, M., and E. E. Willard. 1978. Peregrine falcon and its habitat. U.S.D.A. Forest Service Northern Region. 11 pp.

Johnsgard, P.A. 1990. Hawks, eagles, and falcons of North America. Smithsonian Institution Press, Washington, D.C. 403 pp.

Porter, R. D., M. A. Jenkins, and A. L. Gaski. 1987. Working bibliography of the peregrine falcon. *Natl. Wildl. Fed., Sci. and Tech. Ser. No. 9.* 185 pp.

Snow, C. 1972. Habitat management series for endangered species, Report No. 1: American Peregrine Falcon and Arctic Peregrine Falcon. Bureau of Land Management and the U.S. Dept. of the Interior, Tech. Note: T-N-167. 35 pp.

U.S. Fish and Wildlife Service. 1984. American Peregrine Falcon Rocky Mountains/southwest population recovery plan. Prepared in cooperation with the American Peregrine falcon recovery team. U.S. Fish Wildl. Serv., Denver. 105 pp.



WHITE-TAILED PTARMIGAN

Lagopus leucurus


State rank S2S3

Element code ABNLC10030

Global rank G5

Modeled by C. Tobalske, W. Williams

MTNHP status watch list



State range

Currently found commonly in Glacier National Park and uncommonly in the Swan Range and Mission Mountains of the Bob Marshall/Scapegoat Wilderness complex (Montana Bird Distribution Committee 1996, Wright 1996). Old, questionable records exist for the Beartooth Mountains and Yellowstone National Park, but are not considered reliable (Montana Bird Distribution Committee 1996, Wright 1996).

Habitat description

The White-tailed Ptarmigan is a bird of alpine tundra in all seasons, at or above treeline (Scott 1982); typical habitat consists of snow, steep slopes, rocky areas, valleys, and glacial cirques (Chrest 1971). In Glacier National Park, its habitat consists of glacially leveled benchland covered with small ledges, and ranges from dry, rocky sites to wet and mossy streamside areas; most abundant on level or gently sloping grounds (Choate 1963). This species is not found in timber or shrubby vegetation taller than 18", or in boggy areas without rock cover (Choate 1963). Scott (1982) searched 72 areas in Montana and found birds at 8 sites in Glacier NP and at one site in the Mission Mountains, at elevations ranging from 1951 m to 2795 m. Ptarmigans favored moist habitats and used tundra with few herbaceous plants, more conifers and streams, such as krummholz habitat.

Model assumptions & caveats

References

Braun, C. E. 1993. White-tailed ptarmigan habitat investigations in northeast Oregon. *Oregon Birds* 19:72-73.

Choate, T. S. 1963. Habitat and population dynamics of white-tailed ptarmigan in Montana. *J. Wildl. Manage.* 27:684-699.

Chrest, H. 1971. White-tailed Ptarmigan. Pages 181-183 in T.W. Mussehl and F.W. Howell, eds. Game management in Montana. Montana Fish and Game Dept., Game Manage. Div., Helena, MT.


Frederick, G. P., and R. J. Gutierrez. 1992. Habitat use and population characteristics of the white-tailed ptarmigan in the Sierra Nevada, California. *Condor* 94:889-902.

Giesen, K. M., and C. E. Braun. 1992. Winter home range and habitat characteristics of white-tailed ptarmigan in Colorado. *Wilson Bull.* 104:263-272.

Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Scott, D. M. 1982. Distribution and habitat use of white-tailed ptarmigan in Montana. *Proc. Mont. Acad. Sci.* 41:57-66.



SAGE GROUSE

Centrocercus urophasianus


State rank S5

Element code ABNLC12010

Global rank G5

Modeled by C. Tobalske, W. Williams

MTNHP status --



Predicted habitat: 4,534,810 ha, 11.91 % of state.

State range

Lives year-round locally in southwestern, central, and eastern Montana, east of the Continental Divide; does not occur in mountainous areas west of the Divide (Wallestad 1975, Montana Bird Distribution Committee 1996).

Habitat description


A species strongly associated with mature sagebrush communities such as Big Sagebrush, Black Sage, and Bitterbrush (Dalke et al. 1963, Ehrlich et al. 1988, Klott and Lindzey 1990). In central Montana, Wallestad and Pyrah (1974) found 14 nests in areas with sagebrush canopy cover greater than 15%; this cover also preferred by broods (Wallestad 1971). In north-central Montana, wetlands are used for leks when they occur within sagebrush/grassland habitat; riparian areas, especially near hayfields, are used for brood rearing and also feeding in late summer when forbs become dessicated (H. Nyberg pers. comm.). In southwestern Montana, broods preferred sagebrush shorter in height, adults found in sagebrush consistently taller (Martin and Pyrah 1971). In Colorado, nest success and brood survival was related to the amount of residual grass/forb cover, and in winter survival was found to be dependent upon amount of available sagebrush (Braun 1995). Tall sagebrush on flat or slightly rolling terrain is used in winter in Montana (Eng and Schladweiler 1972, Wallestad and Pyrah 1974, Wallestad 1975). The original range of the Sage Grouse was similar to the distribution of sagebrush, but by the mid 1930s agricultural development and livestock grazing converted/destroyed 50% of this range (Martin and Pyrah 1971). Currently, Sage Grouse are found in sagebrush-grasslands of eastern, central, and southwestern Montana; attempts to establish populations west of the Divide and at former sites have been unsuccessful (Martin and Pyrah 1971).

Model assumptions & caveats

Canopy cover information not available for grass/herbaceous cover to include in model.

References

- Braun, C.E. 1995. Distribution and status of Sage Grouse in Colorado. *Prairie Nat.* 27(1): 1-8.
- Braun, C. E., T. Britt, and R. O. Wallestad. 1977. Guidelines for maintenance of sage grouse habitat. *Wildl. Soc. Bull.* 5:99-106.
- Dalke, P. D., D. B. Pyrah, D. C. Stanton, J. E. Crawford, and E. F. Schlatterer. 1963. Ecology, productivity, and management of sage grouse in Idaho. *J. Wildl. Manage.* 27:811-841.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds.* Simon and Shuster, Inc., New York. xxx + 785 pp.
- Eng, R. L., and P. Schladweiler. 1972. Sage grouse winter movements and habitat use in central Montana. *J. Wildl. Manage.* 36:141-146.
- Kantrud, H.A. and R.L. Kologiski. 1982. Effects of soils and grazing on breeding birds of uncultivated upland grasslands of the northern Great Plains. U.S.D.I., Fish and Wildl. Serv., Wildl. Res. Rep. 15. 33 pp.
- Klott, J. H., and F. G. Lindzey. 1990. Brood habitats of sympatric sage grouse and Columbian sharp-tailed grouse in Wyoming. *J. Wildl. Manage.* 54:84-88.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Wallestad, R. 1975. Life history and habitat requirements of the sage grouse in central Montana. *Mont. Dept. Fish and Game.* 65 pp.
- Wallestad, R., and D. Pyrah. 1974. Movements and nesting of sage grouse hens in central Montana. *J. Wildl. Manage.* 38:630-633.
- Wallestad, R. O. 1971. Summer movements and habitat use by sage grouse broods in central Montana. *J. Wildl. Manage.* 35: 129-136.



COLUMBIAN SHARP-TAILED GROUSE

Tympanuchus phasianellus columbianus


State rank S1

Element code ABNLC13033

Global rank G4T3

Modeled by C. Tobalske, W. Williams

MTNHP status special concern



Predicted habitat: 1,165,422 ha, 3.06 % of state.

State range

The Sharp-tailed Grouse subspecies occurring west of the Continental Divide, populations known as recently as 1970 in the Tobacco Plains area, Blackfoot River Valley, Flathead Valley, and Centennial Valley; recent populations persisting only in the Tobacco Plains and Blackfoot River Valley (Brown 1971, Deeble 1996). Recent status of these populations is discussed in Bown (1980) and Deeble (1994, 1996). Historically, range covered most intermountain valleys in western Montana (Brown 1971, Bown 1980).

Habitat description

A species favoring mesic shrubsteppe and grassland habitats, particularly fescue-wheatgrass and sagebrush-grass associations (Brown 1971, Deeble 1996). In Idaho, Big Sagebrush and Low Sagebrush are preferred nesting cover types, on relatively flat terrain (Marks and Marks 1987), but the breeding habitat can also contain bitterbrush, snowberry, alfalfa, and Crested Wheatgrass (Meints et al. 1992). In Colorado, summer habitat consists of shrub-steppe with a diversity of forbs and bunchgrasses (Giesen and Connelly 1993), whereas in Wyoming nests are located in sagebrush-snowberry and mountain shrub associations (Klott and Linzey 1990). In northwestern Montana, palouse prairie (Bluebunch wheatgrass and rough fescue) habitats used for nesting and brood rearing; brushy draws and hardwood riparian areas in fall/winter (H. Nyberg pers. comm.). In the Tobacco Valley of Montana, nesting and brood rearing occur in areas dominated by grass, with all nests in dense native grass; agricultural areas avoided (Cope 1992). In the Blackfoot River Valley, leks primarily used habitats dominated by Big Sagebrush and Intermediate Wheatgrass (Deeble 1996). Structural diversity is a fundamental habitat component — a mosaic of well-developed forbs, grasses, deciduous trees and shrub/grass communities is optimal — pure stands of any single community unsatisfactory (Deeble 1996). Winter habitat requirements are somewhat narrow; availability of winter habitat is critical in determining the suitability of an area to support sharptails (Giesen and Connelly 1993, Deeble 1996). Mountain shrubs (chokecherry, juniper, snowberry, serviceberry) and riparian shrubby areas provide essential food and thermal cover in winter (Marks and Marks 1987, Meints et al. 1992,

Giesen and Connelly 1993). Columbian Sharptail Grouse are highly philopatric, the longest recorded movements ~20 km between lek and winter range (Meints 1991 in Deeble 1996). Hunting and overgrazing at the turn of the century caused initial population declines, followed by ongoing habitat changes that have greatly reduced suitable habitat: conversion of shrub-steppe to agricultural lands; fire suppression; direct and indirect effects of agricultural chemical use; continued overgrazing (Deeble 1996). Populations in Montana persist only where major, nonfragmented stands of bunch grasses, shrubs and native prairie remain; attempts to establish populations in former range areas or augment current ones, have generally been unsuccessful (Deeble 1996).


Model assumptions & caveats

Model focus is breeding habitat, however suitable winter habitat is defined within breeding habitat parameters. Interspersion of cover types could not be resolved at mapping scale used. Extent of agricultural land use, livestock grazing, and fire management information not available to include in model. Basic habitat requirements demand finer scale resolution. Suitable habitat likely overestimated.

References

- Aldrich, J. W. 1963. Geographic distribution of American Tetraonidae. *J. Wildl. Manage.* 27:529-545.
- Bown, R.R. 1980. The status of the Columbian Sharp-tailed Grouse in the Tobacco Plains, Eureka, Montana. B.S. Thesis. University of Montana, Missoula. 42 pp.
- Brown, R.L. 1971. Sharp-tailed Grouse. Pages 129-133 in T.W. Mussehl and F.W. Howell, eds. *Game management in Montana*. Montana Fish and Game Dept., Game Manage. Div., Helena, MT.
- Cope, M. G. 1992. Distribution, habitat selection and survival of transplanted Columbian sharp-tailed grouse (*Tympanuchus phasianellus columbianus*) in the Tobacco Valley, Montana. M.S. thesis. Montana State University. 60 pp.
- Deeble, B. 1994. Conservation assessment for Columbian sharp-tailed grouse in Montana's Blackfoot valley. Draft report. University of Montana, Missoula. 11 pp.
- Deeble, B.D. 1996. Conservation of Columbia Sharp-tailed Grouse, with special emphasis on the upper Blackfoot Valley, Montana. M.S. Thesis, University of Montana, Missoula. 70 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds*. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Giesen, K. M. and J. W. Connelly. 1993. Guidelines for management of sharp-tailed grouse habitats. *Wildl. Soc. Bull.* 21:325-333.
- Johnsgard, P. A. 1973. *Grouse and quail of North America*. U. of Nebraska, Lincoln. 553 pp.
- Klott, J. H., and F. G. Lindzey. 1990. Brood habitats of sympatric sage grouse and Columbian sharp-tailed grouse in Wyoming. *J. Wildl. Manage.* 54:84-88.
- Marks, J. S., and V. S. Marks. 1987. Habitat selection by Columbian sharp-tailed grouse in west-central Idaho. *Dep. Int. Bureau of Land Manage., Boise District, ID.* 115 pp.
- Meints, D. R., J. W. Connelly, K. P. Reese, A. R. Sands, and T. P. Hemker. 1992. Habitat suitability index procedure for Columbian sharp-tailed Grouse. *Idaho For. Wildl. and Range Exp. Sta. Bull.* 55, Moscow, ID. 27 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's *Montana Bird Distribution, Fifth Edition*. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Oedekoven, O. O. 1985. Columbian sharp-tailed grouse population and habitat use in south central Wyoming. M.S. Thesis, Univ. Wyoming.



YELLOW RAIL

Coturnicops noveboracensis


State rank S1B,SZN

Element code ABNME01010

Global rank G4

Modeled by C. Tobalske, W. Williams

MTNHP status special concern



State range

Few records — main breeding range is north and east of Montana; occurs regularly in the northeastern corner of the state; is rare elsewhere (Montana Bird Distribution Committee 1996, Wright 1996). Secretive species making detection difficult — number of sightings increasing; potential breeding in northern/northeastern areas of the state (T. McEneaney pers. comm.)

Habitat description

Within breeding range (mainly east of the Rockies) are very locally distributed, inhabiting freshwater and brackish marshes, preferring slightly higher, drier margins, and wet meadows (Ehrlich et al. 1988, Bookhout 1995). In the upper peninsula of Michigan, yellow rails select seasonally flooded sedge meadow marshes (Bookhout and Stenzel 1987). In southern Oregon they breed in wet mountain meadows (1266-1524 m) located near a spring, a seep, a creek, or in a river floodplain; vegetation consists of broadleaf sedge associations, with Lodgepole, Ponderosa Pine, aspen and white fir growing around the meadows (Stern et al. 1993). In Montana, breeds in wet sedge meadows dominated by *Carex* (rarely found in cattail), with moist substrate and standing water — quality of the habitat is diminished by presence of woody species, such as willow (Bookhout 1995). Nests beneath dead, procumbent vegetation, situated on ground or up to 15 cm above it (Bookhout 1995). Within marshes and wetlands of breeding range, successful nesting requirements poorly understood; secretive nature precludes adequate population numbers and distribution estimates (Bookhout 1995).

Model assumptions & caveats

Suitable breeding habitat for this species is likely overestimated - hydrography coverage may not be consistently accurate/true to mapping scale used; persistence of intermittent water (e.g., seasonal vs. semipermanent, etc.) information not available; interspersed of emergent vegetation with open water and standing water within emergent stands could not be resolved at mapping scale used.

References

Bookhout, T.A. 1995. Yellow Rail. in *The Birds of North America*, No. 139. A. Poole and F. Gill, eds. The Academy of Natural Sciences, Philadelphia and The American Ornithologist's Union, Washington, D.C.

Bookhout, T. A., and J. R. Stenzel. 1987. Habitat and movements of breeding yellow rails. *Wilson Bull.* 99:441-447.


Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds.* Simon and Shuster, Inc., New York. xxx + 785 pp.

Griese, H. J., R. A. Ryder, and C. E. Braun. 1980. Spatial and temporal distribution of rails in Colorado. *Wilson Bull.* 92:96-102.

Montana Bird Distribution Committee. 1996. P. D. Skaar's *Montana Bird Distribution*, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Stern, M. A., J. F. Morawski, and G. A. Rosenberg. 1993. Rediscovery and status of a disjunct population of breeding yellow rails in Southern Oregon. *Condor* 95:1024-1027.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



WHOOPING CRANE

Grus americana


State rank SZN

Element code ABNMK01030

Global rank G1

Modeled by C. Tobalske, W. Williams

MTNHP status special concern



State range

This species is not frequently observed (four sightings 1991-1995); transient records are scattered throughout the state, east of the Continental Divide; most observations in the extreme northeastern corner of Montana (Montana Bird Distribution Committee 1996, T. McEneaney pers comm.).

Habitat description

One of the rarest birds in North America, the Whooping Crane has recovered from 16 individuals in 1941 to three current wild populations — the only natural, wild population nests in the Northwest Territories and adjacent areas of Alberta (Lewis 1995). The Rocky Mountain population consists of a small number of individuals cross-fostered by Sandhill Cranes that summer in southeastern Idaho, western Wyoming, and southwestern Montana — in the Greater Yellowstone Ecosystem, the birds use shallow wetlands, wet meadows, and adjacent upland sites with minimal human disturbances (Drewien *in* Clark et al. 1989, Lewis 1995). The Northwest Territories population migrates from nesting grounds through northeastern Alberta, south-central Saskatchewan, northeastern Montana, and western North Dakota en route to wintering grounds in Texas (Lewis 1995). In Colorado, a study conducted along the Platte River found the species roosting in wide river channels (155 to 365 m) with low, exposed, bare sandbars, shallow water, and isolation from human disturbance — sites farther than 400 m from roads and houses (U.S. Fish and Wildlife Service 1981). Feeding sites were usually within 4.8 km of the river, in a variety of habitats; 80% were in upland agricultural lands (emerging small grains, small grain stubbles). Feeding habitat was devoid of tall trees or dense shrubs; vegetation shorter than 30 cm, topography generally flat — cranes also fed in wetlands with water lower than 60 cm, and very short or sparse emergent vegetation (U.S. Fish and Wildlife Service 1981). Armbruster (1990) characterized migration habitat as including mainly sites with good visibility, water depth of 30 cm or less, and minimum wetland size of 0.04 ha for roosting. The Northwest Territories population is somewhat flexible in habitat use during migration, generally feeding in a variety of croplands and roosting in freshwater, shallow wetlands — most wetlands used for roosting were <4 ha and were within 1 km of a suitable feeding area (Lewis 1995). Family groups fed mostly at


wetland sites; nonfamily groups fed primarily in croplands (Lewis 1995). The central Platte River, Nebraska, was historically a major fall stopover — the wide, shallow river area provided roosting sites adjacent to wetlands and cropland feeding sites (Allen 1952 in Lewis 1995). The Rocky Mountain population winters in the Rio Grande Valley; stopover in the San Luis Valley, Colorado to roost in ponds, streams, wet meadows and feed in nearby agricultural fields, particularly barley (Lewis 1995). Whooping Cranes are very sensitive to human perturbations and slight disturbance will often cause adults to desert nests — losses also suffered from illegal shooting, powerline collisions, barbed wire entanglement and avian tuberculosis (Drewien in Clark et al. 1989).

Model assumptions & caveats

No information on human disturbance/land use information available to include in model.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Armbruster, M. J. 1990. Characterization of habitat used by whooping cranes during migration. U.S. Fish and Wildl. Serv. Biol. Rep. 90(4). 16 pp.
- Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.
- Lewis, J.C. 1995. Whooping Crane. in *The Birds of North America*, No.153. A. Poole and F. Gill, eds. The Academy of Natural Sciences, Philadelphia and The American Ornithologist's Union, Washington, D.C.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- U. S. Fish and Wildlife Service. 1981. The Platte River ecology study. N. Prairie Wildl. Res. Ctr., Jamestown. 187 pp.



PIPING PLOVER

Charadrius melodus


State rank S2B,SZN

Element code ABNNB03070

Global rank G3

Modeled by W. Williams

MTNHP status special concern



Predicted habitat: 56,912 ha, 0.15 % of state.

State range

Breed extremely locally in northeastern Montana near/in Charles M. Russell NWR, Bowdoin NWR, Nelson Reservoir, Medicine Lake NWR, and the Missouri River below Fort Peck dam; has bred in northcentral Montana in Pondera county (Carlson and Skaar 1976 in MNHP-VCA 1996; Montana Piping Plover Recovery Committee 1991, 1992, 1993, 1994 in MNHP-VCA 1996; Montana Bird Distribution Committee 1996). Transient sightings very rare; no winter records in Montana (Montana Bird Distribution Committee 1996).

Habitat description

Breed in three disjunct areas: Atlantic coastal beaches; sand or gravel shorelines of the Great Lakes; sand, gravel, or alkaline shores of lakes and rivers in the Great Plains (Gaines and Ryan 1988, Reel et al. 1989). Throughout the breeding range, wide beaches with minimal vegetation cover are used for nesting (A.O.U. 1983, Gaines and Ryan 1988). Inland, breed on open/exposed areas/beaches of sand, pebble or gravel, especially those with scattered clumps of vegetation that are associated with shores of islands, rivers, ponds, lakes, reservoirs, and alkaline wetlands (Johnsgard 1981, Ehrlich et al. 1988, Gaines and Ryan 1988, Ziewitz et al. 1992). Prefer salt-encrusted, sandy, gravelly, or pebble-mudded shorelines of alkaline wetlands; larger alkaline lakes of Canada and smaller subsaline, semipermanent potholes of the Great Plains represent optimal habitat (Stewart 1975, Johnsgard 1981, Kantrud and Stewart 1984, Weber and Martin 1991, Espie et al. 1996). Avoid vegetated areas or shorelines; on Atlantic beaches nesting areas were abandoned with encroachment of vegetation (Wilcox 1959). At Lake Diefenbaker, Saskatchewan, an important North American breeding lake, select beaches with greatest proportion of gravel/stones; nests located 50-75m from nearest water (Espie et al. 1996). On the Platte River, NE, favor sparsely vegetated, larger, higher sandbars in wider river channels - river reaches at least 1200m long with 275-400m widths, and at least 1.45ha of dry, sandbar or open, permanent islands represent optimal habitat (Ziewitz et al. 1992). In the northern grasslands of MT, ND, SD, and Manitoba, most nests found on unvegetated gravel or sand

substrates, at wetland edges surrounded by native grassland pasture (Kantrud and Higgins 1992). Use beaches of alkaline wetlands in northern Great Plains despite precipitation conditions; 60% use shorelines around alkaline lakes (Haig and Plissner 1993) - even in dry years will return to the same breeding sites, as basins with very little water are still productive (Weber and Martin 1991). River habitat is important (20% of the Great Plains population use river nest sites), but fluctuating/high water levels limit nest site availability and wave action may destroy nests (Gaines and Ryan 1988, Reel et al. 1989, Haig and Plissner 1993). At Lake Diefenbaker, a dammed reservoir, nesting was precluded after high water levels inundated shoreline habitat; normal nesting numbers still not reestablished two years later, even though ample suitable habitat was available (Goossen 1990). In the Chain-of-Lakes area, ND, beach territories average ~25m in width; narrower beaches not selected due to greater wave action and predation threat (Gaines and Ryan 1988). Established territories are characterized by more gravel, but sparse, scattered clumps of vegetation provide nest camouflage (Gaines and Ryan 1988). In the Great Plains area often nest in or adjacent to Least Tern or American Avocet colonies (MNHP-VCA 1996). Forage for worms, crustaceans, mollusks, insects, and marine invertebrates (Ehrlich et al. 1988, Reel et al. 1989). During migration and winter, use uplands, pond margins, flooded fields, and mudflats (Ehrlich et al. 1988, Reel et al. 1989). Populations have declined steadily in the last century, and migration routes of inland birds are poorly understood (Gaines and Ryan 1988, Haig and Plissner 1993). In winter, use primarily coastal habitats - ocean beaches, or sand/algal flats in protected bays (MNHP-VCA 1996). Piping Plovers are federally endangered in the Great Lakes region, threatened elsewhere in their breeding range, and have been extirpated from five central and eastern states due to human disturbance and habitat destruction (Gaines and Ryan 1988, Reel et al. 1989). Population growth may be limited by nest predation and lack of nesting habitat in wet years, but effects of human activities are an important factor in population declines (Flemming et al. 1988, Gaines and Ryan 1988). Human disturbances destroy nest sites, preclude nesting in suitable areas, and adversely affect fledging success - exclusion of human recreation and cattle from breeding habitat (minimum buffer of 160m around nesting areas), is strongly recommended for the preservation and enhancement of breeding populations (Flemming et al. 1988, Gaines and Ryan 1988, Reel et al. 1989).


Model assumptions & caveats

A TES species in Montana, breeding habitat modeled only - inclusion of migration habitat would diminish relevance of model as regards breeding habitat requirements. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Persistence of intermittent water (seasonal vs. semipermanent, etc.), alkalinity, and salinity information not available to include in model. Resolution of beach characteristics and morphology not obtainable at mapping scale used. Assume adequate nesting habitat defined within selected, buffered hydrography.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Espie, R.H.M., R.M. Brigham and P.C. James. 1996. Habitat selection and clutch fate of Piping Plovers (*Charadrius melodus*) breeding at Lake Diefenbaker, Saskatchewan. *Can. J. Zool.* 74(6): 1069-1075.
- Faanes, C. A. 1983. Aspects of the nesting ecology of least terns and piping plovers in central Nebraska. *Prairie Nat.* 15(4): 145-154.
- Flemming, S. P. et al. 1988. Piping plover status in Nova Scotia related to its reproductive and behavioral responses to human disturbance. *J. Field Ornithol.* 59:321-330.
- Gaines, E. P. and M. R. Ryan. 1988. Piping plover habitat use and reproductive success in North Dakota. *J. Wildl. Manage.* 52:266-273.

- Goossen, J. P. 1990. Piping plover research and conservation in Canada. *Blue Jay* 48:139-153.
- Haig, S. M. and J. H. Plissner. 1993. Distribution and abundance of piping plovers: results and implications of the 1991 international census. *Condor* 95:145-156.
- Johnsgard, P. 1981. *The Plovers, Sandpipers, and Snipes of the World*. University of Nebraska Press, Lincoln, Nebraska. 493 pp.
- Kantrud, H.A. and K.F. Higgins. 1992. Nest and nest site characteristics of some ground-nesting, non-passerine birds of northern grasslands. *Prairie Nat.*, 24(2): 67-84.
- Kantrud, H.A. and R.E. Stewart. 1984. Ecological distribution and crude density of breeding birds on prairie wetlands. *J. Wildl. Manage.* 48(2): 426-437.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Reel, S., L. Schassberger and W. Ruediger. 1989. *Caring for Our Natural Community: Region 1 - Threatened, Endangered and Sensitive Species Program*. USDA, Forest Service Northern Region Wildlife and Fisheries.
- Stewart, R.E. 1975. *Breeding birds of North Dakota*. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.
- Weber, L.M. and B.H. Martin. 1991. Piping Plovers nest on dry and nearly dry alkaline wetlands in the northern Great Plains 1988-1990. *Prairie Nat.*, 23(4): 209.
- Wilcox, L. 1959. A twenty year banding study of the Piping Plover. *Auk* 75:129-152.
- Ziewitz, J.W., J.G. Sidle and J.J. Dinan. 1992. Habitat conservation for nesting Least Terns and Piping Plovers on the Platte River, Nebraska. *Prairie Nat.*, 24(1): 1-20.



MOUNTAIN PLOVER

Charadrius montanus


State rank S2B,SZN

Element code ABNNB03100

Global rank G2

Modeled by W. Williams

MTNHP status special concern



Predicted habitat: 875,932 ha, 2.30 % of state.

State range

Currently breed in central, north-central, and southwest Montana (FaunaWest Wildlife Consultants 1991 in MNHP-VCA 1996, Bergeron et al. 1992). Historic range probably encompassed most of eastern and parts of southwest MT; records are summarized in FaunaWest Wildlife Consultants (1991). Transitory through the Greater Yellowstone Ecosystem, sightings rare in Montana; overwintering does not occur (Knowles in Clark et al. 1989, Montana Bird Distribution Committee 1996).

Habitat description

High and arid plains and shortgrass prairie (particularly Blue Grama-Buffalo Grass communities, although they are very limited in Montana), is primary habitat; secondarily use drier, open shrubland and sandy, semi-arid flats (D. Casey pers. comm., Johnsgard 1981, Hayman et al. 1986, Dobkin, 1994). Flat, open, short (<10cm tall) grasslands with little shrub cover are preferred; nests are simple scrapes usually amidst scattered clumps of cacti or Western Wheatgrass, often near conspicuous objects, such as piles of dried cow manure or similarly-sized rocks (Graul 1975, Olson and Edge 1985, Knowles in Clark et al. 1989, DeGraaf et al. 1991). Typically found in intensively grazed areas - bare ground cover of 30% is considered a minimum breeding habitat requirement (Wallis and Wershler 1981, Knowles et al. 1982, Parrish et al. 1993, Knopf and Miller 1994). Consistent use of areas with bare ground, during breeding and overwintering, indicate it is an important year-round habitat component (Knowles et al. 1982, Prellwitz 1993, Knopf and Rupert 1996). In Canada, nesting habitat is characterized by extensive tracts of heavily grazed pastures (burned or unburned), with short (<8cm), sparse grassland vegetation on level to slightly undulating terrain (Wallis and Wershler 1981, Parrish et al. 1993). In CO and WY similar grazed habitats are used for breeding - shortgrass areas dominated by Blue Grama-Buffalo Grass, and sparse grass/ forb habitat, respectively (Knowles et al. 1982, Parrish et al. 1993). In MT, selectively breed in Black-tailed Prairie Dog towns - grazing pressure of prairie dogs and local livestock keep vegetation cover low, providing greater bare ground and more horizontal visibility providing optimal nesting habitat (Olson and Edge 1985, Knowles et al.

1982). In the Charles M. Russell National Wildlife Refuge (CMRNWR), select prairie dog towns at least 6ha in size (≥ 10 ha preferred), in upland, level areas, characterized by short vegetation and moderate to heavy grazing pressure by cattle (Knowles et al. 1982, Olson and Edge 1985). Often nest far from water, however, prairie dog towns are usually close to livestock watering ponds (Knowles et al. 1982). In northeastern MT, sighted on habitat similar to that found in Black-tailed Prairie Dog towns, but on CMRNWR lands Mountain Plovers did not nest in heavily grazed areas outside of towns (Knowles et al. 1982, Prellwitz 1993). A minimum area of 28ha is needed to raise a brood (Knopf and Rupert 1996). Brood rearing habitat includes areas with forbs or objects such as fence posts for shade; in more disturbed prairie, broods use cattle watering and loafing sites, fallow agricultural fields (Knopf and Rupert 1996). Forage primarily on insects - grasshoppers, beetles, crickets (Ehrlich et al. 1988, DeGraaf et al. 1991). Often seen in flocks during migration and winter; use freshly tilled and recently burned fields, semi-desert or dry agricultural habitats, alkaline flats, heavily grazed grasslands (Hayman et al. 1986, Knopf and Rupert 1996). Populations declined 63% between 1966 and 1991 in North America, and continued to do so through 1993, due to habitat degradation on both breeding and wintering grounds (Page and Gill 1994, Knopf and Rupert 1996). Breeds across the western Great Plains, as well as locally in WY, CO, NM, and TX, however, Weld Co., CO and Phillips Co., MT provide suitable nesting habitat for over half of the North American population (Knopf and Miller 1994). The Mountain Plover has been extirpated from ND and SD, is endangered in Canada, has been petitioned for candidacy for Threatened and Endangered status in the U.S., and is considered a TES species in MT due to severe range contraction as a result of prairie conversion to agriculture and the extermination of prairie dogs (Johnsgard 1981, Olson and Edge 1985, Dobkin 1994, Page and Gill 1994, Montana Bird Distribution Committee 1996). In the breeding range of MT (CMRNWR lands), heavy cattle grazing alone does not create suitable nesting habitat - in this area, plovers were not sighted outside of prairie dog towns even though potential habitat was surveyed (Olson and Edge 1985). Maintaining prairie dog towns on the CMRNWR is critical to preserving breeding habitat; persistent prairie dog control efforts are detrimental to breeding populations on the refuge and limit suitable breeding habitat outside its boundaries (Olson and Edge 1985, Dobkin 1994).

Model assumptions & caveats


Breeding habitat modeled. Rare migrant, does not overwinter. Some migration habitat assumed defined within breeding habitat parameters, but other habitat excluded from model, as inclusion would diminish relevance of model as regards breeding habitat requirements. Cover type area (territory size) not included in model. Black-tailed Prairie Dog town locations were available, but sizes/area coverage of the towns were not available for all locations. Additionally, Mountain Plovers will breed outside of prairie dog towns in Montana, therefore inclusion of prairie dog towns as a habitat filter would underestimate breeding habitat in these areas. Breeding habitat requirements in Montana appear fairly rigid although are still not completely understood - ample heavily grazed areas are probably available in the state, however, suitable breeding habitat is likely overestimated.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Bergeron, D., C. Jones, D.L. Genter, and D. Sullivan. 1992. P. D. Skaar's Montana Bird Distribution, Fourth Edition. Special Publication No. 2. Montana Natural Heritage Program, Helena. 116 pp.
- Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.

Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and

- Great Plains Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Graul, W. D. 1975. Breeding biology of the mountain plover. *Wilson Bull.* 87:6-31.
- Hayman, P., J. Marchant, and T. Prater. 1986. Shorebirds: an identification guide to the waders of the world. Houghton Mifflin Co., Boston. 412 pp.
- Johnsgard, P. 1981. The Plovers, Sandpipers, and Snipes of the World. University of Nebraska Press, Lincoln, Nebraska. 493 pp.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Knopf, F. L. and B. J. Miller. 1994. *Charadrius montanus* -- montane, grassland, or bare-ground bird? *Auk* 111:504-506.
- Knopf, F. L., and J. R. Rupert. 1996. Productivity and monements of Mountain Plovers breeding in Colorado. *Wilson Bull.* 108:28-35.
- Knowles, C. J., C. J. Stoner, and S. P. Gieb. 1982. Selective use of black-tailed prairie dog towns by mountain plovers. *Condor* 84:71-74.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Olson, S.L. and W.D. Edge. 1985. Nest site selection by mountain plovers in northcentral Montana. *J. Range Manage.* 38(3): 280-282.
- Page, G.W. and R.E. Gill, Jr. 1994. Shorebirds in western North America: late 1800s to late 1900s. *Studies in Avian Biology*, No. 15: 147-160.
- Parrish, T.L., S.H. Anderson and W.F. Oelklaus. 1993. Mountain Plover habitat selection in the Powder River Basin, Wyoming. *Prairie Nat.* 25(3): 219-226.
- Prellwitz, D.M. 1993. Additional Mountain Plover sightings in Montana. *Prairie Nat.*, 25(1): 23-26.
- Wallis, C.A. and C.R. Wershler. 1981. Status and breeding of Mountain Plovers (*Charadrius montanus*) in Canada. *Can. Field-Nat.* 95(2): 133-136.



BLACK-NECKED STILT

Himantopus mexicanus


State rank S2B,SZN

Element code ABNND01010

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Breeding range has expanded into WA, MT, Alberta, and Saskatchewan in response to drought throughout traditional ranges southwest (Page and Gill 1994). Breeding in Montana is somewhat rare and localized in central (Benton Lake NWR has largest known breeding population in Montana; > 100prs) and northeast (Bowdoin NWR) areas (S. Martin pers. comm., Montana Bird Distribution Committee 1996). Transient sightings recorded mainly for western Montana; overwintering does not occur in the state (Montana Bird Distribution Committee 1996).

Habitat description


Breed in a wide range of wetland habitats: coastal saltmarshes and lagoons; fresh or saline inland lakes, ponds, and marshes; wet savannas; mudflats (Hayman et al. 1986, Ehrlich et al. 1988). Found inland around grassy or bare sandy/gravelly shorelines of shallow, freshwater or brackish pools of wetlands with extensive mudflat areas; also flooded fields (Dekker et al. 1979, Johnsgard 1979, Johnsgard 1981). Nests may be exposed or well concealed in vegetation, and are always close to foraging areas (Johnsgard 1979, Ehrlich et al. 1988). Nest in loose colonies, occasionally singly and well outside normal breeding range; often associated with American Avocets (Hayman et al. 1986, Johnsgard 1986). In MT, nest in medium to large wetland complexes of open marshes and meadows; often in alkaline areas (MNHP-VCA 1996). Subsist on insect larvae - especially of brine flies and brine shrimp, and crayfish; also snails, tadpoles, seeds (Ehrlich et al. 1988). Migration and winter habitats similar to that of breeding (Hayman et al. 1986). With the substantial amount of wetlands lost during the past 200 years - over 90% of Black-necked Stilt wetland and breeding habitat in the Central Valley, CA has been destroyed - other populations breeding in North America have undoubtedly declined; the Black-necked Stilt is a TES species in MT (Page and Gill 1994, Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Uncommon migrant; potential inland migration habitat is similar to that of breeding and is assumed defined within breeding habitat parameters. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Sizes/areas of wetland complexes difficult to resolve at mapping scale used. Extent of wetland mudflat areas and wetland alkalinity information not available to include in model.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Dekker, D., R. Lisier, T.W. Thormin, D.V. Weseloh and L.M. Wesloh. 1979. Black-necked Stilts nesting near Edmontaon, Alberta. *Can. Field-Nat* 93(1):68-69.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds*. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hayman, P., J. Marchant, and T. Prater. 1986. *Shorebirds: an identification guide to the waders of the world*. Houghton Mifflin Co., Boston. 412 pp.
- Johnsgard, P. 1981. *The Plovers, Sandpipers, and Snipes of the World*. University of Nebraska Press, Lincoln, Nebraska. 493 pp.
- Johnsgard, P.A. 1979. *Birds of the Great Plains: breeding species and their distribution*. Univ. Nebraska Press, Lincoln. 539 pp.
- Johnsgard, P. A. 1986. *Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region*. Colorado Associated University Press, Boulder. xi + 504 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's *Montana Bird Distribution, Fifth Edition*. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Page, G.W. and R.E. Gill, Jr. 1994. Shorebirds in western North America: late 1800s to late 1900s. *Studies in Avian Biology*, No. 15: 147-160.
- Salt, W.R. and J.R. Salt. 1976. *The birds of Alberta*. Hurtig Publishers, Edmonton, Alberta. xv + 498 pp.



FRANKLIN'S GULL

Larus pipixcan


State rank S3B,SZN

Element code ABNNM03020

Global rank G4G5

Modeled by W. Williams

MTNHP status special concern



State range

Breeds locally, mainly in eastern Montana along the Highline; also Red Rock Lakes NWR in southwestern Montana (Montana Bird Distribution Committee 1996). Migrates widely throughout Montana, most commonly east of the Continental Divide; overwintering does not occur in the state (Montana Bird Distribution Committee 1996).

Habitat description

Breed on freshwater prairie and steppe marshes, sloughs, and marshy lakes, rarely in flooded meadows; also shallow river impoundment marshes, as those managed for waterfowl (Stewart 1975, A.O.U. 1983, Johnsgard 1986, Ehrlich et al. 1988). Sensitive to precipitation conditions and human disturbance, stable colonies depend on extensive, more permanent and remote prairie marshes (Burger and Gochfeld 1994). Nesting requirements are large marshes with emergent vegetation (cattails, bulrushes, phragmites) in water deep enough (usually 30-60cm) to persist until young have fledged; in MN, nests were deserted when water reached 80cm deep (Burger and Gochfeld 1994). Nest on large, floating platforms of plant debris anchored to emergent vegetation, preferably in semi-open emergent cover to provide adverse weather protection, yet adjacent to open water to facilitate escape from predators (Johnsgard 1979, Ehrlich et al. 1988, Burger and Gochfeld 1994). Since frequently nest on larger, seasonal or semipermanent marshes, colonies are vulnerable to drought or wetland drainage, and often shift from year to year in response to these conditions; in some years local populations may not breed (Stewart 1975, Burger and Gochfeld 1994). In MT, stable colonies breed at Bowdoin NWR, Freezout Lake WMA, and Benton Lake NWR (Benton Lake NWR hosts the largest breeding population in MT, with 10,000-12,000 pairs), with less stable colonies at Medicine Lake NWR and Red Rock Lakes NWR (S. Martin, pers. comm., Burger and Gochfeld 1994). White-faced Ibis, Black Terns, Black-necked Stilts, Forster's Terns and Black-crowned Night-Herons are often interspersed within Montana colonies (Burger and Gochfeld 1994). American Coots will compete for nest sites where ranges overlap (Burger 1973). Opportunistic feeders; subsist on adult and larval insects, earthworms,


vegetation (seeds and foliage), fish, small mammals, other invertebrates (Ehrlich et al. 1988, Burger and Gochfeld 1994). During nesting, feed in shallower waters adjacent to nest sites in surrounding meadows; forage in upland areas when grasshoppers are abundant (Littlefield and Thompson 1981). Post breeding dispersal in prairie regions extends in all directions from colony sites; mixed adult and juvenile flocks often follow agricultural machinery to forage on flushed insects (Littlefield and Thompson 1981, Ehrlich et al. 1988, Burger and Gochfeld 1994). Migration routes are generally south through the Great Plains, but some migrate over high tundra in CO (3900-4270m elevation), and over passes in MT (Burger and Gochfeld 1994). Migrating flocks feed in flooded fields, pastures, prairies, croplands; also estuaries, bays, mudflats, lagoons, lakes, and roost on inland lakes, bays, estuaries (Burger and Gochfeld 1994). Past wetland drainage coupled with the severe effects of the Dust Bowl have destroyed many colony sites; once abundant in Canada in the 1980s, many important colonies were lost due to wetland drainage (Burger and Gochfeld 1994). With the creation of national wildlife refuges and the protection of extensive marshes, however, populations are increasing (Burger and Gochfeld 1994). Franklin's Gulls are very sensitive to disturbance, especially early in nesting, and will desert a colony entirely with excessive exposure to humans (Burger and Gochfeld 1994).

Model assumptions & caveats

Breeding habitat modeled. Potential inland migration, post breeding, and foraging habitats defined within breeding habitat parameters. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Persistence of intermittent water (seasonal vs. semipermanent, etc.) and water depth information not available to include in model. Cannot resolve interspersion of emergent vegetation and open water areas at mapping scale used. Potential for human disturbance information not available to include in model. Suitable breeding habitat likely overestimated.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Burger, J. 1973. Competition between American Coots and Franklin's Gulls for nest sites and egg predation by the coots. *Wilson Bull.* 85: 449-451.
- Burger, J. and M. Gochfeld. 1994. Franklin's Gull (*Larus pipixcan*). In: A. Poole and F. Gill, (eds.), *The Birds of North America*, No. 116. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union. 28 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds*. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.
- Johnsgard, P.A. 1979. *Birds of the Great Plains: breeding species and their distribution*. Univ. Nebraska Press, Lincoln. 539 pp.
- Johnsgard, P. A. 1986. *Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region*. Colorado Associated University Press, Boulder. xi + 504 pp.
- Littlefield, C.D. and S.P. Thompson. 1981. History and status of the Franklin's gull on Malheur National Wildlife Refuge, Oregon. *Great Basin Nat.* 41(4):440-444.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Stewart, R.E. 1975. *Breeding birds of North Dakota*. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



CASPIAN TERN

Sterna caspia


State rank S2B,SZN

Element code ABNNM08020

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Breed very locally, mainly in northeast and northwest-central Montana; is an uncommon migrant found widely throughout the state; overwintering does not occur (Montana Bird Distribution Committee 1996).

Habitat description


Breed very locally in the Rocky Mountain region (Johnsgard 1986). Favor less-developed, less-polluted areas of the coast, nesting on sparsely vegetated sand, gravel, or shell beaches, barrier or spoil islands, shell berm in salt marshes; also found inland on sandy or gravelly shorelines or islands of large lakes or reservoirs (Johnsgard 1986, Ehrlich et al. 1988, DeGraaf et al. 1991). Nest in dense colonies; occasionally single pairs will nest among other tern or gull species (Godfrey 1966, Johnsgard 1986, DeGraaf et al. 1991). Nests are shallow depressions in the substrate, often concealed among driftwood (Ehrlich et al. 1988, DeGraaf et al. 1991). In WA, all colonies - whether coastal or inland - nested on sandy substrates; breeding population locations often varying with periods of erosion and deposition of sandy nesting habitat (Penland 1982). Subsist almost entirely on fish; also crayfish, insects, nestlings and eggs of other bird species; rarely carion (Ehrlich et al. 1988, DeGraaf et al. 1991). During migration use large lakes, reservoirs, water courses, large marshes (DeGraaf et al. 1991). Winter habitat primarily marine and estuarine, along beaches, spits - often roost with other birds on bars and beaches (Godfrey 1966, DeGraaf et al. 1991). Very sensitive to human disturbance; intrusions into breeding colonies results in significant reductions in reproduction success (Penland 1982, Ehrlich et al. 1988). The Caspian Tern is considered a threatened, endangered, or sensitive species in Montana, as little is known about the breeding/nesting requirements of this colonial breeder in the state (Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Migration habitat not specifically modeled; assume some potential inland migration habitat is defined within breeding habitat parameters. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Since potential for human disturbance information is not available to include in model, selected narrow hydrography buffer (90m) and only those permanent lakes and marshes having islands to represent suitable, predicted breeding habitat that is less likely to be disturbed. Island coverage, however, may not be accurately represented within the hydrography coverage.

References

- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Godfrey, W. Earl. 1966. The birds of Canada. National Museums of Canada. Ottawa. 428 pp.
- Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Penland, S. 1992. Distribution and status of the Caspian Tern in Washington State. Murrelet 63: 73-79.



COMMON TERN

Sterna hirundo


State rank S3B,SZN

Element code ABNNM08070

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Breed locally, mainly in northern and central areas of Montana (Montana Bird Distribution Committee 1996). Transient sightings uncommon and scattered; overwintering does not occur (Montana Bird Distribution Committee 1996).

Habitat description

Breed on coastal sandy or gravelly beaches, small islands in salt marshes, or small rocky islands in lakes, rivers, reservoirs (Stewart 1975, DeGraaf et al. 1991). In the Rocky Mountain region, breed primarily on lakes in the plains, rarely in the montane parks (Johnsgard 1986). Nest colonially on isolated, sparsely vegetated islands in large lakes and reservoirs; in ND, shallow river impoundments managed for waterfowl are also used (Stewart 1975, DeGraaf et al. 1991). Frequently nest in low, sparse vegetation; nests are well built hollow mounds of vegetation or simple depressions in sandy substrates (Godfrey 1966, DeGraaf et al. 1991). In Saskatchewan, prefer very low, small, rocky islands with sparse vegetation; nests usually located near the lower edge of vegetation (Stelfox and Brewster 1979). May not nest near an adequate food source, and often travel 10+ miles to nearby lakes in search of food (Stewart 1975, Pinkowski 1977). Forage primarily for fish and crustaceans; also aquatic worms, insects (Ehrlich et al. 1988, DeGraaf et al. 1991). Winter habitat is typically along shorelines, rocky coasts, bays - over shallow, coastal waters (DeGraaf et al. 1991). The Common Tern is a species of special concern, on the Blue List; populations in the Great Lakes are declining due to nest site loss from a rise in water levels and increased nest site competition by increasing populations of Ring-billed Gulls (Ehrlich et al. 1988). In MT, due to a need for information regarding breeding/nesting requirements, this colonial nester is considered a threatened, endangered, or sensitive species (Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Migration habitat not specifically modeled; assume potential inland migration habitat is defined within breeding habitat parameters. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Unable to resolve island characteristics in the land cover layer at mapping scale used. Island coverage may not be accurately represented within the hydrography layer.

References

DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.

Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.

Godfrey, W. Earl. 1966. The birds of Canada. National Museums of Canada. Ottawa. 428 pp.

Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.


Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Pinkowski, B. C. 1980. Adaptations of Common Terns nesting on an inland reservoir. *Prairie Nat.* 12(3&4): 111-113.

Stelfox, H.A. and G.J. Brewster. 1979. Colonial-nesting Herring Gulls and Common Terns in northeastern Saskatchewan. *Can. Field-Nat.* 93(2): 132-138.

Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



FORSTER'S TERN

Sterna forsteri


State rank S2B,SZN

Element code ABNNM08090

Global rank G5

Modeled by W. Williams

MTNHP status special concern



State range

Breeding is known at 6 scattered locations in Montana; migration occurs throughout the state, although off the breeding areas, few are seen; overwintering does not occur (Montana Bird Distribution Committee 1996).

Habitat description


Breed primarily on large saltwater and freshwater marshes with extensive stands of emergent vegetation; in the Rocky Mountain region are mainly associated with shallow prairie marshes (Johnsgard 1986, Ehrlich et al. 1988, DeGraaf et al. 1991). A variety of habitats are suitable for nesting: marshy areas of bays, ponds, lakes, sloughs, and river impoundments; muddy lake or estuarine islands dominated by emergent vegetation; dikes in evaporation ponds; vegetated dredge spoil islands (Stewart 1975, Scharf and Shugart 1984, DeGraaf et al. 1991). Usually nest in colonies, often with Common and Black Terns, and often associated with Yellow-headed Blackbirds; American Coots and Red-necked Grebes occasionally parasitize nests (Bergman et al. 1970, Scharf and Shugart 1984, Ehrlich et al. 1988). If present, muskrat houses near open pools of water are especially favored for nesting; also nest on compact, usually floating, platforms of emergent/submergent vegetation; sometimes in a simple sand or mud depression; occasionally use abandoned nests of Western or Pied-billed Grebes (Bergman et al. 1970, Stewart 1975, Johnsgard 1986, DeGraaf et al. 1991). Essentially piscivorous; also eats insects, crustaceans, amphibians (DeGraaf et al. 1991). Winter habitat is primarily coastal - harbors, marshy bays, estuaries, lagoons; occasionally inland ponds and lakes (DeGraaf et al. 1991). The Forster's Tern is considered a threatened, endangered, or sensitive species in Montana - conversion of wetlands for agricultural purposes may seriously affect this marshland breeder; more information is needed on the breeding/nesting requirements of this species (Ehrlich et al. 1988, Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Migration habitat not specifically modeled; assume potential inland migration habitat is defined within breeding habitat parameters. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Persistence of intermittent water (seasonal vs. semipermanent, etc.), water depth, presence of muskrat houses, and submerged vegetation information not available to include in model. Interspersion of emergent vegetation near/in water cannot be resolved at mapping scale used. Suitable habitat likely overestimated.

References

- Bergman, R. D., P. Swain, and M. W. Weller. 1970. A comparative study of nesting Forster's and black terns. *Wilson Bull.* 82:435-444.
- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds.* Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.
- Johnsgard, P. A. 1986. *Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region.* Colorado Associated University Press, Boulder. xi + 504 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Scharf, W.C. and G.W. Shugart. 1984. Distribution and phenology of nesting Forster's Terns in eastern Lake Huron and Lake St. Claire. *Wilson Bull.* 96(2): 306-309.
- Stewart, R.E. 1975. *Breeding birds of North Dakota.* Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



LEAST TERN

Sterna antillarum


State rank S1B,SZN

Element code ABNNM08100

Global rank G4

Modeled by W. Williams

MTNHP status special concern



State range

Breeding restricted to east-northeast Montana; recorded in low numbers along the lower Yellowstone River and the Missouri River below Fort Peck dam and on small islands in the Fort Peck Reservoir (USFWS 1990, Montana Bird Distribution Committee 1996). Transient sightings rare in Montana; overwintering does not occur (Montana Bird Distribution Committee 1996).

Habitat description

Breed on open, flat areas of sandy coastal beaches, and inland on sand/gravel bars, shorelines and islands of rivers and reservoirs (Ehrlich et al. 1988, DeGraaf et al. 1991). Interior populations nest on sparsely vegetated to barren areas of sand or gravel along the Mississippi and Missouri River systems (Ziewitz et al. 1992, Kirsch 1996); also the salt plains of Oklahoma (USFWS 1990). In the upper Missouri River Basin (Great Plains area), often nest in association with Piping Plovers (Ziewitz et al. 1992); in the Arkansas River system, American Avocets and Snowy Plovers are breeding associates (Grover and Knopf 1982, USFWS 1990). Throughout the interior breeding range, preferred nesting habitat is similar: Sparsely vegetated riverine sand and gravel bars within open, wide channel areas, or salt flats of lake shorelines (USFWS 1990, Ziewitz et al. 1992, Kirsch 1996). Due to high river flow at the onset of nesting, nest sites are usually higher on the sandbar, away from the waters edge (USFWS 1990, Kirsch 1996). Along the Platte River, NE, ideal nesting habitat is 1200m stretches of river with the channel at least 275m wide, with sandbars or islands 1.45-4.0ha in size and at least 0.5m tall, but ≥ 0.9 m in height offers better protection from inundation (Ziewitz et al. 1992). Also nest on artificial substrates - dikefields, sand and gravel pits, and dredgespoil islands (Reel et al. 1989, USFWS 1990). Nest solitarily or in colonies; nests are shallow depressions in the substrate, often near conspicuous objects such as stones or wood debris (USFWS 1990, DeGraaf et al. 1991). Breeding home range is usually limited to the river reach of/near the nest site sandbar (USFWS 1990). Nest site location is often adjacent to shallow areas of lakes or river backwaters that offer an abundant food source (Reel et al. 1989). Primarily piscivorous; also forage for crustaceans and insects (Reel et al. 1989,

USFWS 1990). Winter habitat includes coastal beaches, bays, and estuaries, as well as inland rivers and lakes (A.O.U. 1983). Interior populations have been estimated to be 3,360 pairs, and are declining due to alteration of river systems and destruction of nesting habitat (e.g., channelization, irrigation, reservoir construction), and lower productivity caused by human disturbance/recreation (Reel et al. 1989, USFWS 1990, Kirsch 1996). Although the silty, fine sand of the Yellowstone River shorelines represents marginal nesting habitat, Least Terns do nest there due to the absence of developments and human disturbance (Kreil and Dryer 1978). Along dammed rivers, irregular fluctuations in water levels flood existing nests, or promote vegetative growth on naturally scoured sandbars rendering them unsuitable for nesting (Reel et al. 1989, USFWS 1990). The Interior Least Tern is considered an endangered species by the USFWS and USFS Region 1 (Reel et al. 1989, Montana Bird Distribution Committee 1996). Conservation efforts include protection of existing breeding habitat by fencing off nesting areas to restrict human recreation (200m buffer distance around nests), developing reservoir control policies, and creation of suitable habitat by dredging sand/gravel to elevate existing sandbars and establish new ones in suitable river reaches (USFWS 1990, Ziewitz et al. 1992, Kirsch 1996).


Model assumptions & caveats

Breeding habitat modeled. Migration habitat not specifically modeled; assume potential inland migration habitat is defined within breeding habitat parameters. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Island or shoreline/river bar substrate characteristics cannot be resolved at mapping scale used; presence of river bars and islands may not be accurately represented in the land cover and hydrography, respectively. Potential for human disturbance information not available to include in model. Suitable breeding habitat likely overestimated.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- DeGraaf, R. M., V. E. Scott, R. H. Harme, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Faanes, C. A. 1983. Aspects of the nesting ecology of least terns and piping plovers in central Nebraska. *Prairie Nat.* 15(4): 145-154.
- Grover, P. B. and F. L. Knopf. 1982. Habitat requirements and breeding success of charadriiform birds nesting at Salt Plains National Wildlife Refuge, Oklahoma. *J. Field Ornithol.* 53:139-148.
- Kirsch, E. M. 1996. Habitat selection and productivity of least terns on the lower Platte River, Nebraska. *Wildl. Monogr.* 132. 48 pp.
- Kreil, R.L. and M.P. Dryer. 1987. Nesting of the Interior Least Tern on the Yellowstone River in North Dakota. *Prairie Nat.* 19(2): 135-136.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Reel, S., L. Schassberger and W. Ruediger. 1989. Caring for Our Natural Community: Region 1 - Threatened, Endangered and Sensitive Species Program. USDA, Forest Service Northern Region Wildlife and Fisheries.
- U.S. Fish and Wildlife Service. 1990. Recovery plan for the interior population of the Least Tern (*Sterna antillarum*). Dept. of the Interior, U.S. Fish and Wildl. Serv., Twin Cities, MN. 90 pp.

Ziewitz, J.W., J.G. Sidle and J.J. Dinan. 1992. Habitat consevation for nesting Least Terns and Piping Plovers on the Platte River, Nebraska. *Prairie Nat.*, 24(1): 1-20.



BLACK TERN

Chlidonias niger


State rank S3B,SZN

Element code ABNNM10020

Global rank G4

Modeled by W. Williams

MTNHP status special concern



State range

A fairly common summer resident in the Rocky Mountain region, breed locally (more prevalent in the western Montana) and migrate widely throughout the state; overwintering does not occur (Montana Bird Distribution Committee 1996).

Habitat description

Breed in shallow freshwater marshes with extensive stands of emergent vegetation and areas of open water, in forested or open water habitats up to 1540m elevation - including prairie sloughs, lake or pond margins, shallow river impoundments; occasionally river or island edges, wet meadows, swampy grasslands; sometimes cultivated rice fields, large stock ponds (Stewart 1975, Johnsgard 1986, DeGraaf et al. 1991, Dunn and Agro 1995). In the Rocky Mountain region, breed mainly in marshland or shallow grassy sloughs on the prairies, sedge dominated ponds in the parklands, and marshy cattail margins of northern lakes (Salt and Salt 1976, Johnsgard 1986). In the Prairie Pothole region of ND, highest numbers and densities breed on semipermanent wetlands (Stewart and Kantrud 1984). In IA, marshes or marsh complexes of 11-20+ha are preferred for nesting (Brown and Dinsmore 1986). Nests consist of low, loose, damp vegetation debris built on floating mats of dead marsh vegetation or boards; occasionally on nonfloating substrates such as muskrat houses or abandoned grebe nests - favored sites are surrounded by emergent vegetation to reduce wind and wave perturbations (Bergman et al. 1970, Ehrlich et al. 1988, Dunn and Agro 1995). Nest colonially or singly; frequent renesting and low site tenacity are adaptations to highly variable reproductive success (DeGraaf et al. 1991, Dunn and Agro 1995). Primarily insectivorous (both aquatic and terrestrial species consumed); also forage for worms, mollusks, crustaceans, small fishes (DeGraaf et al. 1991, Dunn and Agro 1995). During migration, frequent freshwater lakes, reservoirs, rivers, and other interior wetlands; also cultivated fields (DeGraaf et al. 1991, Dunn and Agro 1995). Winter habitat is primarily marine - usually within 30km of the coast; also coastal wetlands, estuaries (DeGraaf et al. 1991, Dunn and Agro 1995). The Black Tern is listed as a federal, category 2 candidate species and is considered a threatened, endangered or


sensitive species in Montana (Dunn and Agro 1995, Montana Bird Distribution Committee 1996). Populations have declined in many areas due to wetland conversion/degradation in both breeding and migration stopover areas (Ehrlich et al. 1988, Dunn and Agro 1995). Agricultural chemicals and pesticides have also contributed to declines by reducing hatching success and the insect food base in some areas (Ehrlich et al. 1988, Dunn and Agro 1995). Will readily nest in artificial (e.g., settling ponds) or restored wetlands, and will tolerate human activity near the nesting marsh as long as colonies are not entered (Delehanty and Svedarsky 1993, Dunn and Agro 1995). River impoundments and wetlands managed for waterfowl are suitable for nesting if water levels are held stable throughout the nesting season and emergent vegetation is maintained (Dunn and Agro 1995).

Model assumptions & caveats

Breeding habitat modeled. Migration habitat not specifically modeled; potential inland migration habitat is assumed defined within breeding habitat parameters - agricultural areas not included, as would diminish relevance of model as regards breeding habitat requirements and greatly overestimate habitat. Hydrography coverage may not be consistently accurate and/or true to mapping scale used. Persistence of intermittent water (e.g., seasonal vs. semipermanent, etc.) and water depth information not available to include in model.

References

- Bergman, R. D., P. Swain, and M. W. Weller. 1970. A comparative study of nesting Forster's and black terns. *Wilson Bull.* 82:435-444.
- Brown, M. and J. J. Dinsmore. 1986. Implications of marsh size and isolation for marsh bird management. *J. Wildl. Manage.* 50:392-397.
- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.
- Delehanty, D.J. and W.D. Svedarsky. 1993. Black Tern colonization of a restored prairie wetland in northwestern Minnesota. *Prairie Nat.* 25:213-218.
- Dunn, E. H. and D. J. Agro. 1995. Black Tern (*Chlidonias niger*). In: A. Poole and F. Gill, (eds.), *The Birds of North America*, No. 147. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union. 24 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds*. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.
- Johnsgard, P. A. 1986. *Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region*. Colorado Associated University Press, Boulder. xi + 504 pp.
- Kantrud, H.A. and R.E. Stewart. 1984. Ecological distribution and crude density of breeding birds on prairie wetlands. *J. Wildl. Manage.* 48(2): 426-437.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Salt, W.R. and J.R. Salt. 1976. *The birds of Alberta*. Hurtig Publishers, Edmonton, Alberta. xv + 498 pp.
- Stewart, R.E. 1975. *Breeding birds of North Dakota*. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



YELLOW-BILLED CUCKOO

Coccyzus americanus


State rank S3B,SZN

Element code ABNRB02020

Global rank G5

Modeled by C. Tobalske

MTNHP status special concern



State range

Breeds in southeastern Montana (T. McEneaney pers. comm.). Although records show no breeding since 1991, there are likely more occurrences in eastern and southeastern Montana than the database reflects, due simply to a shortage of submitted observations (Montana Bird Distribution Committee 1996, T. McEneaney pers. comm.).

Habitat description

Preferred habitat includes moderately dense thickets near watercourses, second-growth shrublands, deserted farmlands overgrown with shrubs and brush, and brushy orchards (Johnsgard 1979). Also nests in extensive, mature riparian woodlands, especially of cottonwood or willow (Dobkin 1992), and open woodlands where undergrowth is thick (AOU 1983). In Arizona at 4100', breeds in cottonwoods and willows (Hamilton and Hamilton 1965). In Montana, rare at high elevations (Johnsgard 1986).

Model assumptions & caveats

References

American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.

Benton, R. M. 1987. The Yellow-Billed Cuckoo. Utah Birds 3:7-11.

Dobkin, D. S. 1992. Neotropical migrant landbirds in the Northern Rockies and Great Plains. U.S.D.A. For. Serv. N. Region Publ. R1-93-34. Missoula, Mont.

Hamilton, W. J., and M. E. Hamilton. 1965. Breeding characteristics of yellow-billed cuckoos in Arizona. Proc. Calif. Acad. Sci. 32:405-432.


Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.

Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Potter, E. 1980. Notes on nesting yellow-billed cuckoos. J. Field Ornith. 51:17-29.

Preble, N. A. 1957. The nesting habits of the yellow-billed cuckoo. Amer. Midl. Nat. 57:474-82.



BARN OWL

Tyto alba


State rank SAB,SAN

Element code ABNSA01010

Global rank G5

Modeled by M. Hart

MTNHP status watch list



State range

An accidental breeder, sparsely and patchily distributed, with a few records spread widely throughout the state (Montana Bird Distribution Committee 1996). Montana is at the northern periphery of the species' breeding range, which is apparently limited by winter weather (Marti 1994). Currently, there are two confirmed breeding records, both near the state borders (one south-central, one southwestern), in Carbon and Ravalli counties (Wright 1996). The first was in 1989 (Holt et al. 1992). A nest also has been reported farther north in Manitoba (Nero 1995).

Habitat description

See review by Marti (1992): Uses a wide range of open habitats at low elevations, including urban and agricultural areas as well as grasslands, deserts, and marshes. Nests in cavities of both natural and anthropogenic sources: trees, cliffs, caves, riverbanks, church steeples, barn lofts, hay stacks, and nest boxes. The availability of nest cavities limits use of suitable foraging habitat. However, Barn Owls show great adaptability in selecting nest sites and foraging habitat, and are able to exist in many landscapes dominated by humans. Breeding and wintering ranges are generally the same for all but the most northerly populations (i.e, Montana), which are reported to be partly migratory. In northern winter often roosts in dense conifers; also roosts in nest boxes if available (Marti and Wagner 1985).


Nero (1995) speculates that an increase in numbers of derelict structures (e.g., houses, barns, sheds) in the northern Great Plains over the past several decades has provided a new source of shelter for Barn Owls. Such a factor could be critical in the species' ability to successfully inhabit Montana and other northern regions, because the Barn Owl is susceptible to starvation during prolonged low temperatures and snow cover (Marti and Wagner 1985). Near the northern end of the normal breeding range in northern Utah, winter weather has been found to cause great variations in reproductive performance (Marti 1994).

Model assumptions & caveats

Modeled, but because of the rarity and peripheral nature of breeding records in the state relative to the large size of the latilongs, overestimation of habitat/predicted distribution seems probable. Presence of human-created structures for nesting could not be evaluated, but is inferred in the selected cover types.

References

- Holt, D.W., K. Hicks, and W.D. Norton. 1992. First nest record for the Barn Owl in Montana. *Prairie Nat.* 24(2):121-122.
- Marti, C.D. 1992. Barn Owl (*Tyto alba*). In *The Birds of North America*, No. 1 (A. Poole, P. Stettenheim, and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. (This account replaces an earlier draft, with limited distribution, published in 1991.) 16 pp.
- Marti, C. D. 1994. Barn Owl reproduction: patterns and variation near the limit of the species' distribution. *Condor* 96:468-484.
- Marti, C. D. and P. W. Wagner. 1985. Winter mortality in Common Barn-Owls and its effect on population density and reproduction. *Condor* 87:111-115.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Nero, R.W. 1995. Barn Owl nesting in Manitoba. *Blue Jay* 53(3):159-166.
- Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.




FLAMMULATED OWL

Otus flammeolus

State rank S2S3B,SZN
Global rank G4
MTNHP status special concern

Element code ABNSB01020
Modeled by M. Hart



State range

Found in western Montana; breeding has been confirmed in six latilongs, generally in the southwest (Montana Bird Distribution Committee 1996). Because of its widespread presence in Missoula and Ravalli counties, should not be considered rare in the state (P.L. Wright 1996).

Habitat description

See review by McCallum (1994): Flammulated Owls occupy primarily open ponderosa pine/Douglas-fir or similar forests, e.g., dry montane conifer or aspen forests. The species is associated with mature to old-growth ponderosa pine and Douglas-fir forests throughout the northern and central Rocky Mountains (Bull and Anderson 1978, Goggans 1986, Holt and Hillis 1987, Howie and Ritcey 1987, Reynolds and Linkhart 1987, Atkinson and Atkinson 1990, Bull et al. 1990, Reynolds and Linkhart 1992, V. Wright 1996). Stands used by Flammulated Owls also tend to be relatively open (Goggans 1986, Howie and Ritcey 1987, McCallum and Gehlbach 1988, Atkinson and Atkinson 1990).

In northeastern Oregon, stands of large-diameter (>50 cm DBH) ponderosa pine and Douglas-fir or grand fir with ponderosa pine in the overstory were identified as nesting habitat (Bull and Anderson 1978, Bull et al. 1990). Similarly, Goggans (1986) described nesting habitat as stands of ponderosa pine/Douglas-fir, 30-50 cm DBH, with less than 50% canopy closure. Owls foraged in the edge between forest and grassland, as well as in ponderosa pine/Douglas-fir forests of low or moderate density. Density appeared to be a crucial aspect of roosting habitat: owls roosted in mixed conifer stands, and avoided open stands of ponderosa pine.

Reynolds and Linkhart (1987, 1992) have found a strong association between Flammulated Owls and old-growth ponderosa pine/Douglas-fir habitat in Colorado, noting that such forests were used more than expected for nesting, foraging, and singing. They speculate that the presence of cavities and snags, the abundance of arthropods, and a stand structure suitable for foraging may be factors in this preference (Reynolds and Linkhart 1992). Owls have also been found to nest in live aspen ($n = 3$) in

Colorado (Richmond et al. 1980).

Marcot and Hill (1980) also recorded use of hardwoods; California black oak (*Quercus kelloggii*) was present at 67% of locations in northwestern California, while ponderosa pine was present at 50% of locations. All territory sites were on xeric midslopes or near ridgetops. Also in California, Bloom (1983) observed 3 owls, all in stands dominated by ponderosa pine.

In central Idaho, territorial owls occupied relatively open, multistoried Douglas-fir, ponderosa pine, and mixed conifer stands with some mature trees usually present (Atkinson and Atkinson 1990). Territories were often near more open areas, including old burns, grassy hillsides, natural clearings, or clearcuts. Atkinson and Atkinson (1990) also noted a clumped distribution of territorial males, leaving apparently suitable habitat vacant. Similarly, Marcot and Hill (1980) found "quasi-colonies" of territorial males, along with unoccupied areas of apparently optimal habitat.

At the northern edge of the owl's range in British Columbia, Howie and Ritcey (1987) identified mature/old-growth (>100 year-old) Douglas-fir and Douglas-fir/ponderosa pine stands as nesting habitat, finding that owl densities were highest in stands 140-200+ years old. Stands were open, with canopy closures between 35-65%, and at least 2 canopy layers were present. Although Howie and Ritcey (1987:253) found a clearer association with mature/old-growth Douglas-fir than with ponderosa pine, they stated that "...the open nature of the fir forests coupled with natural or artificial openings created by logging probably resembles the physical structure of preferred forests in the southern portion of the owl's range."

Most recently, V. Wright (1996) found that owls in the Bitterroot and Sapphire ranges of southwestern Montana used mature and old-growth ponderosa pine/Douglas-fir stands rather than young stands or stands of other conifers, but noted that 48% of the plots in the study area with suitable microhabitat were unoccupied. Examined at multiple scales, owls responded to the presence of snags and large trees (microhabitat); the presence of openings (home range); and the overall abundance of low/moderate canopy closure ponderosa pine/Douglas-fir (landscape). Occupied landscapes were described as a mosaic of grass/xeric shrub and forest edge habitat, which is characteristic of most of the species' range.

Model assumptions & caveats

Model does not include forest stand structure/size class, and thus overestimates potential habitat. Unable to evaluate presence of suitable nest sites (snags) or stand understory. Forest edge habitats were addressed by placing a 90 m buffer around selected cover types, and selecting openings (grass and shrub) when they fall within this narrow buffer. Model queries for nesting and foraging habitat.

References

- Atkinson, E. C. and M. L. Atkinson. 1990. Distribution and status of Flammulated Owls (*Otus flammeolus*) on the Salmon National Forest. Idaho Nat. Heritage Prog., Idaho Dept. Fish Game, Salmon Natl. Forest Unpublished report. 25 pp. plus appendices.
- Bloom, P.H. 1983. Notes on the distribution and biology of the Flammulated Owl in California. West. Birds 14:49-52.
- Bull, E. L., A. L. Wright, and M. G. Henjum. 1990. Nesting habitat of Flammulated Owls in Oregon. J. Raptor Res. 24:52-55.
- Bull, E. L. and R. G. Anderson. 1978. Notes on Flammulated Owls in northeastern Oregon. Murrelet 59:26-28.
- Goggans, R. 1986. Habitat use by Flammulated Owls in northeastern Oregon. M.S. thesis, Oregon State Univ., Corvallis. 54 pp.
- Hayward, G. D., and J. Verner, tech. eds. 1994. Flammulated, Boreal, and Great Gray owls in the United States: A technical conservation assessment. Gen. Tech. Rep. RM-253. U.S.D.A. Forest Service, Rocky Mountain Forest and

Range Experiment Station. Fort Collins, CO. 214 pp. 3 maps.

Holt, D. W. and J. M. Hillis. 1987. Current status and habitat associations of forest owls in western Montana. Pp. 281-288 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings*, Feb. 3-7, Winnepeg, Manitoba. Gen. Tech. Rep. RM-142. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Holt, D.W., J.A. Hoy, and P.L. Wright. 1987. Occurrence and first nest record of Flammulated Owls in Montana. *J. Raptor Res.* 21(3):120-121.

Howie, R.R., and R. Ritcey. 1987. Distribution, habitat selection, and densities of Flammulated Owls in British Columbia. Pp. 249-254 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings*. Gen. Tech. Rep. RM-143. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo. 309 pp.

Marcot, B.G. and R. Hill. 1980. Flammulated Owls in northwestern California. *West. Birds* 11:141-149.

McCallum, D.A. 1994. Flammulated Owl (*Otus flammeolus*). In *The Birds of North America*, No. 93 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 pp.

McCallum, D. A. and F. R. Gehlbach. 1988. Nest-site preferences of Flammulated Owls in western New Mexico. *Condor* 90:653-661.


Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Reynolds, R.T. and B.D. Linkhart. 1987. Fidelity to territory and mate in Flammulated Owls. Pp. 234-238 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings*. Gen. Tech. Rep. RM-143. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, Colo. 309 pp.

Richmond, M.L., L.R. DeWeese, and R.E. Pillmore. 1980. Brief observations on the breeding biology of the Flammulated Owl in Colorado. *West. Birds* 11:35-46.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.

Wright, V. 1996. Multi-scale analysis of Flammulated Owl habitat use: owl distribution, habitat management, and conservation. M.S. thesis, Univ. Montana, Missoula. 91 pp.



EASTERN SCREECH-OWL

Orus asio


State rank S3S4

Element code ABNSB01030

Global rank G5

Modeled by M. Hart

MTNHP status watch list



Predicted habitat: 514,275 ha, 1.35 % of state.

State range

Riparian corridors in eastern Montana extending west as far as the main Rocky Mountain chain; most locations in southeastern Montana (Montana Bird Distribution Committee 1996). Exact range limits are not well known (Johnsgard 1986, Adam 1987). Probably relatively common across much of eastern and central Montana (Wright 1996). May be common in preferred habitat along the Milk River (D. Prellwitz, pers. comm.). Specimen collected at Mammoth, Yellowstone National Park, Wyoming (T. McEaney, pers. comm.) may offer insight on the species' western limits in southern Montana.

Habitat description

Open woodlands, deciduous forest, parklands, residential areas in towns, scrub, and riparian woodland in drier regions (AOU 1983). In Montana, probably is largely restricted to riparian woodlands along major streams and rivers (J. Marks, pers. comm, T. McEaney, pers. comm.). See review by Gehlbach (1995): The Eastern Screech-Owl occupies the broadest ecological niche of any owl in its range; although common, it is poorly studied. It is found in tree-dominated landscapes of most types, natural and anthropogenic, early to late successional, river valleys to mountain slopes. Generally found <1800 m. Marginal in boreal and montane conifer forests. Requires only a suitable cavity tree or substitute (e.g., nest box). Forages in forest and woodland understory, openings, and edges. Most habitat descriptions are based on studies from the eastern U.S. Habitats listed by Gehlbach (1995) that would apply to Montana include northern mixed and deciduous forest, urban and suburban yards, parks, and green belts.

Two attempts have been made to better define the range overlap between *O. asio* and *O. kennicottii* in Wyoming (Fitton 1993, Dorn and Dorn 1994). Fitton (1993) reports Eastern Screech-Owls at four locations, all east of the Continental Divide and in riparian woodlands dominated by Plains cottonwood. The highest location was at 1920 m. Dorn and Dorn (1994) report 21 locations; again, all are east of the divide. All were also <1890 m. Large-sized Plains cottonwoods were noted to be an important habitat component, followed by willow and boxelder, all of which tend to grow along


streams on the plains. Similarly, in southeastern Saskatchewan, Adam (1987) describes the Eastern Screech-Owl as a rare resident of riparian habitats, typically Manitoba maple river bottom woodlands. Of 17 singles or pairs observed by Adam (1987), most were associated with farms, or were near villages or houses.

Model assumptions & caveats

Model does not assess presence of suitable cavities for nesting.

References

- Adam, C. I. G. 1987. Status of the Eastern Screech Owl in Saskatchewan with reference to adjacent areas. Pp. 268-276 in R.W Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings, Feb. 3-7, Winnipeg, Manitoba*. Gen. Tech. Rep. RM-142. USDA, Forest Service., Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.
- American Ornithologists' Union. 1983. *Check-list of North American birds*, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Bull, E. L., A. L. Wright, and M. G. Henjum. 1989. Nesting and diet of Long-eared Owls in conifer forests, Oregon. *Condor* 91:908-912.
- Dorn, R.D. and J.L. Dorn. 1994. Further data on screech-owl distribution and habitat use in Wyoming. *West. Birds* 25:35-42.
- Fitton, S. 1993. Screech-owl distribution in Wyoming. *West. Birds* 24:182-188.
- Gehlbach, F.R. 1995. Eastern Screech-owl (*Otus asio*). In *The Birds of North America*, No. 165 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 pp.
- Holt, D.W. 1997. The Long-eared Owl (*Asio otus*) and forest management: a review of the literature. *J. Raptor Res.* 31(2):175-186.
- Holt, D. W. and J. M. Hillis. 1987. Current status and habitat associations of forest owls in western Montana. Pp. 281-288 in R.W Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings, Feb. 3-7, Winnipeg, Manitoba*. Gen. Tech. Rep. RM-142. USDA, Forest Service., Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.
- Johnsgard, P. A. 1986. *Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region*. Colorado Associated University Press, Boulder. xi + 504 pp.
- Maples, M.T. and D.W. Holt. 1995. Ground-nesting Long-eared Owls. *Wilson Bull.* 107(3):563-565.
- Marks, J. S. 1984. Feeding ecology of breeding Long-eared Owls in southwestern Idaho. *Can. J. Zool.* 62: 1528-1533.
- Marks, J.S., D.L. Evans, and D.W. Holt. 1994. Long-eared Owl (*Asio otus*). In *The Birds of North America*, No. 133 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's *Montana Bird Distribution, Fifth Edition*. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



WESTERN SCREECH-OWL

Otus kennicottii


State rank S3S4

Element code ABNSB01040

Global rank G5

Modeled by M. Hart

MTNHP status watch list



Predicted habitat: 317,674 ha, 0.83% of state.

State range

Western Montana, as far east as the eastern edge of the Rocky Mountain chain; most locations in west-central Montana (Montana Bird Distribution Committee 1996). Easternmost records are from Choteau (north) and Bozeman (south). Exact range limits are not well known (Johnsgard 1986, Holt and Hillis 1987).

Habitat description

Throughout range, woodlands, especially oak and riparian woodlands, and scrub (AOU 1983); in Montana, most often found in riparian areas with gallery forests (T. McEaney, pers. comm.). See Holt and Hillis (1987), who reviewed status and habitat use of forest owls in western Montana. For the Western Screech-Owl, they could only find three nest sites, located in black cottonwood and quaking aspen, but noted that several observers have reported fledglings in cottonwood habitats near Missoula. Holt and Hillis (1987:282) further suggest "This is undoubtedly a more common species closely associated with riparian habitats." They also note that the owls are occasionally seen away from riparian areas, in mixed conifer forests.

Two attempts have been made to better define the range overlap between *O. asio* and *O. kennicottii* in Wyoming (Fitton 1993, Dorn and Dorn 1994). Dorn and Dorn (1994) located no Western Screech-Owls. Fitton (1993) found them at two locations in the Wind River Mountains, elevation 2400 m. In both areas, aspen was most abundant, followed by Douglas-fir.

Model assumptions & caveats

Model does not assess presence of suitable nest sites.

References

American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.


Dorn, R.D. and J.L. Dorn. 1994. Further data on screech-owl distribution and habitat use in Wyoming. *West. Birds* 25:35-42.

Fitton, S. 1993. Screech-owl distribution in Wyoming. *West. Birds* 24:182-188.

Hand, R.L. 1969. A distributional checklist of the birds of western Montana. Unpubl. rep. 55 pp.

Holt, D. W. and J. M. Hillis. 1987. Current status and habitat associations of forest owls in western Montana. Pp. 281-288 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings, Feb. 3-7, Winnepeg, Manitoba*. Gen. Tech. Rep. RM-142. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.



NORTHERN HAWK OWL

Surnia ulula


State rank SAB,SAN

Element code ABNSB07010

Global rank G5

Modeled by M. Hart

MTNHP status watch list



Predicted habitat: 503,817 ha, 1.32 % of state.

State range

Occurs primarily in northwestern Montana, although scattered observations have been recorded throughout other parts of the state (Montana Bird Distribution Committee 1996). The single confirmed breeding took place near Polebridge in Glacier National Park in 1994. Nevertheless, the species should no longer be considered rare in the state (Wright 1996).

Habitat description

Widely distributed across the northern boreal forest from Alaska to Newfoundland, yet among North America's least studied birds (Rohner et al. 1995). Found in open mixed or coniferous forest, forest edge and clearings, old burns in deciduous forest, dense shrubby areas (especially tamarack), swamps, scrubby second-growth woodland and muskeg (AOU 1983). Clark and others (1987) describe Northern Hawk Owl habitat as clearings and patchy areas in northern coniferous forest, and low scrub and trees near water.

Model assumptions & caveats

Model based on very limited information. As a result, cover types included may be overly generous.

References


American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.

Clark, R.J., D.G. Smith, and L. Kelso. 1987. Distributional status and literature of northern forest owls. Pp. 47-55 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings, Feb. 3-7, Winnipeg, Manitoba*. Gen. Tech. Rep. RM-142. USDA, Forest Service., Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Rohner, C., J.N.M. Smith, J. Stroman, M. Joyce, F.I. Doyle, and R. Boonstra. 1995. Northern Hawk-owls in the nearctic boreal forest: prey selection and population consequences of multiple prey cycles. *Condor* 97:208-220.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



BURROWING OWL

Athene cunicularia


State rank S3B,SZN

Element code ABNSB10010

Global rank G4

Modeled by M. Hart

MTNHP status special concern



Predicted habitat: 23,445,126 ha, 61.57 % of state.

State range

Breeds widely east of the Rockies, but also occurs regularly in southwestern Montana, and scattered records exist elsewhere in intermountain valleys west of the Divide (Montana Bird Distribution Committee 1996). In Montana, may be declining even more precipitously than mountain plovers (R. Matchett, pers. comm.).

Habitat description

Open grasslands, especially prairie, plains, and savanna, sometimes in open areas such as vacant lots near human habitation or airports, typically nesting and roosting in burrows dug by mammals (AOU 1983). See Haug and others (1993) for review of this ground-dwelling inhabitant of the western deserts and grasslands, which nests in loose colonies in a wide variety of open environments, including agricultural lands. Presence of a nest burrow seems to be this owl's critical requirement; in fact, this species' close association with burrowing mammals suggests dependence upon them. Typically forages in shortgrass, mowed, or overgrazed pastures, but also uses golf courses, airports, and road rights-of-way. In Colorado, Burrowing Owls tend to select burrows that are close to other burrows and to roads, and that are surrounded by bare ground or short grass (Plumpton 1992).

Model assumptions & caveats

Presence of burrowing mammals assumed, not assessed. Model would be improved by inclusion of prairie dog town locations. An elevation limit was used primarily to filter out the mixed barren cover type (7800), which has a broad range, at higher elevations. Distribution is very likely overpredicted, based on recent declines.

References

Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.

American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.

Green, G. A. and R. G. Anthony. 1989. Nesting success and habitat relationships of Burrowing Owls in the Columbia Basin, Oregon. *Condor* 91:347-354.

Haug, E. A. and L. W. Oliphant. 1990. Movements, activity patterns, and habitat use of Burrowing Owls in Saskatchewan. *J. Wildl. Manage.* 54:27-35.


Haug, E. A., B. A. Milsap, and M. S. Martell. 1993. Burrowing Owl (*Speotyto cunicularia*). In *The Birds of North America*, No. 61 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 20 pp.

Kantrud, H. A. 1981. Grazing intensity effects on the breeding avifauna of North Dakota native grasslands. *Can. Field-Nat.* 95:404-417.

Kantrud, H.A. and R.L. Kologiski. 1982. Effects of soils and grazing on breeding birds of uncultivated upland grasslands of the northern Great Plains. U.S.D.I., Fish and Wildl. Serv., Wildl. Res. Rep. 15. 33 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Plumpton, D.L. 1992. Aspects of nest site selection and habitat use by Burrowing Owls at the Rocky Mountain Arsenal, Colorado. M.S. thesis, Texas Tech. Univ., Lubbock.



GREAT GRAY OWL

Strix nebulosa


State rank S3

Element code ABNSB12040

Global rank G5

Modeled by M. Hart

MTNHP status special concern



State range

A year-round resident of the montane areas of western and central Montana; winter records are less numerous (Montana Bird Distribution Committee 1996).

Habitat description

See review by Bull and Duncan (1993): Primarily a bird of dense, northern boreal forests, the Great Gray Owl finds suitable coniferous habitat southward into the Northern Rocky and Sierra mountains. Favors areas near bogs, forest edge, montane meadows and other openings. Breeding distribution appears to be limited by availability of prey and of nest sites. In the southern parts of the species' range, it is found in deciduous or coniferous forests up to 2800 m. In Idaho and Wyoming, nearly all observations have been made in the lodgepole pine/Douglas-fir/aspen zone (Franklin 1988). In Montana, the species most often is associated with spruce-fir, lodgepole pine, and Douglas-fir forests, as well as mixed conifer and whitebark pine forests farther east on the Lewis and Clark National Forest; elevations are described as "high", ~7000 ft., and ~7200 ft. for various National Forests (Verner 1994). Wintering habitat is the same as breeding habitat (Duncan and Hayward 1994).

Model assumptions & caveats

Presence of suitable nest sites not evaluated. Interspersion of forest/openings not evaluated. Elevation limit set low so that the selected cover types themselves define the elevation range for the most part; used elevation to filter included cover types like 4280 (mixed mesic forest) which occur at lower elevations. Focus placed on habitats used in the breeding season, assuming that those will include the best habitats for the rest of the year as well. Habitat is likely overpredicted.

References

Bull, E.L. and J.R. Duncan. 1993. Great Gray Owl (*Strix nebulosa*). In *The Birds of North America*, No. 41 (A. Poole

and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 16 pp.

Duncan, J.R. 1997. Great Gray Owls (*Strix nebulosa nebulosa*) and forest management in North America: a review and recommendations. *J. Raptor Res.* 31(2):160-166.

Duncan, J.R. and P.H. Hayward. 1994. Review of technical knowledge: Great Gray Owls. Pp. 159-175 in G.D. Hayward and J. Verner, tech. eds. *Flammulated, Boreal, and Great Gray Owls in the United States: A technical conservation assessment*. Gen. Tech. Rep. RM-253. U.S.D.A. Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO. 214 pp. 3 maps.


Franklin, A. B. 1988. Breeding biology of the Great Gray Owl in southeastern Idaho and northwestern Wyoming. *Condor* 90:689-696.

Hayward, G. D., and J. Verner, tech. eds. 1994. *Flammulated, Boreal, and Great Gray owls in the United States: A technical conservation assessment*. Gen. Tech. Rep. RM-253. U.S.D.A. Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO. 214 pp. 3 maps.

Holt, D. W. and J. M. Hillis. 1987. Current status and habitat associations of forest owls in western Montana. Pp. 281-288 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings, Feb. 3-7, Winnipeg, Manitoba*. Gen. Tech. Rep. RM-142. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's *Montana Bird Distribution, Fifth Edition*. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Verner, J. 1994. Current management situation: Great Gray Owls. Pp. 155-158 in G.D. Hayward and J. Verner, tech. eds. *Flammulated, Boreal, and Great Gray Owls in the United States: A technical conservation assessment*. Gen. Tech. Rep. RM-253. U.S.D.A. Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO. 214 pp. 3 maps.



BOREAL OWL

Aegolius funereus


State rank S3S4

Element code ABNSB15010

Global rank G5

Modeled by M. Hart

MTNHP status special concern



State range

Mountainous regions of western and central Montana (Montana Bird Distribution Committee 1996). Should no longer be considered rare in the state (Wright 1996).

Habitat description

See review by Hayward and Hayward (1993): Found in boreal and subalpine forests across North America, with scattered populations extending southward into the subalpine forests of the Rocky, Blue, and Cascade Mountains. Breeding and wintering habitats are similar.

Boreal Owls are typically found in mature/old-growth spruce-fir forests in the northern and central Rocky Mountains (Palmer 1986, Hayward et al. 1987, Holt and Hillis 1987, O'Connell 1987, Ryder et al. 1987, Holt and Ermatinger 1989, Hayward et al. 1993). Although Boreal Owls may be relatively common in certain habitats, until recently they have remained little known in the Rocky Mountain states, probably due to their breeding chronology and high elevation associations (Holt and Hillis 1987).

Based on limited surveys in Montana, Holt and Hillis (1987) noted a preference for mature/overmature Engelmann spruce/subalpine forests above 1500 m. Holt and Ermatinger (1989) recorded the first confirmed nest in Montana, also in an Engelmann spruce/subalpine fir stand over 120 years old.

An extensive description of habitat use by Boreal Owls in the northern Rocky Mountains is provided by Hayward et al. (1993). Forests in Montana, Idaho, and northern Wyoming were surveyed for Boreal Owls, and 49 nests or singing males were found. No owls were detected below 1292 m, and 75% of locations were above 1584 m. Forest cover types in which owls were located included lodgepole pine, Douglas-fir, western hemlock/western larch/subalpine fir, Engelmann spruce, and Engelmann spruce/subalpine fir. Stands were classified as mature or older at 76% of locations.

Hayward et al. (1993) also studied Boreal Owls more intensively in the River of No Return Wilderness (RNRW) in central Idaho. Of 28 breeding sites in RNRW, 39% were in mixed conifer, 25% in Engelmann spruce/subalpine fir, 18% in Douglas-fir, and 18% in aspen stands. Lodgepole pine was not used for nesting, although it was the most common forest type in the study area. Nest sites were found only in structurally complex mature and old forests; characteristic features included high density of large trees, open understory, and multilayered canopy. The mean size of nest stands was 7.6 ha (range 0.8-14.6 ha). Average roost-to-nest distance was approximately 1730 m; Hayward et al. (1993) suggested that this measure is probably a good approximation of foraging distance. The best foraging habitat was associated with Engelmann spruce/subalpine fir stands, where prey densities were highest and open stand structure facilitated hunting. Mature/overmature Engelmann spruce/subalpine fir stands also provided cool sites for summer roosting, and Hayward et al. (1993) proposed that roosting habitat can be maintained through management of foraging habitat. Finally, Hayward et al. (1993) further recommended that all forested sites within the spruce-fir zone be considered as potential Boreal Owl habitat, as well as forests 100-200 m below this zone, which may provide the most important nesting habitats.

Model assumptions & caveats

Assume that broadleaf forest (4140) is most likely to be aspen in the elevation zones in which Boreal Owls are found. Also assume that foraging/roosting habitat will be adequately captured by queries for nesting habitat; similarly, that wintering habitat will be adequately captured by breeding habitat. Model does not include forest stand structure/size class, nor does it assess presence of suitable nest sites. Model probably overestimates habitat because large cavities are likely limited at highest elevations (N. Warren, pers. comm.).

References

Burcham, M. 1992. All-owls transects: final report, Flathead National Forest, March-April, 1992. Unpubl. report on file, Flathead National Forest, Kalispell, Mont.

Hayward, G.D. 1997. Forest management and conservation of Boreal Owls in North America. *J. Raptor Res.* 31(2):114-124.

Hayward, G. D., and J. Verner, tech. eds. 1994. Flammulated, Boreal, and Great Gray owls in the United States: A technical conservation assessment. Gen. Tech. Rep. RM-253. U.S.D.A. Forest Service, Rocky Mountain Forest and Range Experiment Station. Fort Collins, CO. 214 pp. 3 maps.

Hayward, G.D. and P.H. Hayward. 1993. Boreal Owl (*Aegolius funereus*). In *The Birds of North America*, No. 63 (A. Poole and F. Gill, Eds.) Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologist's Union. 20 pp.

Hayward, G.D., P.H. Hayward and E.O. Garton. 1987. Revised breeding distribution of the Boreal Owl in the northern Rocky Mountains. *Condor* 89:431-432.

Hayward, G.D., P.H. Hayward, and E.O. Garton. 1993. Ecology of Boreal Owls in the northern Rocky Mountains, USA. *Wildl. Monogr.* 124. 59 pp.

Holt, D. W. and D. Ermatinger. 1989. First confirmed nest site of Boreal Owls in Montana. *Northwest. Nat.* 70:27-31.

Holt, D. W. and J. M. Hillis. 1987. Current status and habitat associations of forest owls in western Montana. Pp. 281-288 in R.W. Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings*, Feb. 3-7, Winnepeg, Manitoba. Gen. Tech. Rep. RM-142. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

O'Connell, M. W. 1987. Occurrence of the Boreal Owl in northeastern Washington. Pp. 185-188 in R.W. Nero, R.J.


Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings*, Feb. 3-7, Winnipeg, Manitoba. Gen. Tech. Rep. RM-142. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Palmer, D.A. 1986. *Habitat selection, movements and activity of Boreal and Saw-whet Owls*. M.S. thesis, Colo. State Univ., Fort Collins. 101 pp.

Palmer, D. A. 1987. Annual, seasonal, and nightly variation in calling activity of Boreal and Northern Saw-whet Owls. Pp. 162-168 in R.W Nero, R.J. Clark, R.J. Knapton, and R.H. Hamre (eds.), *Biology and conservation of northern forest owls: symposium proceedings*, Feb. 3-7, Winnipeg, Manitoba. Gen. Tech. Rep. RM-142. USDA, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO. 309 pp.

Palmer, D.A., and R.A. Ryder. 1984. The first documented breeding of the Boreal Owl in Colorado. *Condor* 86:215-217.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



BLACK SWIFT

Cypseloides niger


State rank S3B,SZN

Element code ABNUA01010

Global rank G4

Modeled by W. Williams

MTNHP status special concern



State range

Considered rare to uncommon in the U.S.; somewhat rare in Montana, breeding records limited to the northwest; overwintering does not occur (Montana Bird Distribution Committee 1996).

Habitat description


Breed in a variety of open, montane habitats, frequently seen over more rugged terrain, from sea level to 11,000ft (DeGraaf et al. 1991, Chantler and Driessens 1995). Habitat defined by mountainous areas associated with steep cliffs or narrow canyons; always near water (Johnsgard 1986, Dobkin 1994). Nest in small colonies, from 1-15 pairs, in high cliff crevices or ledges, steep rock canyon walls, shallow caves, sea cliffs, occasionally sea caves; preferably near or behind a waterfall (DeGraaf et al. 1991, Dobkin 1994, Chantler and Driessens 1995). Nests are pads of moss, ferns, pine needles bound with little mud (Ehrlich et al. 1988, Chantler and Driessens 1995). High relief, inaccessibility, unobstructed flyways, at least partial darkness, and water are nest site requirements (Knorr 1961, Hunter and Baldwin 1962). Black Swifts show strong nest site fidelity - even when severe drought greatly affects waterfall flow, they will return each year - but apparently do not nest on truly intermittent streams (Knorr 1993). In nw MT, a small colony was found nesting behind or immediately adjacent to a 150m wide, cascading waterfall, at 4,700ft elevation, amidst transition-zone vegetation: Lowland Fir (*Abies grandis*), Douglas-fir, Englemann Spruce, Mountain Maple, willow, alder, dogwood, and juniper (Hunter and Baldwin 1962). Aerially forage over both forested and open montane areas, exclusively for flying insects (Ehrlich et al. 1988). In nw Montana, often forage in valley bottoms, particularly over water; often many kilometers from nesting habitat, especially during stormy weather (D. Casey pers. comm.). Winter from Mexico, south to Costa Rica (Ehrlich et al. 1988, Dobkin 1994), and may be seen in large flocks during migration (Chantler and Driessens 1995). Black Swifts are one of the least known land birds in North America, and are considered a threatened, endangered, or sensitive species in Montana; BBS data, however, are not sufficient enough to determine population trends (Dobkin 1994, Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Migration habitat is assumed similar to that of breeding. Foraging habitat in addition to that defined for breeding, not well described, therefore not specifically included in model. Assume potential migration and foraging habitats are adequately defined within breeding habitat parameters, which include 1km buffer around potential nest areas. Waterfall information not available in hydrography coverage and cliffs could not be accurately resolved at mapping scale used. Attempted to target nest areas by intersecting stream coverage with slope $\geq 40\%$. Unable to model most nesting requirements, due to limitations of mapping scale and resolution; suitable breeding habitat may be overestimated, however, foraging ranges may extend beyond 1km buffer therefore foraging habitat is likely underestimated.

References

- Chantler, P. and G. Driessens. 1995. Swifts: A guide to the swifts and treeswifts of the world. Pica Press, Sussex, England. 237 pp.
- DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hunter, W. F. and P. H. Baldwin. 1962. Nesting of the black swift in Montana. *Wilson Bull.* 74(4):409-416.
- Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.
- Knorr, O. A. 1961. The geographical and ecological distribution of the black swift in Colorado. *Wilson Bull.* 73(2):155-170.
- Knorr, O. A. 1993. Breeding of the Black Swift in the Great Basin. *West. Birds* 24(3): 197-198.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.



BLACK-BACKED WOODPECKER

Picoides arcticus


State rank S3

Element code ABNYF07090

Global rank G5

Modeled by C. Tobalske, W. Williams

MTNHP status special concern



State range

Breed in montane areas of western Montana; winter range may be more restricted to northwestern Montana (Montana Bird Distribution Committee 1996).

Habitat description

A species associated with coniferous forests, often in openings where burns, logged areas, lake and stream shores, swamps, and bogs occur (Campbell et al. 1990). Prefers denser forests containing a mixture of coniferous species (Bock and Bock 1974). In Oregon, favors various mature and overmature mixed conifer types and pure lodgepole forests, and avoids seedling, sapling, pole stands and cut areas (Marshall 1992, Goggans et al. 1989). Nests also in Ponderosa Pine forests with moderate canopy closure (Bull et al. 1986). In Montana, preferred habitat consists of coniferous forests (fir, spruce), especially windfall and burned sites; the species nests and forages in recent burns (younger than 4 years), and also nests in dense forest stands (MNHP-VCA 1996). Management recommendations include the conservation of pine-dominated habitat blocks of about 1000 acres (MNHP-VCA 1996). In Lincoln County it was rarely found in Subalpine Fir and Lodgepole Pine woods, but occurred in Ponderosa Pine forests at low elevation, and in mixed conifer and Douglas-fir associations higher up (Weydemeyer and Weydemeyer 1928). Although an opportunistic species, it is much more abundant in burned forest than other "green" forests (S. Hejl pers. comm.), and may be suffering from the transition from extensive burned areas to second growth stands since the 1940s (McClelland 1977). The Black-backed Woodpecker is virtually restricted to early post-fire habitat in Montana; prevention of stand-replacement fires and post-fire salvage cutting have seriously reduced suitable breeding habitat (Hutto 1995b). Unburned forests may be sinks for birds emigrating from burns when those are 5-6 years old (Hutto 1995).


Model assumptions & caveats

Burn cover type temporally limited by acquisition date of satellite imagery. Burn cover type was

buffered by 5 km to focus on burned areas, yet include alternate habitat, albeit secondary. Suitable breeding habitat is likely over- and inaccurately estimated.

References

- Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.
- Bock, C. E. and J. H. Bock. 1974. On the geographical ecology and evolution of the three-toed woodpeckers, *Picoides tridactylus* and *P. arcticus*. *Amer. Midl. Nat.*, 92(2):397-405.
- Bull, E. L., S. R. Peterson and J. W. Thomas. 1986. Resource partitioning among woodpeckers in northeastern Oregon. USDA For. Serv. Pac. Northw. Res. Station, Portland, Ore., Res. Note PNW-444. 19 pp.
- Campbell, R.W., N.K. Dawe, I. McTaggart-Cowan, J.M. Cooper, G.W. Kaiser, M.C.E. McNall. 1990. The Birds of British Columbia. Vol. 2. Canadian Wildlife Service and the Royal British Columbia Museum and Environment Canada.
- Goggans, R., R. D. Dixon and L. C. Seminara. 1989. Habitat use by three-toed and black-backed woodpeckers, Deschutes National Forest, Oregon. Ore. Dept. Fish Wildl. - Nongame Wildl. Prog. USDA Deschutes Natl. For. Tech. Rep. 87-3-02. 43 pp.
- Harris, M. A. 1982. Habitat use among woodpeckers in forest burns. M.S. thesis. University of Montana, Missoula. 62 pp.
- Hutto, R.L. 1995a. Composition of bird communities following stand-replacement fires in Northern Rocky Mountain (U.S.A.) conifer forests. *Conservation Biology* 9: 1041-1058.
- Hutto, R.L. 1995b. USFS Northern Region Songbird Monitoring Program - Distribution and Habitat Relationships. Second Rept., USFS Region 1 contract #57-0343-5-00012. 120 pp.
- Marshall, D. B. 1992. Status of the black-backed woodpecker in Oregon and Washington. Audubon Society of Portland, OR. 15 pp.
- McClelland, B. R. 1977. Relationships between hole-nesting birds, forest snags and decay in western larch-Douglas fir forests of the northern Rocky Mountains. Ph.D. Thesis., Univ. of Montana, Missoula. 483 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Weydemeyer, W., and D. Weydemeyer. 1928. The woodpeckers of Lincoln County, Montana. *Condor* 30:339-346.



ALDER FLYCATCHER

Empidonax alnorum


State rank S1B,SZN

Element code ABPAE33030

Global rank G5

Modeled by W. Williams

MTNHP status special concern



Predicted habitat: 49,697 ha, 0.13 % of state.

State range

Uncommon summer resident in the northwestern Rocky Mountain region; breed primarily east and north of Montana (Johnsgard 1986, Wright 1996). Although there is no direct evidence of breeding in Montana to date, strong indirect evidence has been recorded, particularly in the Pine Butte Swamp area - nesting is also likely along the northern border of the state (Montana Bird Distribution Committee 1996, Wright 1996).

Habitat description

A common inhabitant of northern alder swamps and aspen parklands; breed in alder or willow thickets and deciduous forest edges bordering streams, lakes, ponds, bogs, swamps, muskegs; also brushy, scrubby growth in damp cut-over areas; occasionally low spruces in bogs (Godfrey 1966, A.O.U. 1983, Johnsgard 1986, DeGraaf et al. 1991). Usually found near water; require open forest areas and edges with low, dense shrubby growth (DeGraaf et al. 1991). Nests are built 2-6ft above ground, saddled in upright fork of low tree or shrub branch (Godfrey 1966, DeGraaf et al. 1991). Forage for flying insects, particularly beetles, by hawking/sallying from a perch; also glean spiders and berries from vegetation (Ehrlich et al. 1988, DeGraaf et al. 1991). In migration and winter also use open woodland habitats; winter in South America (A.O.U. 1983, Ehrlich et al. 1988). Closely related to Willow Flycatchers (*Empidonax traillii*); were formerly considered the same species (A.O.U. 1983, Ehrlich et al. 1988). The Alder Flycatcher is considered a threatened, endangered, or sensitive species in Montana - recently considered a separate species from the Willow Flycatcher, little is known about its breeding habitat requirements in the state (Montana Bird Distribution Committee 1996).

Model assumptions & caveats

Breeding habitat modeled. Migration habitat is somewhat similar to that of breeding and is assumed

defined within breeding habitat parameters. Difficult to model edge habitats at mapping scale used. Undergrowth/shrub cover density information not available to include in model.

References

American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.

DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.


Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.

Godfrey, W. Earl. 1966. The birds of Canada. National Museums of Canada. Ottawa. 428 pp.

Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.




CASSIN'S KINGBIRD

Tyrannus vociferans

State rank S1S3B,SZN
Global rank G5
MTNHP status special concern

Element code ABPAE52030
Modeled by W. Williams



State range

Local breeder in the high plains east of the Rocky Mountains; breeding in Montana is limited to the southeast; overwintering does not occur (Johnsgard 1986, Montana Bird Distribution Committee 1996).

Habitat description

Breed in open, semi-desert habitats, up to 7500ft (DeGraaf et al. 1991). In CA, use open valley woodlands and foothill grasslands with scattered oaks, cottonwoods, sycamores (DeGraaf et al. 1991). In the northern Rocky Mountains/ Great Plains region, breed in open country with scattered trees, especially pinyon-juniper woodlands; also dry savanna, Ponderosa Pine woodlands, open scrub, shrubsteppe, pine-oak associations; extending into grasslands and agricultural areas (Johnsgard 1986, DeGraaf et al. 1991, Dobkin 1994). Prefer habitats less open than do Western Kingbirds, such as pine-oak-juniper woodlands in TX and closed canopy riparian forests in AZ (Gamble and Bergin 1996). Nest near ends of horizontal branches in fairly tall deciduous trees, 8-55ft above ground; occasionally in bushes or on fence posts (Ehrlich et al. 1988, DeGraaf et al. 1991). Winter south into South America, in dry habitats such as highland pine-oak associations and dry scrub (A.O.U. 1983, Dobkin 1994). Forage primarily for insects, usually by sallying from perch; also take spiders, berries (DeGraaf et al. 1991, Dobkin 1994). Populations appear to be declining slightly in the West, but BBS data is insufficient (Dobkin 1994). The Cassin's Kingbird is considered a threatened, endangered, or sensitive species in Montana; eastern Montana is the extreme northern limit of the breeding range (Montana Bird Distribution Committee 1996, Dobkin 1994).

Model assumptions & caveats

Breeding habitat modeled. Cannot resolve interspersions of cover types at mapping scale used.

References

American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.

DeGraaf, R. M., V. E. Scott, R. H. Harne, L. Ernst, and S. H. Anderson. 1991. Forest and rangeland birds of the United States. U.S.D.A. Forest Service Agriculture Handbook 688. 625 pp.


Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.

Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.

Gamble, L.R. and T.M. Bergin. 1996. In The Birds of North America. A. Poole and F. Gill, eds. The Academy of Natural Sciences, Philadelphia and The American Ornithologists Union, Washington, D.C. 19 pp.

Johnsgard, P. A. 1986. Birds of the Rocky Mountains with particular reference to national parks in the Northern Rocky Mountain region. Colorado Associated University Press, Boulder. xi + 504 pp.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.



BLUE-GRAY GNATCATCHER

Polioptila caerulea


State rank S1B,SAN

Element code ABPBJ08010

Global rank G5

Modeled by P. McLaughlin

MTNHP status special concern



State range

The only records are in the Glasgow-Fort Peck area and in the Pryor Mountains, Carbon County. The latter, found in 1996, constitutes a small nesting population, which is the first breeding record for the state (Montana Bird Distribution Committee 1996, Wright 1996).

Habitat description

The blue-gray gnatcatcher is on the Montana Natural Heritage Program's Special Concern list due to the discovery of a breeding population in Bear Canyon in the Pryor Mountains (Wright 1996, P. Hendricks pers. comm.). In this area it is associated with Utah juniper and limber pine on the south side of the Pryors; Rocky Mountain juniper may also be present (P. Hendricks pers. comm.). In the west generally, the species frequents chaparral, scrub-oak, and pinyon-juniper habitats (Farrand 1983, Hejl et al. 1995). The species nests especially where tracts of brush, scrub, or chaparral are intermixed with taller vegetation (e.g. forest edge, riparian corridors), often near water. The nest is in the branch or fork of a tree or shrub, usually 1-25 m above ground, by both adults (Harrison 1978). Populations of this species have increased over the past 25 years, expanding northward (Ellison 1992).

Model assumptions & caveats

References

Ellison, W.G. 1992. Blue-gray gnatcatcher (*Polioptila caerulea*). In A. Poole, P. Stettenheim, and F. Gill, (eds.), *The Birds of North America*, No. 23. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union.


Farrand, J., (ed.). 1983. Audubon Society master guide to birding, 3 vols. Alfred A. Knopf, New York. 1244 pp.

Harrison, C. 1978. A field guide to the nests, eggs and nestlings of North American birds. Collins, Cleveland.

Hejl, S.J., R.L. Hutto, C.R. Preston, and D.M. Finch. 1995. Effects of silvicultural treatments in the Rocky Mountains. Pages 220-244 in T. E. Martin and D. M. Finch, eds. Ecology and Management of Neotropical Migratory Birds. Oxford Univ. Press, New York.

Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.

Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



LOGGERHEAD SHRIKE

Lanius ludovicianus


State rank S4B,SZN

Element code ABPBR01030

Global rank G5

Modeled by P. McLaughlin

MTNHP status watch list



Predicted habitat: 10,795,456 ha, 28.35 % of state.

State range

Confirmed breeding throughout the eastern two thirds of the state; east of the Continental Divide is its primary breeding range in the state (Montana Bird Distribution Committee 1996, T. McEneaney, pers. comm.). Migrants recorded throughout the state (Montana Bird Distribution Committee 1996).

Habitat description

Unique as both a passerine and a top-level predator, the loggerhead shrike is one of the few passerines whose populations have decreased throughout the continent in recent decades (Yosef 1996). Breeding Bird Survey (BBS) data show a current decrease of ~3.5-5% / yr across its range (Robbins et al. 1986). The use of organochlorines and pesticides as well as habitat destruction contribute to the species' decline (Yosef 1996). It is listed as a Species of Management Concern in Region 6 (U.S. Fish & Wildlife Service 1995). Dobkin (1994) notes that the species is more or less stable in Montana, although its numbers fluctuate greatly. Based on BBS data, Peterjohn and Sauer (1995) suggest that Montana had stable or increasing populations from 1966-1989; there is evidence in western states that suggests these populations are stable (C. McCarthy, pers. comm.). The loggerhead breeds in open fields and prairies containing scattered trees and shrubs (often spiny), savanna, and open woodland that are usually interspersed with grasses, forbs, and bare ground (Stewart 1975, Johnsgard 1979, AOU 1983, Ehrlich et al. 1988, Yosef 1996). In Montana this open terrain is often heavily grazed (T. McEneaney, pers. comm.) and can be relatively barren of vegetation, such as badlands with few scattered shrubs and trees (E. Atkinson, pers. comm.). In Idaho the species prefers sage-steppe habitat where it nests in sagebrush (65%), bitterbrush, and greasewood and is equally successful in all three (Woods 1993, Woods and Cade 1996). Jewett et al. (1953) suggest that nest sites are often conspicuous bushes relatively larger and more prominent than surrounding sagebrush. It also regularly uses disturbed sites providing open woody growth such as shelterbelts adjacent to croplands, farmsteads, mowed roadsides, and cemeteries (Stewart 1975, Johnsgard 1979, Dobkin 1994, Yosef 1996). Since it favors fence lines and utility lines and poles for perching and takes invertebrates from areas containing bare ground, roadsides are frequented (AOU 1983, Yosef 1996, E. Atkinson, pers. comm.). Suitable hunting

perches are an important part of the habitat (Yosef and Grubb 1994). In Alberta breeding areas, it was found more often than expected in medium (15-35cm) and tall (>35cm) grass (Prescott and Collister 1993). Increasing the prevalence of medium and perhaps tall grasses could mitigate population decreases of the eastern loggerhead shrike east of the Great Plains (Yosef 1996, E. Atkinson, pers. comm.). However, in western Canada it prefers shortgrass pastures (Telfer 1993). Nest sites, which are generally located in open country, are apparently selected based more on degree of cover provided than on particular tree or shrub species. Trees and thickets with thorns are usually preferred for the nest: Russian olive, sagebrush, bitterbrush, greasewood, buffaloberry, elm, and hawthorn (Johnsgard 1979, Yosef 1996, E. Atkinson, pers. comm.). Woodland draws and shelterbelts provide nest sites in Montana and North Dakota (C. McCarthy, pers. comm.).

Model assumptions & caveats

Shrub and tree height cannot be modeled. Mesic shrub (3200) is buffered 500m; within the buffer, agriculture (2010, 2020) and altered herbaceous, very low-low, low-medium, and medium-high grasslands (3110, 3130, 3150, 3170) are selected. Other cover types are included in their entirety. An elevation limit of 1950m (6398') was added to restrict higher elevation predicted habitat.

References


- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Jewett, S.G., W.P. Taylor, W.T. Shaw, and J.W. Aldrich. 1953. Birds of Washington state. Univ. Washington Press, Seattle. 767 pp.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Peterjohn, B.G. and J.R. Sauer. 1995. Population trends of the loggerhead shrike from the North American breeding bird survey. In Shrikes (Laniidae) of the world: biology and conservation (R. Yosef and F.E. Lohrer, eds.). Proc. West. Found. Vert. Zool. 6: 117-121.
- Prescott, D.R.C. and D.M. Collister. 1993. Characteristics of occupied and unoccupied loggerhead shrike territories in southeastern Alberta. J. Wildl. Manage. 57: 346-352.
- Robbins, C.S., D. Bystrak, and P.H. Geissler. 1986. The breeding bird survey: its first fifteen years, 1965-1979. U. S. Fish Wildl. Serv. Resour. Publ.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.
- Telfer, E.S. 1993. Habitat change as a factor in the decline of the western Canadian loggerhead shrike, *Lanius ludovicianus*, population. Can. Field-Nat. 106: 321-326.
- U.S. Fish and Wildlife Service, Office of Migratory Bird Management. 1995. Migratory nongame birds of management concern in the United States: the 1995 list. U.S. Government Printing Office: 1996-404-911/44014. 22 pp.
- Woods, C.P. 1993. Variation in loggerhead shrike nest composition between two shrub species in southwest Idaho. J.

Field Ornithol. 64: 352-357.

Woods, C.P. and T.J. Cade. 1996. Nesting habits of the loggerhead shrike in sagebrush. *Condor* 98: 75-81.

Yosef, R. 1996. Loggerhead shrike (*Lanius ludovicianus*). In A. Poole and F. Gill, (eds.), *The Birds of North America*, No. 231. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union.

Yosef, R. and T.C. Grubb. 1994. Resource dependence and territory size in loggerhead shrikes. *Auk* 111: 465-469.



TENNESSEE WARBLER

Vermivora peregrina


State rank S3S4B,SZN

Element code ABPBX01040

Global rank G5

Modeled by P. McLaughlin

MTNHP status watch list



Predicted habitat: 398,394 ha, 1.05 % of state.

State range

Migrates widely throughout Montana, with indirect evidence of breeding in the Kootenai National Forest-Libby area, as well as in the the Flathead Indian Reservation area and the eastern Glacier National Park area. Direct evidence of breeding in the Flathead National Forest-Whitefish area. No overwintering exhibited (Montana Bird Distribution Committee 1996).

Habitat description


The clutch size of the Tennessee warbler increases in response to outbreaks of forest insects, especially the spruce budworm (Ehrlich et al. 1988, Dobkin 1994, Dunn and Garrett 1997). Although overall population numbers are abundant, the species responds rapidly to budworm outbreaks and crashes, resulting in regional abundance fluctuations from year to year (Dobkin 1994, Dunn and Garrett 1997). A breeder in boreal forests, the Tennessee warbler prefers open, regenerating, or second-growth forests (Johnsgard 1979, AOU 1983, Dunn and Garrett 1997). Bogs and swamps of deciduous (aspens, poplar, willow, alder) or mixed deciduous-coniferous (especially spruce) woodland are preferred habitats, provided there is a well-developed mossy or herbaceous ground layer and an understory (AOU 1983, Ehrlich et al. 1988, Dobkin 1994, Dunn and Garrett 1997). In Alberta and British Columbia the species may be primarily in deciduous associations of aspen, willow, alder, or poplar, usually along streams (Dunn and Garrett 1997); in Montana it prefers deciduous growth as well (T. McEneaney, pers. comm.). In the northern Rockies, spruce forests are occupied (Hejl et al. 1995, Dunn and Garrett 1997). The nest is often hidden at the base of a bush or on a sphagnum hummock, and occasionally on drier hillsides (Ehrlich et al. 1988, Dunn and Garrett 1997). In migration, the species uses a variety of forest, woodland, scrub, and thicket habitats, including urban areas (AOU 1983, Dunn and Garrett 1997).

Model assumptions & caveats

Canopy closure up to 69% is selected. A relatively open tree canopy is selected with the assumption that a well-developed understory is present. Yearly fluctuations in populations cannot be modeled. Species' density augments in areas of bark beetle infestation; food may be limiting factor. Only breeding habitat is modeled, with the assumption that migratory habitat is included as well.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Dunn, J.L. and K.L. Garrett. 1997. A field guide to warblers of North America. Houghton and Mifflin Publ., Boston, Mass. x + 656 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hejl, S.J., R.L. Hutto, C.R. Preston, and D.M. Finch. 1995. Effects of silvicultural treatments in the Rocky Mountains. Pages 220-244 in T. E. Martin and D. M. Finch, eds. Ecology and Management of Neotropical Migratory Birds. Oxford Univ. Press, New York.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.



BLACK-AND-WHITE WARBLER

Mniotilta varia


State rank S2S3B,SZN

Element code ABPBX05010

Global rank G5

Modeled by P. McLaughlin

MTNHP status watch list



State range

Direct evidence of breeding in northeastern Montana, Glasgow-Fort Peck Reservoir area. Indirect breeding evidence observed throughout the eastern part of the state, extending westward only as far as the Northern Cheyenne Indian Reservation area and northward to Fort Peck reservoir. It is a summer resident in the eastern third of Montana (Dobkin 1994). Migrants have been observed extensively east of the Continental Divide, with 2 of 24 sightings west of the divide. No overwintering exhibited (Montana Bird Distribution Committee 1996).

Habitat description


The species is characteristic of semi-open stands of upland forest that are composed of second-growth or mature deciduous or deciduous-coniferous forests; less commonly it uses coniferous forests (Stewart 1975, Johnsgard 1979, Ehrlich et al. 1988, Kricher 1995). Large trees are a critical component of its habitat (Dunn and Garrett 1997). These forest types include mixed stands of juniper / green ash and green ash, American elm, boxelder, and other deciduous trees (Stewart 1975, Johnsgard 1979). Overall, the black-and-white warbler is a habitat generalist with broad tolerances throughout its range (Kricher 1995). James (1971) found that the species showed a strong preference for mature forest over early successional habitat. In central Ontario, it frequents forests with high tree density and high canopy volume, which indicate a preference for mid- to late-successional forests (Clark et al. 1983). The species may also be found in hillside or ravine groves with thin understories and in riparian forests adjacent to grasslands (Johnsgard 1979). Its breeding is restricted to extreme eastern Montana, where it is a bird of woody draws and cottonwood riparian forests (D. Casey and T. McEneaney, pers. comm.). The black-and-white warbler is adapted for gleaning arthropods from tree trunks similar to creepers or nuthatches (Johnsgard 1979, Dobkin 1994). Its nest is usually in a depression on the ground against a shrub, tree, base of a stump, rock, or log (Kricher 1995). It generally migrates east of the Rocky Mountains and is found in interior forest, forest edge, parkland, woodland, second growth, and suburban areas (AOU 1983, Kricher 1995).

Model assumptions & caveats

Of the 25 sightings for the state, six are from the period 1991-1995 (Montana Natural Heritage database). The species was last reported in 1995. Predicted distribution may be overestimated. Size class cannot be modeled.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Clark, K., D. Euler, and E. Armstrong. 1983. Habitat associations of breeding birds in cottage and natural areas in central Ontario. *Wilson Bull.* 95: 77-96.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Dunn, J.L. and K.L. Garrett. 1997. A field guide to warblers of North America. Houghton and Mifflin Publ., Boston, Mass. x + 656 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. *The birder's handbook: a field guide to the natural history of North American birds.* Simon and Shuster, Inc., New York. xxx + 785 pp.
- James, F.C. 1971. Ordination of habitat relationships among breeding birds. *Wilson Bull.* 83: 215-236.
- Johnsgard, P.A. 1979. *Birds of the Great Plains: breeding species and their distribution.* Univ. Nebraska Press, Lincoln. 539 pp.
- Kricher, J.C. 1995. Black-and-white warbler (*Mniotilta varia*). In A. Poole and F. Gill, (eds.), *The Birds of North America*, No. 158. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Stewart, R.E. 1975. *Breeding birds of North Dakota.* Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



DICKCISSEL

Spiza americana


State rank S1S2B,SZN

Element code ABPBX65010

Global rank G5

Modeled by P. McLaughlin

MTNHP status special concern



State range

There are about 15 records in the state; the most recent is from June 1986 at Toston, Broadwater County, which is also the westernmost state record (Wright 1996). There is confirmed breeding only in latilong 34, Custer County. Summer birds have been observed widely throughout eastern 1/4th of state. There is no evidence of overwintering (Montana Bird Distribution Committee 1996).

Habitat description


The dickcissel, a summer resident, is listed as a Species of Management Concern in Region 6 (U.S. Fish and Wildlife Service 1995) and as a sensitive species by the Montana Natural Heritage Program (Montana Bird Distribution Committee 1996). Its numbers are declining significantly throughout its entire range (Dobkin 1994, Johnson and Igl 1995). Pronounced fluctuations in local numbers between years are the rule rather than the exception (Fretwell 1986, Ehrlich et al. 1988, Dobkin 1994). The species breeds in meadows, savanna, brushy fields, ungrazed native prairie, grasslands having tall grasses, forbs, or shrubs, weedy croplands, and in croplands planted to crops such as timothy, alfalfa and clover (Johnsgard 1979, AOU 1983). In North Dakota the species is especially characteristic of alfalfa hayfields; it also frequents sweetclover fields and weedy cropland fields that have been temporarily retired from agriculture (Stewart 1975). Crop mowing and harvesting destroys nests and nestlings, plus the species is a frequent cowbird host (Dobkin 1994). Agricultural lands may pose a significant biological sink for the species (C. McCarthy, pers. comm.). In North Dakota, Conservation Reserve Program (CRP) fields are especially important for the dickcissel during the breeding season; if CRP habitat were returned to cropland and densities of breeding birds were substituted in cropland for densities in CRP, the dickcissel's numbers would be reduced by 17.1% (Johnson and Igl 1995).

Model assumptions & caveats

There are too few observers in eastern Montana to allow for accurate determination of the species' presence (Wright 1996). Two cover types were chosen for this model: altered herbaceous grasslands (3110) and moderate/high cover grasslands (3170). CRP land (included in 3110) is often non-native species. Agricultural cover types were not selected since crop types (per Johnsgard 1979) cannot be specified and will vary from year to year. The low-medium grassland (3150) cover type, the state's largest at >10 million ha, was not included because its selection would largely overestimate predicted habitat.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Fretwell, S. 1986. Distribution and abundance of the dickcissel. *Current Ornithology* 4: 211-242.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Johnson, D.H. and L.D. Igl. 1995. Contributions of the Conservation Reserve Program to populations of breeding birds in North Dakota. *Wilson Bull.* 107(4): 709-718.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.
- U.S. Fish and Wildlife Service, Office of Migratory Bird Management. 1995. Migratory nongame birds of management concern in the United States: the 1995 list. U.S. Government Printing Office: 1996-404-911/44014. 22 pp.
- Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



BAIRD'S SPARROW

Ammodramus bairdii


State rank S3S4B,SZN

Element code ABPBXA0010

Global rank G4

Modeled by P. McLaughlin

MTNHP status special concern



State range

Summering birds occur in north-central and the eastern half of Montana (Bergeron et al. 1992). Direct evidence of breeding found only in northern half of the state, east of the Continental Divide, i.e. in the Choteau and Great Falls area, and east in the Fort Peck Indian reservation-Plentywood area. Only migrants found west of the divide and no overwintering exhibited throughout the state (Montana Bird Distribution Committee 1996). The species breeds extensively throughout northern Phillips and Blaine counties and Bowdoin NWR, where the most nests have been documented in the state 1994-1997 (D. Prellwitz, pers. comm.). Medicine Lake is also a stronghold for this species (T. McEneaney, pers. comm.).

Habitat description

A local and generally uncommon species, the Baird's sparrow is a mixed-grass prairie specialist endemic to the northern Great Plains (Dobkin 1994, Rising 1996, C. McCarthy, pers. comm.). It is a Species of Management Concern in Region 6 (U.S. Fish and Wildlife Service 1995). However, the species is very common in ungrazed grasslands in northeastern and north-central Montana (D. Prellwitz, pers. comm.). Its optimal habitats include extensive ungrazed or lightly grazed tracts of mixed-grass prairie and local pockets of wet meadow zones or tall-grass prairie that are associated with wetlands and intermittent streams (Stewart 1975, Johnsgard 1979, Rising 1996). These grasslands may be interspersed with scattered clumps of grass and low shrubs with tangled, matted grass on the ground (A.O.U. 1983, Rising 1996), although it generally prefers grasslands having little or no shrub cover (Dobkin 1994). Large populations in northeastern Montana are located in native mixed-grass grasslands that have not been grazed by cattle or disturbed by fire for approximately 20 years (Hammond et al. 1997). Such habitat is found at Bowdoin NWR, where the nest success for the Baird's sparrow in 1997 was 46.6%, slightly above the mean nest success of 42% reported for open-cup nesting neotropical migrants in North America (Hammond et al. 1997). Based on 1995-1997 data, current populations have increased from the period 1979-1981 when transects were done by the Bureau of Land Management (D. Prellwitz, pers. comm.). Conservation Reserve Program (CRP) and crested wheatgrass

(CWG) fields have in part contributed to this population increase (D. Prellwitz, pers. comm.). The species will use vegetation <2' tall, but will not use taller, 3'-3.5' vegetation of moister areas as in parts of North Dakota and the Midwest (D. Prellwitz, pers. comm.). Although it especially prefers native grasslands over disturbed grasslands (Owens and Myres 1973), the Baird's will use disturbed habitats such as retired croplands, hay and alfalfa fields, crested wheatgrass, and weedy stubble fields (Johnsgard 1979, Dobkin 1994, C. McCarthy and D. Prellwitz, pers. comm.). While agricultural practices and grazing have decreased available habitat (Ehrlich et al. 1988), the species can tolerate light grazing regimes (Kantrud and Kologiski 1982). The Baird's sparrow will use a forb or small hummock for a perch, but will not use a fence or wire for elevation (D. Prellwitz, pers. comm.). This may bias population censuses such as Breeding Bird Surveys (BBS) that rely on driven routes to gather data (D. Prellwitz, pers. comm.). The species is thought to be highly nomadic across its breeding range: in wet years it commonly uses suitable grasslands of eastern Montana and the Dakotas; in dry years it appears to be more common in suitable grasslands of the Canadian prairie (C. McCarthy, pers. comm.).


Model assumptions & caveats

Breeding habitat only is modeled; assumed potential migration habitats defined within breeding habitat parameters. Irrigated agriculture (2020) is selected because the species will use croplands that follow a rest-rotation regime; such land-use practices cannot be modeled. Seasonal weather patterns affecting the species' distribution cannot be modeled.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Bergeron, D., C. Jones, D.L. Genter, and D. Sullivan. 1992. P. D. Skaar's Montana Bird Distribution, Fourth Edition. Special Publication No. 2. Montana Natural Heritage Program, Helena. 116 pp.
- Dobkin, D.S. 1994. Conservation and management of neotropical migrant landbirds in the Northern Rockies and Great Plains. Univ. Idaho Press, Moscow, Idaho. 220 pp.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Hammond, J., J. Wood, P. Gouse, and S. Jones. 1997. Grassland songbird reproduction monitoring on the Bowdoin National Wildlife Refuge, special emphasis on Baird's sparrow and Sprague's pipit: results from the 1997 field season. Point Reyes Bird Observatory and U. S. D. I. Fish & Wildlife Service Mountain-Prairie Region. 7 pp.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Kantrud, H.A. and R.L. Kologiski. 1982. Effects of soils and grazing on breeding birds of uncultivated upland grasslands of the northern Great Plains. U.S.D.I., Fish and Wildl. Serv., Wildl. Res. Rep. 15. 33 pp.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Owens, R.A. and M.T. Myres. 1973. Effects of agriculture upon populations of native passerine birds of an Alberta fescue grassland. Can. J. Zool. 51: 697-713.
- Rising, J.D. 1996. A guide to the identification and natural history of the sparrows of the United States and Canada. Academic Press, Inc., San Diego, CA. 365 pp.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.

U.S. Fish and Wildlife Service, Office of Migratory Bird Management. 1995. Migratory nongame birds of management concern in the United States: the 1995 list. U.S. Government Printing Office: 1996-404-911/44014. 22 pp.



LE CONTE'S SPARROW

Ammodramus leconteii


State rank S1S2B,SZN

Element code ABPBXA0040

Global rank G4

Modeled by P. McLaughlin

MTNHP status special concern



State range

Confirmed breeding is only on Camas Creek, Glacier National Park, and in Sheridan County in northeastern Montana. Transients recorded rarely throughout the state, though distributed east and west of the divide, as well as northern and southern Montana. No overwintering exhibited (Montana Bird Distribution Committee 1996, Wright 1996).

Habitat description

Listed as a sensitive species in Montana (Montana Bird Distribution Committee 1996), the LeConte's sparrow is an uncommon breeder with a disjunct and restricted breeding range in the state (T. McEneaney, pers. comm.). Preferred breeding habitats include moist grass or sedge meadows that are occasionally mixed with cattails or phragmites, and often interspersed with small alders and birches; and hummocky alkaline wetlands (fens) (Johnsgard 1979, A.O.U. 1983, Lowther 1996, Rising 1996). The species is less commonly found in lowland tracts of tall-grass prairie and wet-meadow zones on the periphery of prairie ponds and lakes or along intermittent streams (Stewart 1975, Johnsgard 1979, Ehrlich et al. 1988). Disturbed habitats occupied are hayfields and reitured croplands, and in migration it is found in tall, dense grasses, damp weedy fields, or cattails (A.O.U. 1983, Rising 1996). In mountainous areas, it is found in flooded sedge and grass meadows (Semenchuk 1992). Despite its preference for marshy, wet areas, moist habitats do not seem to be a requirement (Lowther 1996): North Dakota censuses have found LeConte's sparrows in 6-9 year-old fields of alfalfa and wheatgrass (Renken and Dinsmore 1987). Population densities can fluctuate yearly; usually this species reaches peak numbers in wet years, and may be scarce or absent during dry years (Stewart 1975, Lowther 1996). In 1994 during surveys of Conservation Reserve Program fields in Sheridan County, 52 pairs were found (Wright 1996).


Model assumptions & caveats

The species responds negatively to any level of livestock grazing in mixed-grass prairie (Maher 1979).

Altered herbaceous (3110) and montane parklands and subalpine meadows (3180) cover types are selected in their entirety. Altered herbaceous (3110) includes, as well as CRP fields, disturbed lands with noxious weeds and bare soil coverages from 10-50%. However, the noxious weed component of 3110 is primarily included in the southern and western parts of the state. Graminoid & forb riparian (6200) is selected only within a 500m hydrography buffer of marshes / swamps. Breeding habitat only is modeled; assumed potential migration habitats defined within breeding habitat parameters.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Lowther, P.E. 1996. Le Conte's Sparrow (*Ammodramus leconteii*). In A. Poole and F. Gill (eds.), The Birds of North America, No. 224. Philadelphia: The Academy of Natural Sciences; Washington D.C.: The American Ornithologists Union. 16 pp.
- Maher, W.J. 1979. Nestling diets of prairie passerine birds at Matador, Saskatchewan, Canada. *Ibis* 121:437-452.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Renken, R.B. and J.J. Dinsmore. 1987. Nongame bird communities on managed grasslands in North Dakota. *Can. Field-Nat.* 101: 551-557.
- Rising, J.D. 1996. A guide to the identification and natural history of the sparrows of the United States and Canada. Academic Press, Inc., San Diego, CA. 365 pp.
- Semenchuk, G.P., ed. 1992. The atlas of breeding birds of Alberta. Fed. Alberta Nat., Edmonton.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.
- Wright, P.L. 1996. Status of rare birds in Montana, with comments on known hybrids. *Northwestern Nat.* 77(3):57-85.



NELSON'S SHARP-TAILED SPARROW

Ammodramus nelsoni


State rank S1B,SZN

Element code ABPBXA0070

Global rank G5

Modeled by P. McLaughlin

MTNHP status special concern



State range

Three records of breeding evidence; one direct and two indirect, both in the Plentywood-Fort Peck Indian Reservation area of northeastern Montana. No transient or overwintering records (Montana Bird Distribution Committee 1996).

Habitat description


A highly localized species, the Nelson's sharp-tailed sparrow is listed as a sensitive species in the state, and breeds only in extreme northeastern Montana (Greenlaw and Rising 1994, Montana Bird Distribution Committee 1996, Rising 1996). In breeding season, the species is restricted to wet meadows and grasslands, the edges of marshes, herbaceous wetlands, and alkaline hummocky bogs (fens) (Johnsgard 1979, A.O.U. 1983, Ehrlich et al. 1988, Greenlaw and Rising 1994, Rising 1996). Within fens, it is largely restricted to patches of coarse emergent vegetation including cattails, bulrushes, and phragmites (Stewart 1975). When water levels are low, the species will use the marshy zones of prairie lakes and ponds and shallow-marsh and deep-marsh zones (Stewart 1975, Johnsgard 1979). Locally fairly common in the prairie pothole region of North Dakota, the species' populations often fluctuate greatly in number from year to year. Unlike the LeConte's sparrow, Nelson's sharp-tailed sparrow populations reach their peak numbers in drought years and are reduced to minimal number during wet years (Stewart 1975). In migration it may occur in wet fields and marshes, often in cattails, and in brushy areas and overgrown fields (A.O.U. 1983, Rising 1996).

Model assumptions & caveats

The Nelson's sharp-tailed sparrow responds negatively to any level of livestock grazing in mixed-grass prairie (Maher 1979). Breeding habitat only is modeled; assumed that potential migration habitats defined within breeding habitat parameters.

References

- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Amer. Ornithologists' Union, Washington, D.C.
- Ehrlich, P.R., D.S. Dobkin, and D. Wheye. 1988. The birder's handbook: a field guide to the natural history of North American birds. Simon and Shuster, Inc., New York. xxx + 785 pp.
- Greenlaw, J.S. and J.D. Rising. 1994. Sharp-tailed sparrow (*Ammodramus caudacutus*). In A. Poole and F. Gill, eds. The birds of North America, No. 112. The Academy of Natural Sciences, Philadelphia, PA, and the American Ornithologists' Union, Washington, DC.
- Johnsgard, P.A. 1979. Birds of the Great Plains: breeding species and their distribution. Univ. Nebraska Press, Lincoln. 539 pp.
- Maher, W.J. 1979. Nestling diets of prairie passerine birds at Matador, Saskatchewan, Canada. *Ibis* 121:437-452.
- Montana Bird Distribution Committee. 1996. P. D. Skaar's Montana Bird Distribution, Fifth Edition. Special Publication No. 3. Montana Natural Heritage Program, Helena. 130 pp.
- Rising, J.D. 1996. A guide to the identification and natural history of the sparrows of the United States and Canada. Academic Press, Inc., San Diego, CA. 365 pp.
- Stewart, R.E. 1975. Breeding birds of North Dakota. Tri-College Center for Environmental Studies, Fargo, North Dakota. 295 pp.



PREBLE'S SHREW

Sorex preblei


State rank S3

Element code AMABA01030

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



State range

Thought to be rare (Flath 1984). However, range has expanded as sampling has increased. Possibly more widespread than previously apparent (Hoffmann and Fisher 1978).

Habitat description

Most Montana specimens from dry sage and sage-grasslands (Hoffman and Pattie 1968). Also riparian shrubs and Douglas-fir forest (Allen et al. 1994; Ports and George 1990). Captured in subalpine conifers in eastern Washington, may be restricted by *S. cinereus*. In Utah: saltgrass, greasewood, sagebrush and sage openings in conifers (Tomasi and Hoffmann 1984).

Model assumptions & caveats

Limited habitat association information. May be found near rock outcrops. Cannot model for rock outcrops, hence distribution likely overestimated.

References

Allen, K.L., T. Weaver, and D. Flath. 1994. Small mammals in Northern Rocky Mountain ecosystems. Unpubl. report to Bureau of Land Management and United States Forest Service, August 31, 1994. Montana State Univ., Bozeman. 54pp.

Cornely, J. E., L. N. Carraway, and B. J. Verts. 1992. *Sorex preblei*. Mamm. Species 416: 1-3.

Flath, D. L. 1984. Vertebrate species of special interest or concern. Mammals, birds, reptiles, amphibians, fishes. Spec.

Publ. Montana Department of Fish, Wildlife, and Parks, Helena. 76 pp.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.


Hoffmann, R. S. and R. D. Fisher. 1978. Additional distributional records of Preble's shrew (*Sorex preblei*). J. Mammal. 59: 883-884.

Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. J. Mammal. 50(3): 579-604.

Long, C. A. and R. S. Hoffmann. 1992. *Sorex preblei* from the Black Canyon, first record from Colorado. Southwest. Nat. 37: 318-319.

Ports, M. A. and S. B. George. 1990. *Sorex preblei* in the northern Great Basin. Great Bas. Nat. 50: 93-95.

Tomasi, T. E. and R. S. Hoffmann. 1984. *Sorex preblei* in Utah and Wyoming. J. Mammal. 65: 708.



DWARF SHREW

Sorex nanus


State rank S3

Element code AMABA01130

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



State range

Known from several high mountain ranges in central Montana: Sweetgrass Hills (Thompson 1977); Bears Paw Mountains (Thompson 1977); Little Belt Mountains (Hoffmann and Taber 1960); Beartooth Mountains (Pattie and Verbeek 1967, Hoffmann and Taber 1960). More recently found at low elevation in southeastern Montana (MacCracken et al. 1985). May be more widespread than currently known (Flath 1984). Has been captured in low elevation prairie habitat in South Dakota. Known from the Holocene in the East Pryor Mountains (Geppert 1984).

Habitat description


Spruce-fir-lodgepole forest, forest meadows and clearcuts, will occupy drier forest than other shrews (Hoffmann and Owen 1980). Open ponderosa pine (Allen et al. 1994). Marsh and riparian areas. Will occupy sage at high population densities (MacCracken et al. 1985). Rocky alpine tundra (Hoffmann and Owen 1980). Individuals found on alpine and subalpine talus slopes in Sweetgrass Hills, Bears Paw Mountains (Thompson 1977), Hellroaring Plateau (Pattie and Verbeek 1967) and Medicine Bow Mountains (Brown 1967).

Model assumptions & caveats

References

Allen, K.L., T. Weaver, and D. Flath. 1994. Small mammals in Northern Rocky Mountain ecosystems. Unpubl. report to Bureau of Land Management and United States Forest Service, August 31, 1994. Montana State Univ., Bozeman. 54pp.

- Berna, H. J. 1990. Observations on the dwarf shrew (*Sorex nanus*) in northern Arizona. *Great Basin Nat.* 50: 161-165.
- Brown, L.N. 1967. Ecological distribution of six species of shrews and comparison of sampling methods in the central Rocky Mountains. *J. Mammal.* 48(4): 617-623.
- Cinq-Mars, R.J., Hoffmann, R.S., Jones, J.K. 1979. New records of the dwarf shrew (*Sorex nanus*) in South Dakota. *Prairie Nat.* 2(1): 7-9.
- George, S. B. 1990. Unusual records of shrews in New Mexico. *Southwest. Nat.* 35:464-465.
- Geppert, T. J. 1984. Small mammals of the Shield Trap, East Pryor Mountain, Montana. M.S. thesis. University of Iowa, Iowa City. 45 pp.
- Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.
- Hoffmann, R.S. and J.G. Owen. 1980. *Sorex tenellus* and *Sorex nanus*. *Mamm. Species* 131:1-4.
- Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.
- MacCracken, J. G., D. W. Uresk, and R. M. Hansen. 1985. Habitat used by shrews in southeastern Montana. *Northwest Science* 59(1):24-27.
- Mickey, A.B. 1948. A record of the shrew *Sorex nanus* for WY. *J. Mammal.* 29(3):295.
- Pattie, D. L. and N. A. M. Verbeek. 1967. Alpine mammals of the Beartooth Plateau. *Northwest Sci.* 41(3): 110-117.
- Spencer, A. W. and D. Pettus. 1966. Habitat preferences of five sympatric species of long-tailed shrews. *Ecology* 47: 677-683.
- Thompson, L.S. 1977. Dwarf shrew in north-central Montana. *J. Mammal.* 58:248-250.




MERRIAM'S SHREW

Sorex merriami

State rank S3
Global rank G5
MTNHP status special concern

Element code AMABA01230
Modeled by P. Thornton



State range

Thought to be rare (Flath 1984). Patchy distribution in the eastern half of the state (Thompson 1982).

Habitat description

Dry sagebrush, sage-grasslands and grasslands (Brown 1967, Armstrong and Jones 1971, MacCracken et al. 1985). Riparian shrubs. Fencerows of agricultural fields (Hooper 1944). Open ponderosa pine (Allen et al. 1994).

Model assumptions & caveats

References

Allen, K.L., T. Weaver, and D. Flath. 1994. Small mammals in Northern Rocky Mountain ecosystems. Unpubl. report to Bureau of Land Management and United States Forest Service, August 31, 1994. Montana State Univ., Bozeman. 54pp.


Armstrong, D. M. and J. Knox Jones, Jr. 1971. *Sorex merriami*. Mamm. Species 2:1-2.

Brown, L.N. 1967. Ecological distribution of six species of shrews and comparison of sampling methods in the central Rocky Mountains. J. Mammal. 48(4): 617-623.

Diersing, V.E. 1979. Noteworthy records of Merriam's shrew from New Mexico southwest. Nat. 24(4):708-709.

Gashwiler, J.S. 1976. A new distribution record of Merriam's shrew in Oregon. Murrelet. 57:13-14.

- George, S. B. 1990. Unusual records of shrews in New Mexico. *Southwest. Nat.* 35:464-465.
- Geppert, T. J. 1984. Small mammals of the Shield Trap, East Pryor Mountain, Montana. M.S. thesis. University of Iowa, Iowa City. 45 pp.
- Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.
- Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.
- Hooper, E.T. 1944. Additional records of Merriam's shrew in Montana. *J. Mammal.* 25:92.
- Johnson, M.L. and C.W. Clanton. 1954. Natural history of *Sorex merriami* in Washington state. *Murrelet.* 35:1-4.
- MacCracken, J. G., D. W. Uresk, and R. M. Hansen. 1985. Habitat used by shrews in southeastern Montana. *Northwest Science* 59(1):24-27.
- Mickey, A.B. 1947. A record of *Sorex merriami* for southeastern Wyoming. *J. Mammal.* 28:293.
- Mickey, A.B. and C.N. Steele, Jr. 1946. A record of *Sorex merriami* for southeastern WY. *J. Mammal.* 28(3):293.
- Mullican, T.R. 1986. Additional records of *Sorex merriami* from Idaho. *Murrelet.* 67:19-20.
- Pefaur, J. E. and R. S. Hoffmann. 1971. Merriam's shrew and hispid pocket mouse in Montana. *Am. Midl. Nat.* 86(1):247-248.
- Ports, M. A. and S. B. George. 1990. *Sorex preblei* in the northern Great Basin. *Great Bas. Nat.* 50: 93-95.
- Thompson, L.S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Bozeman: Montana Audubon Council. 24 pp.




FRINGED MYOTIS

Myotis thysanodes

State rank S3
Global rank G5
MTNHP status special concern

Element code AMACC01090
Modeled by P. Thornton



Predicted habitat: 4,661,200 ha, 12.24 % of state.

State range

Range records of *Myotis thysanodes* are very limited (Hoffmann et al. 1969). Specimens from Lewis and Clark Caverns and Ravalli county exist (Flath 1984). This species is probably found throughout southwestern Montana (Hoffmann et al. 1969).

Habitat description

Primarily at middle elevations of 1,200-2,150 m in desert, grassland, sage, and juniper habitats; has been recorded at 2,850 m in spruce-fir forests in New Mexico, and at low elevations along Pacific Coast. Roosts in caves, mines, rock crevices, buildings, and other protected sites.

Model assumptions & caveats

References

Boyce, Mark S. October 8, 1968. First Record Of The Fringe-Tailed Bat, *Myotis thysanodes*, From Southeastern Wyoming. Southwest. Nat. 25:114-115.

Flath, D. L. 1984. Vertebrate species of special interest or concern. Mammals, birds, reptiles, amphibians, fishes. Spec. Publ. Montana Department of Fish, Wildlife, and Parks, Helena. 76 pp.

Hoffmann, R.S., D.L. Pattie and J.F. Bell. 1969. The distribution of some mammals in Montana. II. Bats. J. Mammal. 50(4): 737-741.


Manning, R. W. and J. K. Jones, Jr. 1988. A new subspecies of fringed myotis, *Myotis thysanodes*, from the

northwestern coast of the United States. Occas. Pap. Mus. Texas Tech Univ. No. 123:1-6.

Manning, R. W., and J. K. Jones, Jr. 1988. A new subspecies of fringed myotis, *Myotis thysanodes*, from the northwestern coast of the United States. Occas. Pap. Mus. Texas Tech Univ. No. 123:1-6.

O'Farrell, M. J., and E. H. Studier. 1973. Reproduction, growth, and development in *Myotis thysanodes* and *M. lucifugus* (Chiroptera: Vespertilionidae). *Ecology* 54:18-30.

O'Farrell, M.J. and E.H. Studier. 1980. *Myotis thysanodes*. *Mamm. Species* 137:1-5.



YUMA MYOTIS

Myotis yumanensis


State rank S3

Element code AMACC01020

Global rank G5

Modeled by P. Thornton

MTNHP status watch list



State range

Widely distributed in western Montana, but probably very limited in occurrence east of the Continental Divide (Hoffmann et al. 1969).

Habitat description

More closely associated with water than most other North American bats. Found in a wide variety of upland and lowland habitats, including riparian, desert scrub, moist woodlands and forests, but usually found near open water. Flies low over flowing water to forage (van Zyll de Jong 1985).

Model assumptions & caveats

References

Aldrich, Hugh. 1986. Manoeuverability and ecological segregation in little brown (*Myotis lucifugus*) and Yuma (*M. yumanensis*) bats (Chiroptera: Vespertilionidae). *Can. J. Zool.* 64:1878-1882.

Brigham, R.M. August 1992. Variation in Habitat Use and Prey Selection by Yuma Bats, *Myotis yumanensis*. *Journal of Mammalogy* 73(3):640-645.

Brigham, R. M., H. D. J. N. Aldridge, and R. L. Mackey. 1992. Variation in habitat use and prey selection by Yuma bats, *Myotis yumanensis*. *J. Mammal.* 73:640-645.

Dalquest, W.W. 1947. Notes on the natural history of the bat, *Myotis yumanensis*, in California with a description of a new race. *Amer. Midl. Nat.* 38:224-247.

Harris, A. H. 1974. *Myotis yumanensis* in interior southwestern North America with comments on *Myotis lucifugus*. *J. Mammal.* 55:589-607.


Herd, R. M. and M. B. Fenton. 1983. An electrophoretic, morphological, and ecological investigation of a putative hybrid zone between *Myotis lucifugus* and *Myotis yumanensis* (Chiroptera: Vespertilionidae). *Can. J. Zool.* 61:2029-2050.

Herd, R. M. and M. B. Fenton. 1983. An electrophoretic, morphological, and ecological investigation of a putative hybrid zone between *Myotis lucifugus* and *Myotis yumanensis* (Chiroptera: Vespertilionidae). *Can. J. Zool.* 61(9):2029-2050.

Herd, R. M., and M. B. Fenton. 1983. An electrophoretic, morphological, and ecological investigation of a putative hybrid zone between *Myotis lucifugus* and *Myotis yumanensis* (Chiroptera: Vespertilionidae). *Can. J. Zool.* 61:2029-2050.

Parkinson, A. 1979. Morphological variation and hybridization in *Myotis yumanensis sociabilis* and *Myotis lucifugus carrisma*. *J. Mammal.* 60:489-504.

van Zyll de Jong, C.G. 1985. Handbook of Canadian mammals. Vol 2. Bats. National Museums of Canada, Ottawa. 212 pp.



NORTHERN MYOTIS

Myotis septentrionalis


State rank S2

Element code AMACC01150

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



State range

This bat is known from a few sites in eastern Montana along the lower Missouri River, and in Montana is probably confined to riparian corridors of the lower Missouri River system (MT-NHP VCA).


Habitat description

In Montana, Northern Myotis has been located hibernating in an abandoned mine in river breaks habitat in Richland County (Swenson and Shanks 1979).

Model assumptions & caveats

References

Swenson, J.E., and G.F Shanks, Jr. 1979. Noteworthy records of bats from northeastern Montana. *J. Mammal.* 60:650-652




SPOTTED BAT

Euderma maculatum

State rank S1
Global rank G4
MTNHP status special concern

Element code AMACC07010
Modeled by P. Thornton



State range

Few definite locations in Montana; all from south-central part of state (Thompson, 1982, MT-NHP VCA).

Habitat description

Found in various habitats from desert to montane coniferous stands, including open ponderosa pine, pinyon-juniper woodland, open pasture, and hayfields. Speculation has been made that captures outside coniferous forests reflect post-breeding wandering (Snow 1974). Roosts in caves and in cracks and crevices in cliffs and canyons, with which this species consistently is associated (Snow 1974, van Zyll de Jong 1985). Winter habits poorly known. Handley (1959) found that spotted bats were found primarily on open or scrub country. Of his 22 recorded occurrences, 13 were around houses.

Model assumptions & caveats

References

Handley, C. O., Jr. 1959. A revision of American bats of the genera *Euderma* and *Plecotus*. Proc. U.S. Natl. Mus. 110:95-246.


Leonard, M. L. and M. B. Fenton. 1983. Habitat use by spotted bats (*Euderma maculatum*, Chiroptera: Vespertilionidae): roosting and foraging behavior. Can. J. Zool. 61:1487-1491.

Snow, C. 1974. Spotted Bat. Habitat Management Series for Endangered Species, Report No. 4. Bureau of Land Management. 13 pp.

Thompson, L.S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Bozeman: Montana Audubon Council. 24 pp.

van Zyll de Jong, C.G. 1985. Handbook of Canadian mammals. Vol 2. Bats. National Museums of Canada, Ottawa. 212 pp.

Watkins, L.C. 1977. *Euderma maculatum*. Mamm. Species 77:1-4.



TOWNSEND'S BIG-EARED BAT

Corynorhinus townsendii


State rank S2S3

Element code AMACC08010

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



Predicted habitat: 14,300,158 ha, 37.55 % of state.

State range

Scattered records imply that this species has a statewide distribution (Thompson 1982). Only two confirmed breeding colonies; several confirmed hibernacula.

Habitat description

Generally found in low densities, occupying a range of habitats including moist forests (Thomas and West 1991) as well as arid regions (Genter and Metzgar 1985), including desert shrub and junipers. In western Montana they are most closely associated with cavernous habitat and rocky outcrops of sedimentary or limestone origin, which are used for roosting. In old-growth forests, large diameter hollow trees may be used for roosting. Occasionally, individuals may be found in buildings. Maternity colonies are found in warm areas of caves, mines and occasionally buildings (Pearson et al. 1952, D. Genter, pers. obs., Idaho State Conservation Effort 1995). Hibernacula are typically in caves or mines with winter temperatures 2-7° C and relative humidity >50%.

Model assumptions & caveats

References

Butts, T. W. 1993. Azure Cave bat surveys, Little Rocky Mountains, Montana, September 1992 and March 1993. Unpublished report for Zortman Mining, Inc. 13 pp.

Fitzgerald, T. 1989. New records of bats from northeastern Colorado. *Journal of the Colorado-Wyoming Academy of Science* 21:22.

Genter, D. L. 1986. Wintering bats of the upper Snake River plain: occurrence in lava-tube caves. *Great Basin Nat.* 46(2):241-244.

Genter, D. L. and L. H. Metzgar. 1985. Survey of the bat species and their habitat use in Grand Teton National Park. Pp. 65-69 in: *Univ. Wyo.-Natl. Park Serv. Res. Center. Annual report 9.*

Hoffmann, R.S., D.L. Pattie and J.F. Bell. 1969. The distribution of some mammals in Montana. II. Bats. *J. Mammal.* 50(4): 737-741.


Idaho State Conservation Effort. 1995. Habitat conservation assessment and conservation strategy for the Townsend's big-eared bat. Draft unpubl. rep. no. 1. Boise, Id.

Madson, M., G. Hanson, S. Martinez, and D. Genter. 1993. Wintering bats in Montana: results of surveys in the Pryor Mountains with annotation on area caves and mines. *Montana Natural Heritage Program. Helena, MT.* 36 pp.

Marcot, B. G. 1984. Winter use of some northwestern California caves by western big-eared bats and long-eared *Myotis*. *Murrelet* 65(2):46.

Pearson, O.P., M.R. Koford, and A.K. Pearson. 1952. Reproduction of the lump-nosed bat (*Corynorhinus townsendii*) in California. *J. Mammal.* 33:273-320.

Thomas, D.W. and S.D. West. 1991. Forest age associations of bats in the southern Washington Cascade and Oregon Coast Ranges. Pp. 295-303 in: Ruggiero, L.F., K.B. Aubry, A.B. Carey, and M.H. Huff (tech. coord.). *Wildlife and vegetation of unmanaged Douglas-fir forests. U.S.D.A. Forest Serv., Pac. Northw. Res. Station. Gen. Tech. Report PNW-GTR-285.* 533 pp.



PALLID BAT

Antrozous pallidus


State rank S1

Element code AMACC10010

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



State range

First state record was in Billings area (Shryer and Flath 1980); otherwise no available information (MT-NHP VCA).

Habitat description

Arid deserts and grasslands, often near rocky outcrops and water. Less abundant in evergreen and mixed conifer woodland. Usually roosts in rock crevice or building, less often in cave, tree hollow, mine, etc. Night roosts often or typically are in caves in Oklahoma (Caire et al. 1989). In Oregon, night roosts were in buildings, under rock overhangs, and under bridges; bats generally were faithful to particular night roosts both within and between years (Lewis 1994). Prefers narrow crevices in caves as hibernation sites (Caire et al. 1989). Young are born in maternity colonies usually in rock crevices or buildings.

Model assumptions & caveats

References

Caire, W., J.D. Tyler, and B.P Glass. 1989. Mammals of Oklahoma. Univ. Oklahoma Press, Norman. xiii + 567 pp.


Hermanson, J. W. and T. J. O'Shea 1983. *Antrozous pallidus*. Mamm. Species 213:1-8.

Lewis, S. E. 1994. Night roosting ecology of pallid bats (*Antrozous pallidus*) in Oregon. Am. Midl. Nat. 132:219-226.

Orr, R.T. 1954. Natural history of the pallid bat, *Antrozous pallidus* (le conte). Proc. California Acad. Sci., 82;

165-246.

Shryer, J. and D. Flath. 1980. First record of the pallid bat (*Antrozous pallidus*) from Montana. *Great Basin Nat.* 40:115.



EASTERN COTTONTAIL

Sylvilagus floridanus


State rank S2?

Element code AMAEB01040

Global rank G5

Modeled by P. Thornton

MTNHP status watch list



State range

Only known to occur in Carter County, southeastern Montana, where it is apparently restricted to riparian habitats (Lampe et al. 1974). Western edge of its range is in the Great Plains.

Habitat description

Wide variety of habitats (Chapman et al. 1980), but prefers dense, shrubby undergrowth (Hoffmann and Pattie 1968). Riparian shrubs. Hardwood draws along intermittent streams (Bergeron and Seabloom 1981).

Model assumptions & caveats


References

Bergeron, D.J. and R.W. Seabloom. 1981. Habitat partitioning by eastern and desert cottontails in southwest North Dakota. *Prairie Nat.* 13:105-110.

Chapman, J. A., J. G. Hockman, and M. M. Ojeda. 1980. *Sylvilagus floridanus*. *Mamm. Species* 136:1-8.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Lampe, R. P., Jones Jr., J. K., Hoffmann, R. S., and E. C. Birney. 1974. The mammals of Carter County, southeastern Montana. *Occa. Pap. Mus. Nat. Hist. Univ. Kan.* 25:1-39.



BLACK-TAILED JACKRABBIT

Lepus californicus


State rank S2S3

Element code AMAEB03050

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



State range

May have entered Montana during this century. Occurs in Beaverhead and Madison Counties.

Habitat description

Sagebrush; short grass prairie with or without brush (Hoffmann and Pattie 1968, Flinders and Hansen 1973, Johnson and Hansen 1979).

Model assumptions & caveats

Number of snow-free days important for defining range limits, but not included in the model.

References


Chapman, J. A. and G. A. Feldhamer, eds. 1982. Wild mammals of North America: biology, management, and economics. Johns Hopkins Univ. Press. Baltimore, MD. 1147 pp.

Flinders, J. T. and R. M. Hansen. 1973. Abundance and dispersion of Leporids within a shortgrass ecosystem. J. Mammal. 54:287-29.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Johnson, M. K. and R. M. Hansen. 1979. Foods of cottontails and woodrats in southcentral Idaho. J. Mammal. 60:213-215.

Smith, G. W. 1990. Home range and activity patterns of black-tailed jackrabbits. Great Basin Nat. 50:249-256.



PYGMY RABBIT

Brachylagus idahoensis


State rank S2S3

Element code AMAEB04010

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



State range

Suitable sagebrush habitat in Beaverhead County and possibly adjacent counties (Hoffmann et al. 1969).

Habitat description

Sagebrush and greasewood. Tall clumps of Big Sage (Green and Flinders 1980), shrub canopy cover > 21%. Soil depth > 36cm; loose, friable soil (Weiss and Verts 1984). At least 46% sage cover on US Sheep Experimental Station; bushes 56cm in height; Snake River Plain sage habitat (Johnson and Hansen 1979).

Model assumptions & caveats

Loose soil originally defined as soil texture other than clay; soil later eliminated from the model because scale of available statewide layer was too coarse.

References

Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.

Campbell, T.M., Clark, T.W. and Groves, C.R. 1982. First record of pygmy rabbits (*Brachylagus idahoensis*) in Wyoming. *Great Basin Nat.* 42:100.

Green, J.S. and Flinders, J.T. Habitat and dietary relationships of the pygmy rabbit. *J. Range Manage.* 33(2):136-142.


Green, J. S. and J. T. Flinders. 1980. *Brachylagus idahoensis*. *Mamm. Species* 125:1-4.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. J. Mammal. 50(3): 579-604.

Johnson, M. K. and R. M. Hansen. 1979. Foods of cottontails and woodrats in southcentral Idaho. J. Mammal. 60:213-215.

Weiss, N. T. and B. J. Verts. 1984. Habitat and distribution of pygmy rabbits (*Sylvilagus idahoensis*) in Oregon. Great Basin Nat. 44(4):563-571.



UINTA CHIPMUNK

Tamias umbrinus


State rank S3?

Element code AMAFB02190

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



State range

Recorded at high elevations in the Beartooth Mountains. May occur in the Absaroka Range and/or other areas around Yellowstone N.P. (Hoffmann et al. 1969, Flath 1984).

Habitat description

Subalpine spruce-fir forest, mountain meadows (Dice 1923, Hoffmann and Pattie 1968); talus slopes up into alpine (Pattie and Verbeek 1967).

Model assumptions & caveats

References


Dice, L.R. 1923. Mammal associations and habitats of the Flathead Lake Region, Montana. *Ecology* 4(3):247-260.

Flath, D. L. 1984. Vertebrate species of special interest or concern. Mammals, birds, reptiles, amphibians, fishes. Spec. Publ. Montana Department of Fish, Wildlife, and Parks, Helena. 76 pp.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.

Pattie, D. L. and N. A. M. Verbeek. 1967. Alpine mammals of the Beartooth Plateau. *Northwest Sci.* 41(3): 110-117.



BLACK-TAILED PRAIRIE DOG

Cynomys ludovicianus


State rank S3S4

Element code AMAFB06010

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



Predicted habitat: 10,318,759 ha, 27.10 % of state.

State range

Does not occur west of the Continental Divide. Distribution reduced because of eradication efforts (MT-NHP VCA).

Habitat description

Formerly very abundant in grasslands and sagebrush-grassland semi-desert (Hoffmann and Pattie 1968). An indicator of overgrazing (Knowles 1982).

Model assumptions & caveats

Model would better reflect species' distribution if prairie dog town layer were incorporated.

References

Apa, A. D., D. W. Uresk, and R. L. Linder. 1990. Black-tailed prairie dog populations one year after treatment with rodenticides. *Great Basin Nat.* 50:107-113.

Clippinger, N. W. 1989. Habitat suitability index models: black-tailed prairie dog. *U.S. Fish Wildl. Serv. Biol. Rep.* 82(10.156). 21 pp.

Flath, D. L. 1978. At home with the prairie dog. *Montana Outdoors.* 9(2):3-8.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Hoogland, J. L. 1995. The black-tailed prairie dog. Social life of a burrowing mammal. University of Chicago Press, Chicago. xiv. plus 557 pp.

Knowles, C. J. 1982. Habitat affinity, populations, and control of black-tailed prairie dogs on the Charles M. Russell National Wildlife Refuge. Ph.D. Thesis, Univ. Montana, Missoula. 171 pp.


Knowles, C. J. 1987. Reproductive ecology of black-tailed prairie dogs in Montana. *Great Basin Nat.* 47:202-206.

Knowles, C. J., C. J. Stoner, and S. P. Gieb. 1982. Selective use of black-tailed prairie dog towns by mountain plovers. *Condor* 84:71-74.

Licht, D. S. and K. D. Sanchez. 1993. Association of black-tailed prairie dog colonies with cattle point attractants in the northern Great Plains. *Great Basin Nat.* 53:385-389.

Linder, R.L. and C.N. Hillman. 1973. Proceedings of the black-footed ferret and prairie dog workshop. September 4-6, Rapid City, South Dakota. South Dakota State University. 208 pp.

Tileston, J.V., and R.R. Lechleitner. 1966. Some comparisons of the black-tailed and white-tailed prairie dogs in north-central Colorado. *Am. Midl. Nat.* 75: 292-316.



WHITE-TAILED PRAIRIE DOG

Cynomys leucurus


State rank S2

Element code AMAFB06020

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



State range

Peripheral; reaches northern limit in Carbon County, south-central Montana (Thompson 1982, MT-NHP VCA).

Habitat description

Salt sage (*Ivs axillaris*) and white sage (*Eurotia lanata*) (Hoffmann and Pattie 1968). Grassland and rangeland (Clark et al. 1971). Found along the Clarks Fork of the Yellowstone and its tributaries in arid, and atypical, habitat; elsewhere, the species is associated with mountain meadows (MT-NHP VCA).

Model assumptions & caveats

Model would better reflect species' distribution if prairie dog town layer were incorporated.

References

Clark, T. W. 1977. Ecology and ethology of the white-tailed prairie dog (*Cynomys leucurus*). 103 pp.

Clark, T. W., R. S. Hoffmann, and C. F. Nadler. 1971. *Cynomys leucurus*. Mamm. Species 7:1-4.


Flath, D. L. 1978. At home with the prairie dog. Montana Outdoors. 9(2):3-8.

Flath, D. L. 1979. Status of the white-tailed prairie dog in Montana. Proc. Mont. Acad. Sci. 38:63-67.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Linder, R.L. and C.N. Hillman. 1973. Proceedings of the black-footed ferret and prairie dog workshop. September 4-6, Rapid City, South Dakota. South Dakota State University. 208 pp.

Thompson, L.S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Bozeman: Montana Audubon Council. 24 pp.



GREAT BASIN POCKET MOUSE

Perognathus parvus


State rank S2S4

Element code AMAFD01070

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



State range

Known from Beaverhead, suspected in Madison County (Flath 1984). Probably limited to arid areas in southwestern Montana.

Habitat description

Sage, greasewood, saltbrush, and dry grassland with sandy soils, not more than 50% clay (Feldhamer 1979, Verts and Kirkland 1988). Big Sage of shrub-steppe with continuous cover (O'Farrell et al. 1975). Arid, semi-arid habitat (Hoffmann and Pattie 1968).

Model assumptions & caveats

Soil texture not included because scale of available statewide soils map was too coarse.

References

Feldhamer, G.A. 1979. Vegetative and edaphic features affecting the abundance and distribution of small mammals in southeast Oregon. *Great Basin Nat.* 39:207-218.

Flath, D. L. 1984. Vertebrate species of special interest or concern. Mammals, birds, reptiles, amphibians, fishes. Spec. Publ. Montana Department of Fish, Wildlife, and Parks, Helena. 76 pp.

Hayward, C.L. and M.L. Killpack. 1958. Distribution and variation of the Utah population of the great basin pocket mouse. *Great Basin Nat.* 18:26.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.


Kritzman, E.B. 1973. Ecological relationships of *Peromyscus maniculatus* & *Perognathus parvus* in eastern WA 1974. *J. Mammal.* 55:172-188.

Medin, D. E. and W. P. Clary. 1991. Small mammals of a beaver pond ecosystem and adjacent riparian habitat in Idaho. USDA, Forest Service, Res. Paper INT-445.

O'Farrell, T. P., R. J. Olson, R. O. Gilbert, and J. D. Hedlund. 1975. A population of Great Basin pocket mice (*Perognathus parvus*) in the shrub-steppe of south-central Washington. *Ecol. Monogr.* 45(1):1-28.

Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.

Verts, B. J., and G. L. Kirkland, Jr. 1988. *Perognathus parvus*. *Mamm. Species* 318:1-8.



HISPID POCKET MOUSE

Chaetodipus hispidus


State rank S1

Element code AMAFD05050

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



State range

One capture in Montana on a north-facing grassland slope in Carter County at 1,040m elevation (Pefaur and Hoffmann 1971). Has been recorded 20 miles south of Montana in Wyoming and in North Dakota near eastern Montana (Jones et al. 1983).

Habitat description

Prefers prairie areas with sparse or moderate vegetation; various dry grassland habitats. Has been found in irrigated cornfields. In underground burrow when inactive. Can occur in a variety of upland habitats. Usually loamy soil areas with some bare ground; not restricted to sandy soils like other pocket mice. Sometimes rocky prairie areas. Mid- and shortgrasses, shrubs, forbs, cactus, and yucca vegetation (Jones et al. 1983).


Model assumptions & caveats

References

Jones, J. K. Jr., D. M. Armstrong, R. S. Hoffmann and C. Jones. 1983. Mammals of the northern Great Plains. Univ. Neb. Press, Lincoln. 379 pp.

Paulson, D. D. 1988. *Chaetodipus hispidus*. Mamm. Species 320:1-4.

Pefaur, J. E. and R. S. Hoffmann. 1971. Merriam's shrew and hispid pocket mouse in Montana. Am. Midl. Nat. 86(1):247-248.



NORTHERN BOG LEMMING

Synaptomys borealis


State rank S2

Element code AMAFF17020

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



State range

Until recently, there were few locations known in Montana: several in Glacier National Park (Wright 1950, Weckwerth and Hawley 1962) and one in the Rattlesnake drainage north of Missoula (Adelman 1979). During 1992-93, 10 additional sites were found, with locations ranging from the northwestern corner of Montana south to just north of Lost Trail Pass in Beaverhead County and east to the Rocky Mountain Front (Reichel and Beckstrom 1993, 1994). The northern bog lemming may eventually be found to occur locally across much of western Montana.

Habitat description

Sedge-alder-willow bogs within or on edge of dense spruce-fir and lodgepole forest (Dice 1923, Hoffmann and Pattie 1968). Sphagnum moss mats important (Reichel and Beckstrom 1993).

Model assumptions & caveats

Marshes and bogs not well mapped.

References


Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.

Adelman, E. B. 1979. A survey of the nongame mammals in the Upper Rattlesnake Creek drainage of western Montana. M.S. thesis. University of Montana, Missoula. 129 pp.

Banfield, A. W. F. 1974. The mammals of Canada. University of Toronto Press, Toronto. Reprinted, 1981.

Clough, G. C., and J. J. Albright. 1987. Occurrence of the northern bog lemming, *Synaptomys borealis*, in the northeastern United States. *Can. Field-Naturalist* 101:611-613.

- Dice, L.R. 1923. Mammal associations and habitats of the Flathead Lake Region, Montana. *Ecology* 4(3):247-260.
- Groves, C. and E. Yensen. 1989. Rediscovery of the northern bog lemming (*Synaptomys borealis*) in Idaho. *Northw. Nat.* 70:14-15.
- Hall, E. R. 1981. *Mammals of North America*, 2nd edition. (2 vols.). John Wiley and Sons, New York.
- Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.
- Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.
- Jonkel, C. J. 1959. An ecological and physiological study of pine marten. M.S. thesis. Montana State Univ., Missoula. 81 pp.
- Laysen, E. F. and T. E. Burke. 1973. The northern bog lemming and its unique habitat in northeastern Washington. *Murrelet* 54:7-8.
- Reichel, J.D. and S.G. Beckstrom. 1993. Northern bog lemming survey: 1992. Unpublished report. Montana Natural Heritage Program, Helena, MT. 64 pp.
- Reichel, J.D. and S.G. Beckstrom. 1994. Northern bog lemming survey: 1993. Unpublished report. Montana Natural Heritage Program. Helena, MT. 87 pp.
- Weckwerth, R. P. and V. D. Hawley. 1962. Marten food habits and population fluctuations in Montana. *J. Wildl. Manage.* 26(1): 55-74.
- Wright, P.L. 1950. *Synaptomys borealis* from Glacier National Park, Montana. *J. Mammal.* 31(4): 460.



MEADOW JUMPING MOUSE

Zapus hudsonius


State rank S2S3

Element code AMAFH01010

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



Predicted habitat: 2,134,775 ha, 5.61 % of state.

State range

Several counties in eastern Montana (Flath 1984).

Habitat description

Grassland, meadows, and streamsides with thick and tall grass or forbs (Hoffmann and Pattie 1968, Pefaur and Hoffmann 1971, Matthews 1980).

Model assumptions & caveats

References

Douglass, R.J., and McNaughton, A.E.L. 1977. A recent record of the meadow jumping mouse, *Zapus hudsonius*, in the Northwest Territories. *Can. Field Nat.* 91: 96-97.

Flath, D. L. 1984. Vertebrate species of special interest or concern. Mammals, birds, reptiles, amphibians, fishes. *Spec. Publ. Montana Department of Fish, Wildlife, and Parks*, Helena. 76 pp.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. *Univ. Mont., Missoula*. 133 pp.


Hoyle, J. A. and R. Boonstra. 1986. Life history traits of the meadow jumping mouse, *Zapus hudsonius*, in southern Ontario. *Can. Field-Nat.* 100:537-544.

Matthews, W.L. 1980. Note: the meadow jumping mouse in southeastern Montana. *Prairie Nat.* 12(2):63-64.

Pefaur, J. E. and R. S. Hoffmann. 1971. Merriam's shrew and hispid pocket mouse in Montana. *Am. Midl. Nat.* 86(1):247-248.

Whitaker, J.O., and R. E. Wrigley. 1972. *Zapus hudsonius*. Mamm. Species No. 11. 7 pp.

Whitaker, J.O., and R. E. Wrigley. 1972. *Zapus hudsonius*. Mamm. Species No. 11. 7 pp.




GRAY WOLF

Canis lupus

State rank S1
Global rank G4
MTNHP status special concern

Element code AMAJA01030
Modeled by P. Thornton



State range

Resides in the Glacier N.P. area adjacent to Canada — present on east side, pack on west side. Also reports from northern Yellowstone NP and movements along Bitterroot divide. Occasional reports from other areas (Hoffmann et al. 1969, Northern Rocky Mountain Wolf Recovery Team 1980, Kaminski and Hansen 1984, Robbins 1986).

Habitat description

No particular habitat preference. Requires areas with low human population and high prey densities. In Minnesota and Wisconsin, usually found in areas with few roads, which increase human access and land uses incompatible with wolf presence (Thiel 1985, Mech et al. 1988, Mech 1989). Minimum of 10,000-13,000 sq km (with low road density) might be necessary to support a viable population (USFWS 1990); a single pack does not constitute a "minimum viable population." Young are born in an underground burrow that has been abandoned by another mammal or dug by wolf. In Minnesota, dens usually were not near territory boundaries, although they were used traditionally by most denning alpha females studied for more than 1 year; possibly the availability of a stable food supply source helped determine den location (Ciucci and Mech 1992). Remote montane and forest areas; uses valley bottoms and flood plains, and deer/elk winter ranges (Ream 1985, Ream et al. 1985, Ream 1986). Occasionally moves through atypical habitat (Hoffmann et al. 1969).

Model assumptions & caveats

References

Bangs, E. E., and S. H. Fritts. 1993. Reintroduction of gray wolves to Yellowstone National Park and central Idaho. *Endangered Species Tech. Bull.* 18(3):1, 18-20.


- Bergerud, A. T. and W. B. Ballard. 1988. Wolf predation on caribou: the Nelchina herd case history, a different interpretation. *J. Wildl. Manage.* 52:344-357.
- Bjorge, R. R., and J. R. Gunson. 1989. Wolf, *Canis lupus*, population characteristics and prey relationships near Simonette River, Alberta. *Can. Field-Nat.* 103:327-334.
- Carbyn, L. 1983. *Wolves in Canada and Alaska*. Ottawa. 135 pp.
- Ciucci, P., and L. D. Mech. 1992. Selection of wolf dens in relation to winter territories in northeastern Minnesota. *J. Mammal.* 73: 899-905.
- Cohn, J. P. 1990. Endangered wolf population increases. *BioScience* 40(9):628-632.
- Fritts, S.H. and L.D. Mech. 1981. Dynamics, movements, and feeding ecology of a newly protected wolf population in northwestern Minnesota. *Wildl. Monogr.* No. 80:1-79.
- Gese, E.M. and L.D. Mech. 1991. Dispersal of wolves (*Canis lupus*) in northeastern Minnesota, 1969-1989. *Can. J. Zool.* 69:2946-2955.
- Harrington, F. H. and P. C. Paquet. 1982. *Wolves of the world: perspectives of behavior, ecology, and conservation*. Noyes Publications, Park Ridge, New Jersey.
- Hoffmann, R. S. and D. L. Pattie. 1968. *A guide to Montana mammals: identification, habitat, distribution, and abundance*. Univ. Mont., Missoula. 133 pp.
- Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.
- Kaminski, T. and J. Hansen. 1984. *Wolves of central Idaho*. MT Coop. Wildl. Res. Unit, Missoula. 197 pp.
- Klinghammer, E. (ed). 1978. *The behavior and ecology of wolves*. Garland STPM Press, New York. 588 pp.
- Licht, D. S. and S. H. Fritts. 1994. Gray wolf (*Canis lupus*) occurrences in the Dakotas. *Am. Midl. Nat.* 132:74-81.
- Mech, L.D. 1974. *Canis lupus*. *Mamm. Species No.* 37. 6 pp.
- Mech, L. D. 1989. Wolf population survival in an area of high road density. *Am. Midl. Nat.* 121: 387-389.
- Mech, L.D., S.H. Fritts, G.L. Radde, and W.J. Paul. 1988. Wolf distribution and road density in Minnesota. *Wildl. Soc. Bull.* 16(1): 85-87.
- Murie, A. 1944. *The wolves of Mt. McKinley*. National Park Service, Fauna Ser. 5, 238 pp.
- Northern Rocky Mountain Wolf Recovery Team. 1980. *Northern Rocky Mountain wolf recovery plan interagency report*. 67 pp.
- Ream, R.R. 1985. *Wolf ecology project: annual report, July 1984 - June 1985*. MT Coop. Wildl. Research Unit. Univ. Montana, Missoula, MT 59812.
- Ream, R.R. 1986. Wolf movement in southern Alberta. *National Geographic Society Res. Rep.* Vol. 21:405-409.
- Ream, R.R., R.B. Harris, J. Smith, and D. Boyd. 1985. Movement patterns of a lone wolf (*Canis lupus*) in unoccupied wolf range, southeastern British Columbia. *Can. Field Nat.* 99(2):234-239.
- Robbins, J. 1986. Wolves across the border. *Nat. Hist.* 95(5): 6-15.

Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.

Thiel, R. P. 1985. Relationship between road densities and wolf habitat suitability in Wisconsin. *Am. Midl. Nat.* 113: 404-407.

U.S. Fish and Wildlife Service. 1990. Draft revised recovery plan for the eastern timber wolf. 93 pp.

U.S. Fish and Wildlife Service. 1990. Endangered and threatened species recovery program: report to Congress. 406 pp.




SWIFT FOX

Vulpes velox

State rank S1
Global rank G3
MTNHP status special concern

Element code AMAJA03030
Modeled by P. Thornton



State range

Once common and widespread on the prairie grasslands east of the Rocky Mountains, the species was believed to have been extirpated from Montana sometime between 1918 and 1978; however, recent records and sightings indicate that the species is recolonizing in southeastern and north-central Montana. These observations presumably represent dispersing individuals from populations outside of Montana, but a small resident population may actually exist in the north-central portion of the state north of Chinook and around Circle, Terry, and Wilsall (Giddings and Knowles 1995, Knowles et al. 1995).

Habitat description

Great Basin habitat types, prairie, badlands, shrub and shrub/grassland (Egoscue 1962, Hoffmann and Pattie 1968, Uresk and Sharps 1986, Allen 1987). Soils from clay-loam to sandy or gravelly loam; easy to dig (Scott-Brown et al. 1987).

Model assumptions & caveats

Soil texture not included because scale of available statewide soils map is too coarse.

References

- Allen, A.W. 1987. The relationship between habitat and furbearers. Pages 164-179 in M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. Wild Furbearer Management and Conservation in North America. Ontario Trappers Assn. and Ontario Ministry Nat. Res., Toronto, Ontario.
- Daneke, D., M. Sunquist, and S. Berwick. 1984. Notes on kit fox biology in Utah. Southwest. Nat. 29:361-362.
- Egoscue, H. J. 1962. Ecology and life history of the kit fox in Tooele County, Utah. Ecology 43(2):481-497.

Egoscue, H. J. 1979. *Vulpes velox*. Mamm. Species 122:1-5.

Floyd, Barry L. New Records of the Swift Fox (*vulpes velox*) in Wyoming. J. Mammal. V.62(3). pp.650-651.

Giddings, B. and C. J. Knowles. 1995. The current status of swift fox in Montana. Pp. 101-105 in: Allen, S. H., J. W. Hoagland, and E. D. Stukel, (eds.), Report of the Swift Fox Conservation Team. 170 pp.

Hines, Terrence D. 1991. Diet, Home Range, Movements, and Activity Periods of Swift Fox in Nebraska. Prairie Nat. 23(3):131-138.


Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Jones, J. K. Jr., D. M. Armstrong, R. S. Hoffmann and C. Jones. 1983. Mammals of the northern Great Plains. Univ. Neb. Press, Lincoln. 379 pp.

Knowles, C. J., P. R. Knowles, B. Giddings, and A. R. Dood. 1995. Status of the Swift fox in Montana. [document submitted for publication]. FaunaWest Wildlife Consultants, Boulder, MT, and MT Dept. of FWP, Helena, MT. 16 pp.

Scott-Brown, J.M., S. Herrero, and J. Reynolds. 1987. Swift fox. Pages 432-441 in M. Novak, J. A. Baker, M. E. Obbard, and B. Malloch, eds. Wild Furbearer Management and Conservation in North America. Ontario Trappers Assn. and Ontario Ministry Nat. Res., Toronto, Ontario, Canada.

Uresk, D. W., and J. C. Sharps. 1986. Denning habitat and diet of the swift fox in western South Dakota. Great Basin Nat. 46:249-253.



GRIZZLY OR BROWN BEAR

Ursus arctos


State rank S1S2

Element code AMAJB01020

Global rank G4

Modeled by P. Thornton

MTNHP status special concern



Predicted habitat: 6,923,086 ha, 18.18 % of state.

State range

Occurs in the Northern Continental Divide ecosystem and Yellowstone ecosystems, and also at low numbers in the Cabinet-Yaak ecosystem. Suspected in the Selway-Bitterroot ecosystem. Scattered individuals may occur in other mountainous areas of western Montana. Extinct in eastern Montana (Thompson 1982, MT-NHP VCA).

Habitat description

Now found mostly in arctic tundra, alpine tundra, and subalpine mountain forests. Once found in a wide variety of habitats including open prairie, brushlands, riparian woodlands, and semidesert scrub. Ranges widely at the landscape level. Most populations require huge areas of suitable habitat. Common only where food is abundant and concentrated (e.g., salmon runs, caribou calving grounds, berry patches, glacier lilies). Typically digs own hibernation den, usually on steep northern slope where snow accumulates. See LeFranc et al. (1987). Young are born in den in cave, crevice, hollow tree; hollow dug under rock, or similar site. Use of summit or ridge for mating (in May-June) reported for Banff National Park, Alberta, but not elsewhere (Hamer and Herrero 1990). In Spain, remnant deciduous forests and upland creek drainages were prime feeding areas (Clevenger et al. 1992). Meadows, seeps, riparian zones, mixed shrub fields, closed timber, open timber, sidehill parks, snow chutes, alpine slabrock. Varies between area, season, local populations and individuals (Chapman and Feldhamer 1982, Craighead et al. 1982, Servheen 1983, Aune 1984, Kasworm 1985).

Model assumptions & caveats

Distribution likely overestimated due to lack of road density data.

References

- Aune, K. 1984. Rocky Mountain Front grizzly bear monitoring and investigation. Montana Dept. Fish, Wildlife and Parks, Helena, 239 pp.
- Banci, V. 1991. Status report on the grizzly bear *Ursus arctos horribilis*. Committee on the Status of Endangered Wildlife in Canada. 171 pp.
- Blanchard, B. M., R. R. Knight, and D. J. Mattson. 1992. Distribution of Yellowstone grizzly bears during the 1980s. *Am. Midl. Nat.* 128:332-338.
- Chapman, J. A. and G. A. Feldhamer, eds. 1982. Wild mammals of North America: biology, management, and economics. Johns Hopkins Univ. Press. Baltimore, MD. 1147 pp.
- Clevenger, A. P., F. J. Purroy, and M. R. Pelton. 1992. Food habits of brown bears (*Ursus arctos*) in the Cantabrian Mountains, Spain. *J. Mammal.* 73: 415-421.
- Craighead, J. J., J. S. Sumner, and G. B. Scaggs. 1982. A definitive system for analysis of grizzly bear habitat and other wilderness resources. Wildlife-Wildlands Institute Monograph #1. University of Montana Foundation, Missoula. 279 pp.
- Darling, L. and R. Archibald, editors. 1990. Bears: their biology and management. Proc. Eighth Int. Conf. on Bear Res. and Manage., Victoria, British Columbia, Feb. 1989. 438 pp.
- Edge, W. D., C. L. Marcum and S. L. Olson-Edge. 1990. Distribution and grizzly bear (*Ursus arctos*) use of yellow sweetvetch (*Hedysarum sulphurescens*) in northwestern Montana and southeastern British Columbia. *Can. Field-Nat.* 104:435-438.
- Hamer, D. and S. Herrero. 1990. Courtship and use of mating areas by grizzly bears in the Front Ranges of Banff National Park, Alberta. *Can. J. Zool.* 68: 2695-2697.
- Hamer, D., S. Herrero and K. Brady. 1991. Food and habitat used by grizzly bears, *Ursus arctos*, along the Continental Divide in Waterton Lakes National Park, Alberta. *Can. Field-Nat.* 105:325-329.
- Hayward, G. D. 1989. Historical grizzly bear trends in Glacier National Park, Montana: a critique. *Wildl. Soc. Bull.* 17:195-197.
- Herrero (ed.). Bears: their biology and management. IUCN Publ. new ser., No. 23, 371 pp.
- Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.
- Kasworm, W. 1985. Cabinet Mountains grizzly bear study: 1984 annual progress report. Montana Department of Fish, Wildlife and Parks, Helena. 81 pp.
- LeFranc, M.N., Jr., M.B. Moss, K.A. Patnode, and W.C. Sugg III, eds. 1987. Grizzly bear compendium. Interagency Grizzly Bear Committee. iii + 540 pp.
- Martinka, C. J. and K. L. McArthur, (eds). 1980. Bears: their biology and management. Bear Biology Assoc. Conf. Ser. No. 3. Bear Biology Assoc., Boise, Idaho. 375 pp.
- Mattson, D. J. and D. G. Despain. 1985. Grizzly bear habitat component mapping handbook for the Yellowstone ecosystem. Interagency Grizzly Bear Study Team: National Park Service and U.S. Forest Service, May 1985. 37 pp.
- Mattson, D. J. and M. M. Reid. 1991. Conservation of the Yellowstone grizzly bear. *Conservation Biology* 5(3):364-372.
- Mattson, D. J., B. M. Blanchard and R. R. Knight. 1991. Food habits of Yellowstone grizzly bears, 1977-1987. *Can. J. Zool.* 69:1619-1629.


Mattson, D. J., C. M. Gillin, S. A. Benson and R. R. Knight. 1991. Bear feeding activity at alpine insect aggregation sites in the Yellowstone ecosystem. *Can. J. Zool.* 69:2430-2435.

Pattie, D. L. and N. A. M. Verbeek. 1967. Alpine mammals of the Beartooth Plateau. *Northwest Sci.* 41(3): 110-117.

Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.

Servheen, C. 1983. Grizzly bear food habits, movements and habitat selection in the Mission Mountains, Montana. *J. Wildl. Manage.* 47:1026-1035.

Thompson, L.S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Bozeman: Montana Audubon Council. 24 pp.



FISHER

Martes pennanti


State rank S2

Element code AMAJF01020

Global rank G5

Modeled by P. Thornton

MTNHP status special concern



Predicted habitat: 5,035,902 ha, 13.22 % of state.

State range

Apparently was previously extirpated in state. No documentation for 1920-1960. In 1959, 36 fishers from central British Columbia were released at three sites in western Montana; at least one transplant was successful and is apparently increasing (Weckwerth and Wright 1968). Recently recorded in southwestern Montana (Thompson 1982).

Habitat description

Prefers continuous canopy, dense old-growth conifer forest (Powell 1982, Allen 1987, Clark et al. 1989, Heinemeyer and Jones 1994). Also found at timberline (Aubry and Houston 1992). At least 70% canopy closure, fallen logs, stumps and seedlings, shrubs and herbaceous cover important (Powell 1982, Clark et al. 1989). Riparian areas especially important (Powell 1982, Heinemeyer and Jones 1994). Preys on snowshoe hares, voles, shrews, red squirrels, and birds (Raine 1987).

Model assumptions & caveats

Size class not available, so old-growth conditions are not well mapped. No information on coarse woody debris, stumps, and understory so suitable habitat is overestimated.

References


Allen, A. W. 1983. Habitat suitability index models: fisher. U.S. Fish and Wildl. Serv. FWS/OBS-82/10.45. 19 pp.

Allen, A.W. 1987. The relationship between habitat and furbearers. Pages 164-179 in M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. Wild Furbearer Management and Conservation in North America. Ontario Trappers Assn. and Ontario Ministry Nat. Res., Toronto, Ontario.

- Arthur, S. M. 1989. Habitat use and diet of fishers. *J. Wildl. Manage.* 53(3):680-688.
- Arthur, S. M. 1989. Home range characteristics of adult fishers. *J. Wildl. Manage.* 53(3):674-679.
- Arthur, S. M., W. B. Krohn, and J. R. Gilbert. 1989. Home range characteristics of adult fishers. *J. Wildl. Manage.* 53:674-679.
- Aubry, K.B. and D. B. Houston. 1992. Distribution and status of the fisher (*Martes pennanti*) in Washington. *Northwest Nat.* 73:69-79.
- Banci, V. 1994. Wolverine. Pages 99-127 in: Ruggiero, L. F., K. B. Aubry, S. W. Buskirk, L. J. Lyon, and W. J. Zielinski (eds.), *The scientific basis for conserving forest carnivores, American marten, fisher, lynx, and wolverine in the western United States.* U.S.D.A, Forest Serv. Rocky Mountain Forest and Range Exper. Stat. Gen. Tech. Report RM-254. 184 pp.
- Berg, W. E. 1982. Reintroduction of fisher, pine marten, and river otter. Pp. 159-173 in G. C. Sanderson (ed). *Midwest furbearer management. Proc. Symp. 43rd Midwest Fish and Wildlife Conference, Wichita, Kansas.*
- Brown, L. N. 1965. A fisher, *Martes pennanti*, in Sheridan County, Wyoming. *SW Nat.* 10(2):143.
- Buskirk, S. W. 1992. Conserving circumboreal forests for martens and fishers. *Cons. Biol.* 6:318-320.
- Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem. Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- Hash, H. S. and M. G. Hornocker. 1979. Range and habitat of male fishers in northwest Montana. *Idaho Coop. Wildl. Res. Unit. Univ. Idaho, Moscow.* 10 pp.
- Heinemeyer, K. S. and J. L. Jones 1994. Fisher biology and management: a literature review and adaptive management strategy. *USDA For. Serv. Northern Region, Missoula, MT.* 108 pp.
- Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. *Univ. Mont., Missoula.* 133 pp.
- Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.
- Maj, M. and E.O. Garten. 1994. Fisher, lynx, wolverine: Summary of distribution information. Pages 169-175 in Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon, and W.J. Zielinski (eds.). *The scientific basis for conserving forest carnivores, American marten, fisher, lynx, and wolverine in the western United States.* U.S.D.A, Forest Serv. Rocky Mountain Forest and Range Exper. Stat. Gen. Tech. Rep. RM-254. 184 pp.
- Powell, R.A. 1982. *The fisher: life history, ecology, and behavior.* Univ. Minnesota Press. xvi + 217 pp.
- Raine, R.M. 1983. Winter habitat use and responses to snow cover of fisher (*Martes pennanti*) and marten (*Martes americana*) in southeastern Manitoba. *Can. J. Zool.* 61:25-34.
- Raine, R.M. 1987. Winter food habits and foraging behavior of fishers (*Martes pennanti*) and martens (*Martes americana*) in southeastern Manitoba. *Can. J. Zool.* 65: 745-747.
- Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.
- Thomasma, L. E., T. D. Drummer, and R. O. Peterson. 1991. Testing the habitat suitability index model for the fisher. *Wildl. Soc. Bull.* 19:291-297.

Thompson, L.S. 1982. Distribution of Montana amphibians, reptiles, and mammals. Bozeman: Montana Audubon Council. 24 pp.

Weckwerth, R. P. and P. L. Wright. 1968. Results of transplanting fishers in Montana. J. Wildl. Manag. 32(4):977-981.



BLACK-FOOTED FERRET

Mustela nigripes


State rank SH#

Element code AMAJF02040

Global rank G1

Modeled by P. Thornton

MTNHP status special concern



State range

Historically found throughout eastern Montana. Reintroduced at UL Bend in Phillips County in fall 1994 with continued releases through 1997. Also released on Fort Belknap Reservation in fall 1997. Extirpated elsewhere in the state. Could possibly occur in several counties of southeastern Montana: Carter, Custer, and Rosebud (MT NHP VCA).

Habitat description

Limited to open habitat, the same habitat used by prairie dogs: grasslands, steppe, and shrub steppe. Occupies underground burrows made by prairie dogs; either black-tailed or white-tailed prairie dog colonies offer acceptable habitat. Young are born in underground burrows made by prairie dogs. It has been estimated that about 40-60 ha of a prairie dog colony is needed to support one ferret, but little is known about what truly constitutes ferret habitat, and reintroductions are providing new insights every year (R. Matchett, pers. comm.). See Biggins et al. (*in Oldemeyer et al. 1993*) for information on evaluating areas as potential ferret habitat. Such habitat factors include size of prairie dog complex, prairie dog population density, spatial arrangement of prairie dog colonies, potential for disease in prairie dogs and ferrets, potential for prairie dog expansion, abundance of predators, future resource conflicts and ownership stability, and public and landowner attitudes. Only large prairie dog complexes (several thousand acres of closely-spaced colonies) can support a breeding ferret population.

Model assumptions & caveats

Model would be greatly improved by inclusion of prairie dog town layer.

References

Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.


Biggins, D. and M.H. Schroeder. 1988. Historical and present status of the black-footed ferret. Pages 93-97 in G. L. Shenbeck and R. Lifkin, eds. Proc. eighth Great Plains wildlife damage control workshop. U.S. For. Serv. Gen. Tech. Rep. RM-154.

Hoffmann, R. S. and D. L. Pattie. 1968. A guide to Montana mammals: identification, habitat, distribution, and abundance. Univ. Mont., Missoula. 133 pp.

Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.

Linder, R.L. and C.N. Hillman. 1973. Proceedings of the black-footed ferret and prairie dog workshop. September 4-6, Rapid City, South Dakota. South Dakota State University. 208 pp.

Oldemeyer, J.L., D.E. Biggins, B.J. Miller, and R. Crete, editors. 1993. Proc. of the symposium on the management of prairie dog complexes for the reintroduction of the black-footed ferret. U.S. Fish Wildl. Serv. Biol. Rep., No. 13. 96 pp.




WOLVERINE

Gulo gulo

State rank S2
Global rank G4
MTNHP status special concern

Element code AMAJF03010
Modeled by P. Thornton



State range

Nearly extinct from 1920-1940, with increasing numbers and range since. "Recovery" originated in northwestern Montana-Glacier Park area and spread from there (Newby and Wright 1955, Newby and McDougal 1964). Current distribution primarily from western Montana through Greater Yellowstone area (Thompson 1982). May exhibit dispersal movements in and through atypical habitat (MT-NHP VCA).

Habitat description

Found in mature and old-growth fir, pine and larch forests (Hornocker and Hash 1981, Banci and Harestad 1990, Reichel et al. 1992). Also in alpine shrub, talus slopes, and riparian cottonwoods (Banci and Harestad 1990, Reichel et al. 1992). Requires large areas of habitat with little or no human disturbance (Hash 1987).

Model assumptions & caveats

Old-growth conditions are not well mapped.

References

Banci, V. 1986. The wolverine in the Yukon: myths and management. *Discovery* 15:134-137.

Banci, V. 1994. Wolverine. Pages 99-127 in: Ruggiero, L. F., K. B. Aubry, S. W. Buskirk, L. J. Lyon, and W. J. Zielinski (eds.), *The scientific basis for conserving forest carnivores, American marten, fisher, lynx, and wolverine in the western United States*. U.S.D.A, Forest Serv. Rocky Mountain Forest and Range Exper. Stat. Gen. Tech. Report RM-254. 184 pp.

Banci, V. and A. S. Harestad. 1990. Home range and habitat use of wolverines in Yukon, Canada. *Holarc. Ecol.* 13:195-200.

Butts, T.W. 1992. Wolverine (*Gulo gulo*) biology and management: a literature review and annotated bibliography. USDA For. Serv. N. Reg. Threatened, Endangered, and Sensitive Prog. 106 pp. + appendix.

Gardner, C. L., W. B. Ballard and R. H. Jessup. 1986. Long distance movements by an adult wolverine. *J. Mammal.* 67:603.

Groves, C. R. 1988. Distribution of the wolverine in Idaho as determined by mail questionnaire. *Northw. Sci.* 62:181-185.

Hash, H. S. 1987. Wolverine. Pages 574-585 in M. Novak, J. Baker, M. Obbard and B. Malloch, eds. *Wild furbearer management and conservation in North America*. Ministry Nat. Res., Toronto, Ontario.

Hoak, J., J. Weaver and T. Clark. 1980. Wolverines in western Wyoming. *Northwest Sci.* 56(3):159-161.

Hoffmann, R. S. and D. L. Pattie. 1968. *A guide to Montana mammals: identification, habitat, distribution, and abundance*. Univ. Mont., Missoula. 133 pp.

Hornocker, M. G. and H. S. Hash. 1981. Ecology of the wolverine in northwestern Montana. *Can. J. Zool.* 59:1286-1301.

Johnson, R. E. 1977. An historical analysis of wolverine abundance and distribution in Washington. *Murrelet* 58:13-16.

Maj, M. and E.O. Garten. 1994. Fisher, lynx, wolverine: Summary of distribution information. Pages 169-175 in Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon, and W.J. Zielinski (eds.). *The scientific basis for conserving forest carnivores, American marten, fisher, lynx, and wolverine in the western United States*. U.S.D.A, Forest Serv. Rocky Mountain Forest and Range Exper. Stat. Gen. Tech. Rep. RM-254. 184 pp.

Newby, F.E. and J.J. McDougal. 1964. Range extension of the wolverine in Montana. *J. Mammal.* 45: 485-487.


Newby, F.E. and P.L. Wright. 1955. Distribution and status of the wolverine in Montana. *J. Mammal.* 36: 248-253.

Reichel, J. D., D. L. Genter and E. Atkinson. 1992. *Sensitive animal species in the Elkhorn and Big Belt Mountains of the Helena National Forest*. Montana Natural Heritage Program. Helena, MT. 102 pp.

Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.

Thompson, L.S. 1982. *Distribution of Montana amphibians, reptiles, and mammals*. Bozeman: Montana Audubon Council. 24 pp.

van Zyll de Jong, C.G. 1975. The distribution and abundance of the wolverine (*Gulo gulo*) in Canada. *Can. Field-Nat.* 89(4):431-437.



WESTERN SPOTTED SKUNK

Spilogale gracilis


State rank SU

Element code AMAJF05020

Global rank G5

Modeled by P. Thornton

MTNHP status watch list



Predicted habitat: 1,370,006 ha, 3.60 % of state.

State range

Occurs in southwestern Montana (Thompson 1982, MT-NHP VCA).

Habitat description

Rocky brush/grassland, grassland, riparian areas, and forest/shrub ecotones (Verts 1967, Hoffmann and Pattie 1968, Chapman and Feldhamer 1982, Allen 1987)

Model assumptions & caveats

References

Allen, A.W. 1987. The relationship between habitat and furbearers. Pages 164-179 in M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. *Wild Furbearer Management and Conservation in North America*. Ontario Trappers Assn. and Ontario Ministry Nat. Res., Toronto, Ontario.


Chapman, J. A. and G. A. Feldhamer, eds. 1982. *Wild mammals of North America: biology, management, and economics*. Johns Hopkins Univ. Press. Baltimore, MD. 1147 pp.

Hoffmann, R. S. and D. L. Pattie. 1968. *A guide to Montana mammals: identification, habitat, distribution, and abundance*. Univ. Mont., Missoula. 133 pp.

Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.

Thompson, L.S. 1982. *Distribution of Montana amphibians, reptiles, and mammals*. Bozeman: Montana Audubon Council. 24 pp.

Verts, B.J. 1967. *The biology of the striped skunk*. Univ. Illinois Press, Urbana. vii+218 pp.




LYNX

Lynx canadensis

State rank S2
Global rank G5
MTNHP status special concern

Element code AMAJH03010
Modeled by P. Thornton



Predicted habitat: 6,045,562 ha, 15.88 % of state.

State range

Resident in western Montana (Thompson 1982). Appearances in eastern Montana may be residents or movements from Canada, possibly a response to periodic shortages of snowshoe hares (MT-NHP VCA).

Habitat description

Dependent on snowshoe hares; found in same habitats (Reichel et al. 1992). Dense, mature old-growth lodgepole pine, Douglas-fir, Engelmann spruce and subalpine fir forests (Koehler et al. 1979, Chapman and Feldhamer 1982, Koehler 1990, Reichel et al. 1992). Willow flats (Nellis in Clark et al. 1989). Above 4000' hunt in dense, early successional stage lodgepole pine forest, den in mature lodgepole pine and spruce-fir forests (Koehler and Brittell 1990). Well-developed understory important (Allen 1987).

Model assumptions & caveats

Understory conditions not mapped, therefore suitable habitat may be overestimated.

References

Abstracts from Montana Rare Animal Meeting. 1992. [November 5-6, 1992]. Lewistown, MT. 20 pp.

Allen, A.W. 1987. The relationship between habitat and furbearers. Pages 164-179 in M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. Wild Furbearer Management and Conservation in North America. Ontario Trappers Assn. and Ontario Ministry Nat. Res., Toronto, Ontario.

Banci, V. 1994. Wolverine. Pages 99-127 in: Ruggiero, L. F., K. B. Aubry, S. W. Buskirk, L. J. Lyon, and W. J. Zielinski (eds.), The scientific basis for conserving forest carnivores, American marten, fisher, lynx, and wolverine in the western United States. U.S.D.A, Forest Serv. Rocky Mountain Forest and Range Exper. Stat. Gen. Tech. Report RM-254. 184 pp.

- Barash, D. P. 1971. Cooperative hunting in the lynx. *J. Mammal.* 52:480.
- Brainerd, S. M. 1985. Reproductive ecology of bobcats and lynx in western Montana. M.S. thesis. University of Montana, Missoula. 85 pp.
- Brand, C. J., and L. B. Keith. 1979. Lynx demography during a snowshoe hare decline in Alberta. *J. Wildl. Manage.* 43:827-849.
- Brand, C. J., L. B. Keith and C. A. Fischer. 1976. Lynx responses to changing snowshoe hare densities in central Alberta. *J. Wildl. Manage.* 40(3):416-428.
- Butts, T.W. 1992. Lynx (*FELIS LYNX*) biology and management: a literature review and annotated bibliography. USDA Forest Service, Northern Region Threatened, Endangered, and Sensitive Program, Missoula, Mont. 114 pp. + appendix.
- Chapman, J. A. and G. A. Feldhamer, eds. 1982. *Wild mammals of North America: biology, management, and economics.* Johns Hopkins Univ. Press. Baltimore, MD. 1147 pp.
- Clark, T.W., A.H. Harvey, R.D. Dorn, D.L. Genter, and C. Groves, eds. 1989. *Rare, sensitive, and threatened species of the Greater Yellowstone Ecosystem.* Northern Rockies Conservation Cooperative, Montana Natural Heritage Program, The Nature Conservancy, and Mountain West Environmental Services. 153 pp.
- Hash, H. 1990. Montana lynx population status and considerations (1990). Unpubl. Rep. Mont. Dept. Fish, Wildl. Parks, Helena. 13 pp.
- Hoffmann, R. S. and D. L. Pattie. 1968. *A guide to Montana mammals: identification, habitat, distribution, and abundance.* Univ. Mont., Missoula. 133 pp.
- Hoffmann, R.S., P.L. Wright, and F.E. Newby. 1969. Distribution of some mammals in Montana. I. Mammals other than bats. *J. Mammal.* 50(3): 579-604.
- Jones, J. K. Jr., D. M. Armstrong, R. S. Hoffmann and C. Jones. 1983. *Mammals of the northern Great Plains.* Univ. Neb. Press, Lincoln. 379 pp.
- Koehler, G.M. 1990. Population and habitat characteristics of lynx and snowshoe hares in north central Washington. *Can. J. Zool.* 68:845-851.
- Koehler, G.M. and J.D. Brittell, 1990. Managing spruce-fir habitat for lynx and snowshoe hares. *J. For.* 88: 10-14.
- Koehler, G. M., M. G. Hornocker, and H. S. Hash. 1979. Lynx movements and habitat use in Montana. *Can. Field-Nat.* 93:441-442.
- Maj, M. and E.O. Garten. 1994. Fisher, lynx, wolverine: Summary of distribution information. Pages 169-175 in Ruggiero, L.F., K.B. Aubry, S.W. Buskirk, L.J. Lyon, and W.J. Zielinski (eds.). *The scientific basis for conserving forest carnivores, American marten, fisher, lynx, and wolverine in the western United States.* U.S.D.A, Forest Serv. Rocky Mountain Forest and Range Exper. Stat. Gen. Tech. Rep. RM-254. 184 pp.
- Reichel, J. D., D. L. Genter and E. Atkinson. 1992. Sensitive animal species in the Elkhorn and Big Belt Mountains of the Helena National Forest. Montana Natural Heritage Program. Helena, MT. 102 pp.
- Rust, H. J. 1946. Mammals of northern Idaho. *J. Mammal.* 27(4): 308-327.
- Thompson, L.S. 1982. *Distribution of Montana amphibians, reptiles, and mammals.* Bozeman: Montana Audubon Council. 24 pp.
- Tumilson, R. 1987. *Felis lynx.* *Mamm. Species* 269:1-8.

